

# Comparing the effects of different exercise sequences on shoulder recovery in male racket sport players

<b>Submission date</b> 27/03/2026	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 30/03/2026	<b>Overall study status</b> Completed	<input checked="" type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 30/03/2026	<b>Condition category</b> Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Racket sports like tennis and badminton place high mechanical demands on the shoulder, often causing discomfort and injury. Recent rehabilitation approaches suggest that whole-body coordination, especially from the pelvis and core, is vital for shoulder recovery. This study aims to find the most effective sequence of exercises by comparing two rehabilitation programs: one starting with pelvic-lumbar (core) coordination exercises followed by shoulder stabilization, and the other performing the reverse order.

### Who can participate?

Adult male recreational racket-sport participants who have experienced persistent shoulder discomfort.

### What does the study involve?

Participants are randomly divided into two groups. Both groups complete the same two 20-minute exercise sessions (a pelvic-lumbar core program and a shoulder stabilization program) but in opposite orders. The two sessions are separated by a 24-hour rest period. Shoulder movement and coordination are evaluated using a smartphone camera (motion analysis) at three time points: 5 minutes before the first program, 5 minutes after the first program, and 5 minutes after the second program.

### What are the possible benefits and risks of participating?

The possible benefits include improvements in shoulder mobility, function, and whole-body coordination during movements. The risks are minimal, primarily involving temporary mild muscle fatigue or soreness typical of standard exercise rehabilitation programs.

### Where is the study run from?

The study is run from the Republic of Korea, involving researchers from Seoul National University and Ajou University, with ethical approval from Jeonbuk National University.

When is the study starting and how long is it expected to run for?  
March 2025 to July 2025.

Who is funding the study?  
Investigator initiated and funded.

Who is the main contact?  
Dr. Donghwi Suh, donghwisuh@hanmail.net

## Contact information

### Type(s)

Public, Scientific, Principal investigator

### Contact name

Dr Donghwi Suh

### Contact details

Department of Practical Physical Education, Daegu Catholic University, 13-13 Hayang-ro, Hayang-eup, Gyeongsan-si, Gyeongsangbuk-do  
Gyeongsan-si  
Korea, South  
38430  
82-10-4131-9808  
donghwisuh@hanmail.net

## Additional identifiers

## Study information

### Scientific Title

The effects of a lumbar-shoulder exercise rehabilitation sequence compared to a shoulder-lumbar sequence on shoulder joint function recovery in male recreational racket sport participants with shoulder discomfort

### Acronym

CSAT

### Study objectives

To compare the effects of two exercise rehabilitation sequences (pelvic-lumbar coordination training followed by shoulder stabilization, and the reverse order) on shoulder joint function and coordination recovery in male recreational racket-sport participants.

### Ethics approval required

Ethics approval required

### Ethics approval(s)

approved 05/03/2025, Institutional Review Board of Jeonbuk National University (567, Baekje-daero, Deokjin-gu, Jeonju-si, Jeollabuk-do, 54896, Republic of Korea, 3rd Floor, Jeonbuk National University Main Building, Jeonju, 54896, Korea, South; +82-063-270-4889; ahnth2@jbnu.ac.kr), ref: 2025-02-021-001

## **Primary study design**

Interventional

## **Allocation**

Randomized controlled trial

## **Masking**

Open (masking not used)

## **Control**

Active

## **Assignment**

Crossover

## **Purpose**

Treatment

## **Study type(s)**

## **Health condition(s) or problem(s) studied**

Shoulder discomfort

## **Interventions**

Participants were randomly assigned to the groups using a computer-generated random number sequence. The allocation sequence was concealed and implemented using sealed opaque envelopes.

LS Group: Completes a 20-minute lumbar-focused program (Core Stabilization Adjustment Training) first, followed by a 24-hour rest, and then a 20-minute shoulder-focused stabilization program.

SL Group: Completes a 20-minute shoulder-focused stabilization program first, followed by a 24-hour rest, and then a 20-minute lumbar-focused program (Core Stabilization Adjustment Training). The total duration of the intervention for each participant spans two days, with testing conducted 5 minutes before and after each session.

## **Intervention Type**

Behavioural

## **Primary outcome(s)**

1. Dominant and non-dominant shoulder Range of Motion (ROM) during discrete and continuous movements measured using a video-based motion analysis (calculating joint angles using MediaPipe Pose Landmarker to extract 3D joint coordinates) at 5 minutes before the first program (pre-test), 5 minutes after the first program (post-test 1), and 5 minutes after the second program (post-test 2)

## **Key secondary outcome(s)**

1. Dynamical degrees of freedom (DDOF) during discrete and continuous movements measured using a Principal Component Analysis (PCA) of 3D joint angles extracted via video-based motion analysis (MediaPipe Pose) at 5 minutes before the first program, 5 minutes after the first program, and 5 minutes after the second program

2. Biomechanical degrees of freedom (BDOF) during discrete and continuous movements measured using the total number of unique joint motions across significant principal components calculated from video-based motion analysis at 5 minutes before the first program, 5 minutes after the first program, and 5 minutes after the second program

## **Completion date**

31/07/2025

## **Eligibility**

### **Key inclusion criteria**

1. Adult male recreational racket-sport participants.
2. Individuals who have received medical clearance to engage in shoulder rehabilitation exercises following the onset of shoulder pain.
3. Individuals reporting persistent shoulder discomfort lasting at least three months prior to study participation.

### **Healthy volunteers allowed**

No

### **Age group**

Adult

### **Lower age limit**

32 years

### **Upper age limit**

44 years

### **Sex**

Male

### **Total final enrolment**

32

### **Key exclusion criteria**

1. Individuals with neurological disorders.
2. Individuals with biomechanical alterations resulting from pathological conditions.

### **Date of first enrolment**

05/03/2025

### **Date of final enrolment**

31/07/2025

## Locations

### Countries of recruitment

Korea, South

## Sponsor information

### Organisation

Daegu Catholic University

### ROR

<https://ror.org/04fxknd68>

## Funder(s)

### Funder type

### Funder Name

Catholic University of Daegu

### Alternative Name(s)

Daegu Catholic University, , DCU, CU

### Funding Body Type

Private sector organisation

### Funding Body Subtype

Universities (academic only)

### Location

Korea, South

## Results and Publications

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not expected to be made available

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
-------------	---------	--------------	------------	----------------	-----------------

[Statistical Analysis Plan](#)

30/03/2026

No

No