

A two-arm pragmatic randomised control trial comparing the clinical and cost-effectiveness of serratus anterior plane block (SAP) in reducing the rate of respiratory infections 5 days post randomisation in patients with multiple rib fractures (MRF) when compared with the usual care in the NHS trauma setting

Submission date 13/10/2022	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 24/10/2022	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 28/02/2025	Condition category Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

People admitted to the hospital with multiple rib fractures often experience severe pain that reduces their coughing or movement. This leads to secretions building up in the lungs. As a result, it increases the risk of developing severe lung and blood infections, which delay recovery and hospital discharge, reduce quality of life after discharge and reduce the rate of survival. As part of the usual care, patients usually receive physiotherapy, deep breathing exercises (incentive spirometry) and rib fixation. Doctors would prescribe opiate-based pain relief to help patients to manage their pain, however this is often ineffective and comes with undesirable side effects.

ERASER aims to find out whether a regional pain relief technique, SAP (Serratus Anterior Plane block), used in addition with usual care, reduces the chance of developing lung infection, for people with multiple rib fractures. SAP is a technique where pain relief medication is infused into the chest at a regular interval, through a small tube (catheter) that is inserted from the side of the body under the guidance of ultrasound. This provides pain relief to the fracture area.

Who can participate?

Patients aged 16 years or older, with multiple rib fractures

What does the study involve?

Whilst in hospital, participants will be asked to rate the pain they are experiencing at regular intervals. The researchers will gather further data needed for this trial from the medical notes. A small blood sample will be taken for use in future research. After discharged from hospital, the

researchers will give participants a phone call at 1 and 3 months from when they joined the trial, to see the progress of recovery.

What are the possible benefits and risks of participating?

While there is no direct benefit or financial incentives to participants that take part in this trial, the information provided by participants may help in the long-term, to improve and shape future care for patients with multiple rib fractures.

SAP has been used across the NHS and has been shown to be relatively safe. However, as with most medical treatments, there are some uncommon risks associated with it, this includes possible infection or bleeding at to the site of catheter insertion, arrhythmias and nerve injury. The research team will closely monitor your health. If you have any concerns during your time in the trial, please do not hesitate to talk to the clinical care team or researcher team. There will be an independent safety committee that will review the trial data anonymously at regular intervals and on-demand where necessary to ensure that the trial is safe to continue.

Where is the study run from?

University of Birmingham (UK)

When is the study starting and how long is it expected to run for?

August 2021 to July 2026

Who is funding the study?

National Institute for Health and Care Research (NIHR) (UK).

Who is the main contact?

Dr Naveed Saeed, n.saeed@bham.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Naveed Saeed

Contact details

University of Birmingham
Birmingham Clinical Trials Unit (BCTU)
Edgbaston
Birmingham
United Kingdom
B15 2TT
+44 121 415 8445
n.saeed@bham.ac.uk

Type(s)

Principal investigator

Contact name

Prof Tonny Veenith

ORCID ID

<https://orcid.org/0000-0002-4125-8804>

Contact details

University of Birmingham
Institute of Inflammation and Ageing
Edgbaston
Birmingham
United Kingdom
B15 2TT

-
tonny.veenith1@nhs.net

Additional identifiers**Clinical Trials Information System (CTIS)**

Nil known

Integrated Research Application System (IRAS)

307628

Protocol serial number

CPMS 53786, NIHR130632, IRAS 307628

Study information**Scientific Title**

A pragmatic randomised controlled trial evaluating the clinical and cost-effectiveness of serratus anterior plane block with catheter insertion compared to usual care in patients with multiple rib fractures

Acronym

ERASER

Study objectives

SAP is more effective than usual care for the treatment of pain after multiple rib fractures (MRF) through improving clinical outcomes and cost-effectiveness

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 11/11/2022, London - Harrow Research Ethics Committee (Level 3, Block B, Whitefriars, Lewins Mead, Bristol, BS1 2NT, UK; +44 207 104 8246; harrow.rec@hra.nhs.uk), ref: 22/LO/0607

Study design

Interventional randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Multiple rib fractures

Interventions

Participant will be asked to rate their pain on a numerical scale and donate a blood sample for future research use.

Researchers will enter the baseline data in the secure online database and the computer will randomly assign the participant to receive SAP with usual care or usual care (at 1:1 ratio). A minimisation algorithm (built into the database) will be used to ensure balance in the intervention allocation over the following variables: Gender, Age, Centre, Injury Severity Score, Pulmonary contusion, Anatomical location of fracture, Presence of flail segment, and a random element (to avoid the possibility of allocation becoming predictable).

Due to the urgent nature of the condition, randomisation and intervention should occur on the same or next day after consent has been taken.

Participants will be asked to rate their pain on a numerical scale 4x a day on day 1-5, day 14 and on discharge from hospital.

Researchers will continue to collect follow up data from the medical notes until participant is discharged from the hospital. At month 1 and 3 from randomisation, the researcher will give a telephone call to the participant to monitor their progress of recovery. They will be asked to rate their quality of life (EQ-5D-5L) and confirm if they have used any community care resources for the same injury. They will also be asked to rate their pain (MRC dyspnoea scale, Brief Pain Index, McGill Pain Questionnaire).

Intervention Type

Procedure/Surgery

Primary outcome(s)

New diagnosis of pneumonia, as agreed by a blinded end point assessment committee, five days after randomisation

Key secondary outcome(s)

1. Pain assessed using a numerical rating scale (0-10), recorded every 4 hours, day 1 to 5, day 14 and before discharge, and by the patient-reported BPI at one and three-months post-randomisation.
2. Ventilatory function measured by incentive spirometry and peak expiratory flow twice a day via the daily assessment log until discharge
3. Number of days ventilated and the requirement for invasive/non-invasive methods
4. 30 day and 3 month mortality
5. Complications and safety data associated with SAP
6. Opiate consumption such as morphine and related compounds (daily until discharge and at 1 and 3 months post randomisation)

7. Length of stay in hospital and critical care level two/three facility
8. Hospital re-admission within 30 days of discharge
9. Patient-Reported Outcome Measures (PROMs) at 1 and 3 months:
 - 9.1. Quality of life: EQ-5D-5L
 - 9.2. Medical Research Council (MRC) dyspnoea scale at 3 months only
 - 9.3. Brief Pain Index (BPI)
 - 9.4. McGill Pain Questionnaire (SF-MPQ-2)

Completion date

31/07/2026

Eligibility

Key inclusion criteria

1. Patients ≥ 16 years with unilateral or bilateral ≥ 3 rib fractures following blunt chest trauma
2. Injury occurred ≤ 72 hours of hospital admission

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

16 years

Sex

All

Key exclusion criteria

1. Severe traumatic brain injury with a predicted ventilatory requirement >7 days
2. Acute quadriplegia
3. Spinal fracture precluding mobilisation
4. Penetrating trauma or open rib fractures
5. Upper airway injury requiring intubation and mechanical ventilation with an expected dependency of more than 5 days (e.g. tracheal disruption)
6. Any chest wall injuries that preclude a catheter insertion
7. Not anticipated to survive ≥ 48 hours
8. Contamination or infection at site of potential SAP insertion
9. Any contraindication for the treating clinician for SAP block catheter insertion
10. Patients of childbearing age who have tested positive for pregnancy

Date of first enrolment

01/02/2023

Date of final enrolment

01/10/2025

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Queen Elizabeth Hospital Birmingham

University Hospitals Birmingham NHS Foundation Trust

Mindelsohn Way

Birmingham

United Kingdom

B15 2TH

Study participating centre

Walsgrave General Hospital

Coventry and Warwickshire NHS Trust

Clifford Bridge Road

Coventry

United Kingdom

CV2 2DX

Study participating centre

Queen's Medical Centre

Nottingham University Hospitals NHS Trust

Derby Road

Nottingham

United Kingdom

NG7 2UH

Study participating centre

Royal Stoke University Hospital

University Hospitals of North Midlands NHS Trust

Newcastle Road

Stoke-on-Trent

United Kingdom

ST4 6QG

Sponsor information

Organisation

University of Birmingham

ROR

<https://ror.org/03angcq70>

Funder(s)

Funder type

Government

Funder Name

NIHR Evaluation, Trials and Studies Co-ordinating Centre (NETSCC)

Results and Publications

Individual participant data (IPD) sharing plan

All data generated or analysed during this study will be included in the subsequent results publication

IPD sharing plan summary

Published as a supplement to the results publication

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes