

ODD SOCKS - Outcomes of Displaced Distal tibial fractures - Surgery Or Casts in Kids

Submission date 05/02/2024	Recruitment status Recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 13/02/2024	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 24/07/2025	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Broken ankles in children often involve the area from which the bone grows – the growth plate. Following growth plate injuries, the growth of the main shin bone in the lower leg (the tibia) can be altered permanently, which can cause the bone to not grow at all, or to grow wonky. Altered growth may affect how well the leg works. The younger the child at the time of injury (i.e. the more they have to grow), the worse the problem may become once the child has fully grown. There are different ways to treat this injury, but it is currently unclear whether one type of treatment is better than another. Some doctors believe that children with growth plate injuries need surgery to reset their bones to ensure that the growth plate is restored to its original position. They believe that this will lower the chance of abnormal growth. However, other doctors believe that attempting to reset the bones to restore the growth plate with surgery could bring about further damage. These doctors recommend the bones be treated in a plaster cast, without surgery to reset the bones. This study aims to look at whether when children aged between 8 and 15 years old break their ankles, surgery to reset the bones leads to better function than letting the bones heal using a plaster cast without resetting the bones. At the end of the study, the information will be combined about all the children that took part. This will help everyone to understand which treatment is best. To make sure people learn about the best treatment, the doctors who help with this study will talk to other doctors and other people in the NHS who write national guidelines. Patient co-investigators will help deliver the message to families and will be invited to share their experience of the study with medical professionals.

Who can participate?

Children aged between 8 and 15 years old who have a fracture through the growth plate at the bottom of the shin bone, where the bone ends have moved apart from each other.

What does the study involve?

In the study, half the children and young people will have their broken bones treated with surgery, whilst the other half will have a plaster cast with no surgery. Those who agree to join the study, with the support of their families, will be split fairly into two groups, using a research process called 'randomisation'. Children will be assigned one of two treatments:

1. Surgical reduction – the children in this group will have an anaesthetic or be sedated so their bones can be reset in theatre, and a plaster cast put on their leg. Sometimes, if the doctor thinks

it necessary, wires, screws or a plate and screws will be inserted to hold the broken bones in position.

2. Conservative treatment – the children in this group will not have the bones reset in position, they will receive a plaster cast for support to allow the bones to heal naturally. The plaster casts will stay on for around 4-6 weeks for both treatments.

All children will be followed up for two years to keep track of their function, and the length and appearance of the leg. They will be asked about pain, whether they needed any more surgery, school attendance, complications, the number of hospital visits, their quality of life and satisfaction with treatment. Follow-up will occur at 6 weeks, 3, 6, 12 & 24 months. Participant follow-up will be organised by the University of Oxford – either electronically by email or text message or by telephone

What are the possible benefits and risks of participating?

Each of the two routinely used treatments has potential advantages and disadvantages.

1. Resting the leg in a plaster cast for up to 6 weeks, to allow it to heal by itself. The benefit is avoiding surgery. However, the growth plate is not in the perfect position, which may mean that it doesn't grow normally. This might cause a difference in the shape of the leg (i.e. the ankle could grow wonky) or the injured leg could become shorter than the uninjured leg. This may need future surgery to correct these problems and could cause pain and arthritis in the future.

2. Surgery to fix the bone, usually with screws and a plaster cast for up to 6 weeks. The benefit is that surgery puts the growth plate in the natural position, which many surgeons believe may help the leg grow normally. However, there are risks of surgery, such as the risk of an anaesthetic, infection, wound problems, pain or stiffness, injury to nerves supplying the foot and problems related to the metal implants. There is also still a chance that the leg doesn't grow normally, and often a need for a second operation to remove any metal implants.

Where is the study run from?

Alder Hey Children's NHS Foundation Trust is the sponsor for the study and has overall responsibility for the management of it. This study will be overseen by Oxford Clinical Trials Research Unit (OCTRU) with the day-to-day running of the study being completed by Oxford Trauma and Emergency Care at the University of Oxford.

The research team has a lot of experience in caring for children and young people with injuries and is active in health research. Parents and children have been involved in the development of this study, and are involved in the management.

When is the study starting and how long is it expected to run for?

September 2021 to May 2027

Who is funding the study?

The National Institute for Health and Care Research (NIHR) Health Technology Assessment (HTA) Programme (Reference NIHR132675)

Who is the main contact?

Mr Nick Peterson: 0151 2284811, ODDSocks@ndorms.ox.ac.uk

Contact information

Type(s)

Principal investigator

Contact name

Mr Nick Peterson

Contact details

Chief Investigator, Consultant Children's Orthopaedic Surgeon, Orthopaedic Department, Alder Hey Children's Hospital Foundation Trust
Liverpool
United Kingdom
L12 2AP
+44 (0)151 228 4811
Nicholas.D.Peterson@alderhey.nhs.uk

Type(s)

Scientific

Contact name

Dr ODD SOCKS Study team

Contact details

The University of Oxford, NDORMS Kadoorie Centre, Level 3, John Radcliffe Hospital
Oxford
United Kingdom
OX3 9DU
+44 (0)1865 227902
ODDSocks@ndorms.ox.ac.uk

Type(s)

Public

Contact name

Miss Kinzah Abbasi

Contact details

Trial Coordinator, University of Oxford, NDORMS, Kadoorie Centre, Level 3, John Radcliffe Hospital
Oxford
United Kingdom
OX3 9DU
+44 (0)1865 (2)27902
kinzah.abbasi@ndorms.ox.ac.uk

Additional identifiers**Clinical Trials Information System (CTIS)**

Nil known

Integrated Research Application System (IRAS)

324571

Protocol serial number

CPMS 60457, IRAS 324571

Study information

Scientific Title

ODD SOCKS Study- Outcomes of Displaced Distal tibial fractures- Surgery Or Casts in KidS Study

Acronym

ODD SOCKS

Study objectives

This pragmatic randomised controlled trial aims to evaluate the clinical and cost-effectiveness of surgical reduction, compared to conservative treatment, for the management of displaced Salter Harris-II fractures of the distal tibial physis in children.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 18/01/2024, East Midlands - Nottingham 1 Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 (0)207 104 8115, (0)207 104 8271, (0) 207 104 8089; Nottingham1.rec@hra.nhs.uk), ref: 24/EM/0006

Study design

Pragmatic randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Childhood displaced Salter Harris-II fractures of the distal tibial physis

Interventions

The proposed project is a two-phased study. Phase 1 (internal pilot) will confirm the expected rate of recruitment and pilot data collection procedures in a large-scale multi-centre RCT. Phase 2 is the expansion of the internal pilot into the full definitive trial (main RCT). Phase 2 is the expansion of the pilot into the full definitive trial. A full trial report for the funder and peer-reviewed publications of the main results will be generated after the completion of this phase.

All children aged 8-15 years inclusive presenting to the trial centres with a displaced fracture of the distal tibia involving the physis and metaphysis (i.e. Salter-Harris II) are potentially eligible to take part. Upon presentation, children will receive analgesia and their ankles will be assessed to ensure that the fracture does not require an emergency realignment (i.e. compromising the blood or nerve supply to the foot, or causing potential damage to the skin and other soft tissue

structures). Once any emergency realignment is either performed or confirmed to not be required, temporary immobilisation of the limb for comfort will be applied as per the usual practice of the treating centre. In many hospitals the decision related to definitive treatment is taken in the emergency department by the on-call orthopaedic surgical teams; in others the child may be discharged to an early appointment in the fracture clinic. Owing to the nature of the condition and treatment pathways, the study will be introduced to the patient at the point where definitive care is planned.

After informed consent/assent has been obtained, baseline demographic and injury data, physical function using the PROMIS Mobility CAT, pain intensity using the Wong-Baker FACES Pain Scale and health-related quality of life using the EQ-5D-Y will be collected.

The children will be split into two groups, using a research process called 'randomisation' using a computer randomisation service to fairly allocate treatments:

1. Surgical Reduction- the children in this group will have an anaesthetic or be sedated so their bones can be reset in theatre, and a plaster cast put on their leg. Sometimes, if the doctor thinks it necessary, a small cut will be made and wires, screws or a plate and screws will be inserted to hold the broken bones in position.

2. CONSERVATIVE TREATMENT group – the children in this group will not have the bones reset in position, they will receive a plaster cast for support and to allow the bones to heal.

After treatment, the parents and/or participants will be asked to complete further questionnaires at 6 weeks, 3 months, 6 months, 12 months and 24 months after randomisation.

Data will be collected primarily electronically (with a telephone interview where required) with email and/or text message prompts.

Participants will usually attend an orthopaedic follow-up clinic regularly, as part of standard care and until they are approximately 16 years old, or at least for 2 years after the initial surgery to monitor for signs of complications. In both groups, children will have up to two additional X-rays to what they would have if they were not taking part in this study.

Intervention Type

Procedure/Surgery

Primary outcome(s)

To determine whether children treated with surgical correction have improved function compared with children treated with conservative care measured using the PROMIS Mobility questionnaire, at 24 months post-randomisation

Key secondary outcome(s)

The following secondary outcome measures will be assessed during the first 2-years post-randomisation between surgical reduction and conservative treatment:

1. Function measured using the PROMIS-Mobility Score at 6 weeks, 3, 6 and 12 months post-randomisation
2. Pain scores measured using the Wong-Baker faces pain rating scale at 6 weeks, 3, 6, 12 and 24 months post-randomisation
3. Quality of life measured using EQ-5D-Y at 6 weeks, 3, 6, 12 and 24 months post-randomisation
4. Complication rate measured using data collected in medical records at 6/8 weeks, 12 and 24 months post-randomisation

5. Satisfaction with cosmetic appearance of the leg measured using the Visual Analogue Scale (VAS) of cosmesis at 3, 12 and 24 months post-randomisation
6. Satisfaction with the treatment received measured using a Likert Scale at 3, 12 and 24 months post-randomisation
7. Child educational participation recording educational absences measured using a parent-reported questionnaire at 6 weeks, 3, 6, 12 and 24 months post-randomisation
8. Leg Length Measurement measured using the 'Tape Measurement Method' by a clinician at 12 and 24 months
9. Angular deformity and PPC (growth arrest) measured using radiographic images 24 months post-randomisation
10. Resource use measured using a bespoke electronic resource use questionnaire, and parental absence from work at 6 weeks, 3, 6, 12 and 24 months post-randomisation

Completion date

31/05/2027

Eligibility

Key inclusion criteria

1. Aged 8 to 15 years old inclusive
2. There is radiographic evidence of a displaced fracture of the distal tibia involving the physis and metaphysis (Salter Harris II); with or without a corresponding fibula fracture
3. The treating clinician believes that they may benefit from surgical reduction +/- fixation

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

8 years

Upper age limit

15 years

Sex

All

Key exclusion criteria

1. The injury is more than 7 days old
2. The fracture is open
3. They have an intra-articular fracture that requires fixation to restore the joint surface
4. They have any other contralateral (Opposite-sided) ankle fracture/injury
5. There is evidence that the patient and/or parent/guardian would be unable to adhere to trial procedures or complete follow-up
6. The patient has previously been enrolled on the ODD SOCKS Study

Date of first enrolment

22/05/2024

Date of final enrolment

01/12/2026

Locations

Countries of recruitment

United Kingdom

England

Scotland

Wales

Study participating centre**NHS Grampian**

Summerfield House

2 Eday Road

Aberdeen

United Kingdom

AB15 6RE

Study participating centre**NHS Lothian**

Waverley Gate

2-4 Waterloo Place

Edinburgh

United Kingdom

EH1 3EG

Study participating centre**Cardiff & Vale University Lhb**

Woodland House

Maes-y-coed Road

Cardiff

United Kingdom

CF14 4HH

Study participating centre

Bradford Royal Infirmary

Duckworth Lane
Bradford
United Kingdom
BD9 6RJ

Study participating centre

Doncaster Royal Infirmary

Armthorpe Road
Doncaster
United Kingdom
DN2 5LT

Study participating centre

Macclesfield District General Hospital

Victoria Road
Macclesfield
United Kingdom
SK10 3BL

Study participating centre

Gloucester Royal Hospital

Great Western Road
Gloucester
United Kingdom
GL1 3NN

Study participating centre

Basingstoke and North Hampshire Hospitals

Aldermaston Road
Basingstoke
United Kingdom
RG24 9NA

Study participating centre

Leeds General Infirmary

Great George Street
Leeds
United Kingdom
LS1 3EX

Study participating centre
Kent and Canterbury Hospital
Ethelbert Road
Canterbury
United Kingdom
CT1 3NG

Study participating centre
Milton Keynes University Hospital
Standing Way
Eaglestone
Milton Keynes
United Kingdom
MK6 5LD

Study participating centre
Musgrove Park Hospital (taunton)
Musgrove Park Hospital
Taunton
United Kingdom
TA1 5DA

Study participating centre
Northwick Park Hospital
Watford Road
Harrow
United Kingdom
HA1 3UJ

Study participating centre
John Radcliffe Hospital
Headley Way
Headington
Oxford
United Kingdom
OX3 9DU

Study participating centre

Royal Berkshire Hospital

London Road
Reading
United Kingdom
RG1 5AN

Study participating centre

St Georges Hospital

Blackshaw Road
London
United Kingdom
SW17 0QT

Study participating centre

Royal Cornwall Hospital (treリスケ)

Treliske
Truro
United Kingdom
TR1 3LJ

Study participating centre

St Peters Hospital

Guildford Road
Chertsey
United Kingdom
KT16 0PZ

Study participating centre

Alder Hey Hospital

Eaton Road
West Derby
Liverpool
United Kingdom
L12 2AP

Study participating centre

University Hospital Coventry & Warwickshire

Clifford Bridge Road
Walsgrave

Coventry
United Kingdom
CV2 2DX

Study participating centre

Hull Royal Infirmary

Anlaby Road
Hull
United Kingdom
HU3 2JZ

Study participating centre

Royal Derby Hospital

Uttoxeter Road
Derby
United Kingdom
DE22 3NE

Study participating centre

University Hospitals of North Midlands NHS Trust

Newcastle Road
Stoke-on-trent
United Kingdom
ST4 6QG

Study participating centre

Southampton General Hospital

Tremona Road
Southampton
United Kingdom
SO16 6YD

Study participating centre

Mid and South Essex NHS Foundation Trust

Prittlewell Chase
Westcliff-on-sea
United Kingdom
SS0 0RY

Study participating centre
Sunderland Royal Hospital
Kayll Road
Sunderland
United Kingdom
SR4 7TP

Study participating centre
Queen Elizabeth the Queen Mother Hospital
St. Peters Road
Margate
United Kingdom
CT9 4AN

Study participating centre
Whiston Hospital (site)
Whiston Hospital
Warrington Road
Prescot
United Kingdom
L35 5DR

Study participating centre
Nottingham University Hospitals NHS Trust
Trust Headquarters
Queens Medical Centre
Derby Road
Nottingham
United Kingdom
NG7 2UH

Study participating centre
Mid Yorkshire Teaching NHS Trust
Pinderfields Hospital
Aberford Road
Wakefield
United Kingdom
WF1 4DG

Study participating centre

Portsmouth Hospitals University National Health Service Trust

Queen Alexandra Hospital
Southwick Hill Road
Cosham
Portsmouth
United Kingdom
PO6 3LY

Study participating centre

Lancashire Teaching Hospitals NHS Foundation Trust

Royal Preston Hospital
Sharoe Green Lane
Fulwood
Preston
United Kingdom
PR2 9HT

Study participating centre

West Suffolk NHS Foundation Trust

West Suffolk Hospital
Hardwick Lane
Bury St. Edmunds
United Kingdom
IP33 2QZ

Study participating centre

Betsi Cadwaladr Uhb

Royal Alexandra Hospital
Marine Drive
Rhyl
United Kingdom
LL18 3AS

Study participating centre

South Tees Hospitals NHS Foundation Trust

James Cook University Hospital
Marton Road
Middlesbrough
United Kingdom
TS4 3BW

Study participating centre
Sheffield Childrens Hospital
Western Bank
Sheffield
United Kingdom
S10 2TH

Study participating centre
Mersey and West Lancashire Teaching Hospitals NHS Trust
Whiston Hospital
Warrington Road
Prescot
United Kingdom
L35 5DR

Study participating centre
University Hospitals of Leicester NHS Trust
Leicester Royal Infirmary
Infirmary Square
Leicester
United Kingdom
LE1 5WW

Study participating centre
Belfast Health and Social Care Trust
Trust Headquarters
A Floor - Belfast City Hospital
Lisburn Road
Belfast
United Kingdom
BT9 7AB

Study participating centre
Royal Free London NHS Foundation Trust
Royal Free Hospital
Pond Street
London
United Kingdom
NW3 2QG

Study participating centre

Birmingham Women's NHS Foundation Trust

Birmingham Womens Hospital
Metchley Park Road
Birmingham
United Kingdom
B15 2TG

Study participating centre

Norfolk and Norwich University Hospitals NHS Foundation Trust

Colney Lane
Colney
Norwich
United Kingdom
NR4 7UY

Study participating centre

Barts Health NHS Trust

The Royal London Hospital
80 Newark Street
London
United Kingdom
E1 2ES

Study participating centre

Maidstone and Tunbridge Wells NHS Trust

The Maidstone Hospital
Hermitage Lane
Maidstone
United Kingdom
ME16 9QQ

Study participating centre

Kettering General Hospital NHS Foundation Trust

Rothwell Road
Kettering
United Kingdom
NN16 8UZ

Study participating centre

Royal Devon and Exeter NHS Foundation Trust

Royal Devon & Exeter Hospital

Barrack Road
Exeter
United Kingdom
EX2 5DW

Study participating centre
University Hospitals Plymouth NHS Trust
Derriford Hospital
Derriford Road
Derriford
Plymouth
United Kingdom
PL6 8DH

Study participating centre
Manchester University NHS Foundation Trust
Cobbett House
Oxford Road
Manchester
United Kingdom
M13 9WL

Sponsor information

Organisation
Alder Hey Children's NHS Foundation Trust

ROR
<https://ror.org/00p18zw56>

Funder(s)

Funder type
Government

Funder Name
Health Technology Assessment Programme

Alternative Name(s)
NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Study website	Study website	11/11/2025	11/11/2025	No	Yes