

AFI-targeted confocal endomicroscopy in Barrett's oesophagus

Submission date 12/06/2017	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 25/07/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 23/09/2022	Condition category Cancer	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

<https://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-study-looking-at-a-new-test-to-find-cell-changes-for-people-with-barretts-oesophagus-ace-b>

Contact information

Type(s)

Public

Contact name

Ms Michele Bianchi

Contact details

MRC Cancer Centre
Hutchison MRC Research Centre
University of Cambridge
Cambridge
United Kingdom
CB2 0XZ

Type(s)

Scientific

Contact name

Dr Massimiliano di Pietro

Contact details

MRC Cancer Centre
Hutchison MRC Research Centre
University of Cambridge
Cambridge
United Kingdom

CB2 0XZ
+44 122 3763994
md460@mrc-cu.cam.ac.uk

Additional identifiers

Protocol serial number

33291

Study information

Scientific Title

Cross-over randomised study to evaluate the combination of autofluorescence imaging and confocal laser endomicroscopy to diagnose dysplasia in Barrett's oesophagus

Study objectives

The aim of this study is to test a combination of enhanced endoscopic imaging (autofluorescence imaging and confocal laser endomicroscopy) in combination to molecular tests on tissue samples to allow diagnosis of dysplasia and early cancer in Barrett's oesophagus.

Ethics approval required

Old ethics approval format

Ethics approval(s)

East of England - Cambridgeshire and Hertfordshire Research Ethic Committee, 05/04/2017, ref: 16/EE/0554

Study design

Randomised; Interventional; Design type: Diagnosis, Imaging

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Barrett's oesophagus

Interventions

As part of this study, patients referred with flat Barrett's oesophagus without evidence of visible lesions receive two endoscopies in a randomised order (standard procedure and experimental procedure).

Standard procedure: This includes a standard endoscopy with multiple random biopsies according to the Seattle protocol (targeted biopsies on visible lesions + 4 biopsies every 2 cm within the Barrett's oesophagus). The endoscopist is only be allowed to use white light high resolution endoscopy.

Experimental procedure: This procedure uses multimodal imaging. The endoscopist uses autofluorescence imaging (AFI) to identify area at risk of dysplasia. These are analysed with Probed-Based Confocal Laser Endomicroscopy (pCLE) to make a real-time optical diagnosis (Barrett's with or without dysplasia). Two targeted biopsies stored in formalin are then taken for histology and biomarkers from each AFI targeted location. In patients with no AFI positive areas one random location are used for pCLE analysis and molecular biomarkers for every 5cm of maximum extent of the Barrett's. The biopsies on AFI targeted areas processed for standard histology are used for clinical purpose as well and are part of the final histologic diagnosis and potentially inform treatment as per clinical guidelines.

In addition to histological diagnosis, the biopsies taken during the experimental procedure are tested for a panel of 3 molecular biomarkers.

After the first study procedure, the patient arecheduled for a second endoscopy 8-12 weeks after with the alternative protocol.

With the second procedure the patient completes the study and referred for standard clinical management based on the histopathological result of the biopsies taken during the two study procedures.

Intervention Type

Other

Primary outcome(s)

The diagnostic accuracy for any grade of dysplasia of pCLE is measured comparing real-time optical diagnosis of dysplasia by pCLE on AFI-positive areas (experimental procedure) with the gold standard histologic diagnosis (overall pathological diagnosis from experimental and standard procedures).

Key secondary outcome(s)

1. Added value of the use of molecular biomarkers to the optical biopsy for the diagnosis of any grade of dysplasia. This will be measured by testing molecular biomarkers on tissue biopsies. The biomarkers result will be integrated with that of the optical diagnosis and the results will be compared with that of the gold standard histologic diagnosis.
2. Diagnostic accuracy for any grade of dysplasia of a panel of biomarkers performed on AFI-targeted biopsies. This will be measured by testing molecular biomarkers on tissue biopsies and comparing the results with the gold standard histologic diagnosis
3. Time to perform AFI-targeted pCLE vs gold standard (Seattle protocol). This will be measured as time from the beginning to the end of each endoscopic procedure. Standard and experimental procedure time will be compared.
4. Costs to perform AFI-targeted pCLE +/- biomarkers and conventional endoscopic surveillance with Seattle protocol. This will be measured by the costs of a single use of pCLE probe and laboratory costs of molecular biomarkers for the experimental procedure and costs for processing biopsies and costs of pathology time for histologic diagnosis for the standard procedure.
5. Patient-reported experience and outcome measures, including acceptability and anxiety levels. This outcome will be measured using 2 validated questionnaires: a 10-point visual analogue scale (VAS, 0 = worst and 10 = best), filled by participants before and after each procedure and 6-item state-trait anxiety inventory (STAI -6), completed after each procedure.

Completion date

29/02/2020

Eligibility

Key inclusion criteria

1. Able to read, comprehend, and complete the consent form
2. Aged ≥ 18
3. Diagnosed with dysplastic or non-dysplastic BO at least 2 cm in length if circumferential (C2) or 3 cm if not circumferential (M3)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

146

Key exclusion criteria

1. Oesophagitis (Los Angeles grade $\geq B$)
2. Previous oesophagectomy or known oesophageal abnormality (e.g. fistula or severe oesophageal stricture)
3. Previous evidence of oesophageal adenocarcinoma
4. Previous history of endoscopically visible BO-related neoplasia
5. Known allergy to fluorescein
6. Severe or uncontrolled asthma
7. Coagulopathy or anticoagulant/antiplatelet therapy for high risk conditions
8. Active or severe cardiopulmonary disease or decompensated liver disease
9. Pacemaker or other intra-cardiac electric device

Date of first enrolment

02/05/2017

Date of final enrolment

31/12/2019

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Cambridge University Hospital
Hills Road
Cambridge
United Kingdom
CB2 0QQ

Study participating centre
Nottingham University Hospital
The Queen's Medical Centre Campus
Derby Road
Nottingham
United Kingdom
NG7 2UH

Sponsor information

Organisation
Cambridge University Hospitals NHS Foundation Trust

ROR
<https://ror.org/04v54gj93>

Funder(s)

Funder type
Charity

Funder Name
Cancer Research UK

Alternative Name(s)
CR_UK, Cancer Research UK - London, Cancer Research UK (CRUK), CRUK

Funding Body Type
Private sector organisation

Funding Body Subtype

Other non-profit organizations

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from from the Chief Investigator, Dr Massimiliano di Pietro [md460@mrc-cu.cam.ac.uk]

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Plain English results			23/09/2022	No	Yes