

Radiofrequency denervation for low back pain

Submission date 17/08/2021	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 09/11/2021	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 30/01/2026	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Long-term low back pain is common, affecting 10-15% of adults. It can significantly impair the health, mood, and daily lives of people who have it. One type of low back pain is caused by the small joints between the bones in the lower back. Treatments include painkillers, exercise and talking therapies. However, if people do not get better with these treatments, they can be offered radiofrequency “denervation”. Denervation involves placing a needle in the nerve to the painful joint, which is heated up to cause a break in the nerve. The purpose of this is to stop the nerve from sending pain messages to the brain. Denervation is low risk and is used widely in the National Health Service (NHS) but it is not known if this procedure definitely reduces pain or is a good way to spend NHS money. This study aims to find out if denervation reduces low back pain and is good value for money.

Who can participate?

Patients aged 18 years or older with chronic moderate to severe low back pain who are eligible for radiofrequency denervation treatment

What does the study involve?

Participants are randomly allocated into one of two groups. Half will have the denervation and half will have a placebo treatment, which involves placement of the needle in the nerve but without heating it up so the nerve is not affected. Participants whose symptoms do not improve after 3 months will be offered the chance to receive the other treatment. This means that patients who had no improvement because they had the placebo treatment first would have the opportunity for denervation the second time. Participants will be asked questions about their low back pain, ability to carry out daily tasks including their work, their general health, and mental well-being over the next 2 years. Information will also be collected to find out if denervation is good value for money.

What are the possible benefits and risks of participating?

There are no guaranteed benefits of participating, but the results from this study may help improve the treatment of people with low back pain in the future. The risks associated with taking part in this study are the same as the risks of having this procedure as part of usual care, and are the same for both the denervation and the placebo treatment. However, if participants do not experience an improvement in pain after 3 months, they will be offered the chance to receive the other treatment. Therefore, there is the possibility that participants will have two

procedures (denervation and placebo treatment). The procedure is low risk and serious side effects are rare.

Where is the study run from?
University of Bristol (UK)

When is the study starting and how long is it expected to run for?
August 2018 to September 2025

Who is funding the study?
National Institute for Health Research (NIHR) (UK)

Who is the main contact?
Vikki Wylde (Chief Investigator) or Kate Ashton (Trial Manager)
radical-study@bristol.ac.uk

Contact information

Type(s)

Public

Contact name

Ms Kate Ashton

ORCID ID

<https://orcid.org/0000-0002-9163-0512>

Contact details

Bristol Trials Centre
Bristol Medical School
University of Bristol
1-5 Whiteladies Road
Bristol
United Kingdom
BS8 1NU

Type(s)

Scientific

Contact name

Dr Vikki Wylde

ORCID ID

<https://orcid.org/0000-0002-8460-1529>

Contact details

Musculoskeletal Research Unit
Translational Health Sciences
Bristol Medical School
University of Bristol
Learning and Research Building

Southmead Hospital
Bristol
United Kingdom
BS10 5NB

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

285322

Protocol serial number

CPMS 46543, NIHR127457

Study information

Scientific Title

Radiofrequency denervation for chronic and moderate to severe low back pain (RADICAL)

Acronym

RADICAL

Study objectives

Radiofrequency denervation compared to a placebo treatment reduces the severity of pain at 3 months after the intervention.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 30/07/2021, London - Fulham Research Ethics Committee (Barlow House, 3rd Floor, 4 Minshull Street, Manchester, M1 3DZ, UK; +44 (0)207 104 8035; fulham.rec@hra.nhs.uk), REC ref: 21/LO/0471

Study design

Randomized; Both; Design type: Treatment, Other, Qualitative

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Chronic and moderate to severe low back pain

Interventions

Trial participants will be randomised in a 1:1 ratio to receive either radiofrequency denervation (RFD) of the lumbar medial branches of the dorsal rami; or placebo treatment, which will follow the same protocol, but the electrode tip temperature will not be raised.

Randomisation will be performed by a member of the theatre staff, not involved in participant follow-up, via a secure internet-based randomisation system ensuring allocation concealment. Participants will be allocated in a 1:1 ratio to RFD or placebo treatment. The allocation, prepared by a statistician independent of the trial team, will be computer-generated and blocked with varying block sizes. Randomisation will be stratified by operator to ensure that any operator effect is distributed equally across groups.

Participants who do not experience a clinically meaningful improvement in pain 3 months after randomisation will be offered "repeat RFD" but with the alternative intervention to the one provided at the outset without disclosing the original allocation.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Patient-reported low back pain (LBP) pain severity over the past week, measured using a 0-10 pain Numeric Rating Scale (NRS); Timepoint(s): 3 months post-randomisation

Key secondary outcome(s)

1. Functional disability measured using the Oswestry Disability Index (ODI) version 2.1b at baseline, 3, 6, 12, 18 and 24 months post randomisation
2. Health-related quality of life (HRQoL) measured using EQ-5D-5L at baseline, 6 weeks, and 3, 6, 12, 18 and 24 months post randomisation
3. General health measured using SF-12 Physical Component Score at baseline, 3, 6, 12, 18 and 24 months post randomisation
4. Mental health measured using SF-12 Mental Component Score at baseline, 3, 6, 12, 18 and 24 months post randomisation
5. Time to pain recovery measured using time from randomisation until the first timepoint at which the patient reports a pain reduction of $\geq 60\%$ that remains at $\geq 60\%$ lower than baseline at their subsequent timepoint. Pain severity over the past week will be measured using a 0-10 pain Numerical Rating Scale (NRS), administered at baseline, 2, 4, 6, 8 and 10 weeks, and 3, 6, 12, 18 and 24 months post randomisation.
Updated 08/07/2024: Pain severity over the past week will be measured using a 0-10 pain Numerical Rating Scale (NRS), administered at baseline, 2 and 6 weeks, and 3, 6, 12, 18 and 24 months post randomisation
6. Uptake of offer for repeat RFD. This will be offered to participants who are eligible 3 months after randomisation and can be taken up by participants up to 24 months
7. Satisfaction with treatment outcome measured using Likert scale at 3, 6, 12, 18 and 24 months post randomisation
8. Adverse events measured using active capture of adverse events at 2 and 6 weeks, and 3, 6, 12, 18 and 24 months post randomisation
9. Work outcomes: Work status and days lost from work and usual activities due to LBP measured using the Work Productivity and Activity Impairment (WPAI) questionnaire at baseline, 3, 6, 12, 18 and 24 months post randomisation
10. Healthcare utilisation, including medications, measured using a patient-reported resource use questionnaire at baseline, 3, 6, 12, 18 and 24 months post randomisation, and medical records

Completion date

30/09/2026

Eligibility

Key inclusion criteria

Current inclusion criteria as of 22/05/2023:

1. 18 years of age or older
2. LBP is the primary source of pain
3. Positive response to a single diagnostic MBB with no steroids administered
4. Chronic LBP (>3 months duration), assumed due to the fact patient was listed for MBB
5. Moderate to severe LBP (pain NRS score ≥ 5 on Baseline Questionnaire)
6. Listed for RFD by their clinical care team

Previous inclusion criteria:

1. 18 years of age or older
2. Chronic moderate to severe LBP (>3 months duration, pain NRS score ≥ 5 for usual pain over the past week at the time of screening)
3. LBP is the primary source of pain
4. Referred to a pain or spinal clinic
5. Listed for MBB by their clinical care team (due to clinical suspicion or clinical features suggesting that the main source of LBP is from a facet joint)
6. Positive response to a single diagnostic MBB with 1 ml or less of local anaesthetic at each level (no steroids)
7. Listed for RFD by their clinical care team

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

100 years

Sex

All

Total final enrolment

180

Key exclusion criteria

Current exclusion criteria as of 22/05/2023:

1. Known pregnancy

2. Severe depression (Hospital Anxiety and Depression Scale (HADS) depression score ≥ 15)
3. Known previous RFD
4. Known previous back surgery where metal-work has been used in the lumbar spine
5. Pacemaker or implantable cardioverter-defibrillator
6. Clinical suspicion that an alternative diagnosis is the reason for low back pain (as defined by NICE, including, but not limited to: metastatic spinal cord compression, spinal injury, spondyloarthritis, or cancer)
7. Prisoner
8. Patient lacks capacity to consent
9. Existing co-enrolment in another clinical study if: i) the intervention in the other study is expected to influence the primary outcome (this will be considered by a senior clinician on a case-by-case basis); ii) it is considered too burdensome for the patient; or iii) it is not permitted by the other study

Previous exclusion criteria:

1. Known pregnancy
2. Unwilling or unable to tolerate procedure
3. Severe depression (Hospital Anxiety and Depression Scale (HADS) depression score ≥ 15)
4. Known previous RFD
5. Known previous back surgery where metal-work has been used in the lumbar spine
6. Pacemaker or implantable cardioverter-defibrillator
7. Clinical suspicion that an alternative diagnosis is the reason for low back pain (as defined by NICE, including, but not limited to: metastatic spinal cord compression, spinal injury, spondyloarthritis, or cancer)
8. Prisoner
9. Patient lacks capacity to consent
10. Existing co-enrolment in another clinical study if: i) the intervention in the other study is expected to influence the primary outcome (this will be considered by a senior clinician on a case-by-case basis); ii) it is considered too burdensome for the patient; or iii) it is not permitted by the other study

Date of first enrolment

16/03/2022

Date of final enrolment

29/01/2026

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

St. James's University Hospital

Beckett Street

Leeds
England
LS9 7TF

Study participating centre
Queen Elizabeth Hospital
Mindelsohn Way
Edgbaston
Birmingham
England
B15 2GW

Study participating centre
Southmead Hospital
Southmead Road
Westbury-On-Trym
Bristol
England
BS10 5NB

Study participating centre
John Radcliffe Hospital
Headley Way
Headington
Oxford
England
OX3 9DU

Study participating centre
Kings Mill Hospital
Mansfield Road
Sutton-In-Ashfield
England
NG17 4JL

Study participating centre
Solent NHS Trust
Highpoint Venue
Bursledon Road

Southampton
England
SO19 8BR

Study participating centre
Walsgrave General Hospital
Clifford Bridge Road
Coventry
England
CV2 2DX

Study participating centre
The Royal London Hospital
80 Newark Street
London
England
E1 2ES

Study participating centre
The Walton Centre
Lower Lane
Liverpool
England
L9 7LJ

Study participating centre
James Cook University Hospital
Marton Road
Middlesbrough
England
TS4 3BW

Study participating centre
Freeman Hospital
Freeman Road
High Heaton
Newcastle upon Tyne
England
NE7 7DN

Study participating centre

NHS Grampian

Summerfield House
2 Eday Road
Aberdeen
Scotland
AB15 6RE

Study participating centre

Royal Berkshire Hospital

London Road
Reading
England
RG1 5AN

Study participating centre

City Hospital

Dudley Road
Birmingham
England
B18 7QH

Study participating centre

Epsom and St Helier University Hospitals NHS Trust

St Helier Hospital
Wrythe Lane
Carshalton
England
SM5 1AA

Study participating centre

Liverpool University Hospitals NHS Foundation Trust

Royal Liverpool University Hospital
Prescot Street
Liverpool
England
L7 8XP

Study participating centre

Royal Orthopaedic Hospital

The Woodlands
Bristol Road South
Northfield
Birmingham
England
B31 2AP

Study participating centre

Lancashire Teaching Hospitals NHS Foundation Trust

Royal Preston Hospital
Sharoe Green Lane
Fulwood
Preston
England
PR2 9HT

Study participating centre

Sherwood Forest Hospitals NHS Foundation Trust

Kings Mill Hospital
Mansfield Road
Sutton-in-ashfield
England
NG17 4JL

Study participating centre

Whittington Health NHS Trust

The Whittington Hospital
Magdala Avenue
London
England
N19 5NF

Study participating centre

East Kent Hospitals University NHS Foundation Trust

Kent & Canterbury Hospital
Ethelbert Road
Canterbury
England
CT1 3NG

Study participating centre
The Royal Orthopaedic Hospital NHS Foundation Trust
The Woodlands
Bristol Road South
Northfield
Birmingham
England
B31 2AP

Study participating centre
Solent NHS Trust
Solent NHS Trust Headquarters
Highpoint Venue
Bursledon Road
Southampton
England
SO19 8BR

Sponsor information

Organisation
North Bristol NHS Trust

ROR
<https://ror.org/036x6gt55>

Funder(s)

Funder type
Government

Funder Name
NIHR Evaluation, Trials and Studies Co-ordinating Centre (NETSCC); Grant Codes: NIHR127457

Results and Publications

Individual participant data (IPD) sharing plan

Data will not be made available for sharing until after the publication of the main results of the study. Thereafter, anonymised individual patient data will be made available for secondary research, conditional on assurance from the secondary researcher that the proposed use of the

data is compliant with the MRC Policy on Data Sharing regarding scientific quality, ethical requirements and value for money. A minimum requirement with respect to scientific quality will be a publicly available pre-specified protocol describing the purpose, methods and analysis of the secondary research, e.g. a protocol for a Cochrane systematic review. Anonymised recruitment consultation and interview transcripts may also be used to support the teaching of qualitative research methods. Please contact Vikki Wylde using the following email: radical-study@bristol.ac.uk.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article		27/07/2024	29/07/2024	Yes	No
HRA research summary			28/06/2023	No	No