

A comparison of rituximab and modified Ponticelli regimen (alternating steroids and cyclophosphamide) for treatment of primary membranous nephropathy

Submission date 04/04/2022	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 15/04/2022	Overall study status Completed	<input checked="" type="checkbox"/> Protocol
Last Edited 23/05/2022	Condition category Urological and Genital Diseases	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Membranous nephropathy (MN) is one of the most common causes of nephrotic syndrome in adults, which is characterised by the presence of protein in the urine. There are two treatment options for this disease, both of which aim to achieve remission – the reduction of protein excretion in the urine. The modified Ponticelli regimen with cyclical steroids and cyclophosphamide is used as first-line treatment but is associated with several adverse effects such as infections, infertility and low blood cell counts. Rituximab has emerged as a newer option for treatment but it has not been compared against the modified Ponticelli regimen in head-to-head trials. The aim of this study is to assess the effectiveness of rituximab versus the modified Ponticelli regimen in inducing remission in MN.

Who can participate?

Patients aged over 18 years with primary membranous nephropathy with more than 3.5 g protein excretion in urine

What does the study involve?

Some patients can have spontaneous resolution of protein excretion in urine. Therefore all patients with primary MN will be given a 3-month observation period to watch for spontaneous resolution. Patients with severe symptoms or those with persistent protein excretion will be randomly allocated into one of the two groups and will receive either rituximab or the modified Ponticelli regimen. The rituximab group will receive two injections of rituximab on days 1 and 15. Their CD19 level (which indicates their B cell levels) will be checked at 1 and 6 months. B cells are suppressed by rituximab and a low level indicates that the rituximab was effective at suppressing the B cells. The modified Ponticelli regimen consists of alternating months of steroids and cytotoxic agents for a total of 6 months. Steroids are given in months 1, 3 and 5 which will include three injections of methylprednisolone on days 1-3 followed by prednisolone tablets. Cyclophosphamide tablets will be given in months 2, 4 and 6.

What are the possible benefits and risks of participating?

The participants would receive two treatments that are routinely being used for this disease. All drugs used in this study have an established role in treatment of MN and all are associated with specific side effects. Steroids can cause weight gain, high blood sugar, bone weakness, muscle wasting and an increased risk of infections. Cyclophosphamide can cause infections, infertility, bladder problems, low blood counts as well as cancer. Rituximab is typically associated with reactions during infusion including flushing, itching and sometimes low blood pressure.

Where is the study run from?

Muljibhai Patel Urological Hospital, Nadiad (India)

When is the study starting and how long is it expected to run for?

June 2018 to June 2021

Who is funding the study?

Muljibhai Patel Society for Research in Nephrology, Nadiad, Gujarat (India)

Who is the main contact?

Dr Sandhya Suresh
sandymmc@gmail.com

Contact information

Type(s)

Principal investigator

Contact name

Dr Sandhya Suresh

ORCID ID

<https://orcid.org/0000-0001-8821-0579>

Contact details

Flat 1 A, PGP Manor
29, Barnaby Road
Kilpauk
Chennai
India
600010
+91 9381093878
sandymmc@gmail.com

Additional identifiers

Protocol serial number

525

Study information

Scientific Title

INdian COntrolled trial of RiTuximab vs Modified Ponticelli regimen for treatment of primary membranous nephropathy

Acronym

INCORIT-M

Study objectives

Rituximab is non-inferior to modified Ponticelli regimen in inducing remission (complete or partial) at 6 months in patients with primary membranous nephropathy (PMN).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 16/08/2018, Muljibhai Patel Society for Research in Nephro-Urology Ethics Committee (Muljibhai Patel Urological Hospital, Dr. Virendra Desai Road, Nadiad - 387001, Gujarat, India; +91 (0)268 2520330; info@mpuh.org), ref: EC/525/2018

Study design

Interventional randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Primary membranous nephropathy

Interventions

Intervention arm - Rituximab injection 500mg IV given on days 1 and 15

Control arm - Modified Ponticelli regimen (cyclical steroids/cyclophosphamide for 6 months)

Months 1, 3 and 5: 1 g IV methylprednisolone daily (Days 1–3), then oral prednisolone (0.5 mg/kg /day) for 27 days (Days 4–30).

Months 2, 4 and 6: Oral cyclophosphamide (2.0 mg/kg/day) for 30 days.

The researchers used a random number-producing algorithm for block randomisation using sealed envelope online software. Randomisation and treatment allocation were then done by the primary investigator using this randomisation list in a concealed manner in a 1:1 manner using sealed envelopes.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Rituximab, cyclophosphamide, steroids

Primary outcome(s)

The proportion of patients reaching complete or partial remission at 6 months, defined according to the 2012 KDIGO guidelines as follows:

Complete remission: a reduction of proteinuria to <0.3 g/24 h (uPCR <300 mg/g) with normal serum albumin concentration and normal serum creatinine

Partial remission: a reduction of proteinuria to <3.5 g/day (uPCR <3500 mg/g) and a 50% or greater reduction from peak values accompanied by an improvement or normalisation of serum albumin concentration and stable serum creatinine

1. Proteinuria measured using the urine protein creatinine ratio (using Pyrogallol red and Jaffe kinetic reaction test methodology) at baseline and 6 months
2. Serum albumin concentration measured using the Bromocresol green test at baseline and 6 months
3. Serum creatinine measured using the Jaffe kinetic test at baseline and 6 months

Key secondary outcome(s)

1. The proportion of patients reaching either complete or partial remission at 12 months after therapy (measured as per the primary outcome measure)
2. The proportion of patients with relapsing nephrotic syndrome measured among patients who previously underwent partial remission or complete remission at any time during the study period as determined by proteinuria (measured as per the primary outcome measure)
3. The time to relapse of nephrotic syndrome after initial remission as determined by proteinuria (measured as per the primary outcome measure)
4. Serum anti-PLA2R levels measured using EUROIMMUN Anti-PLA2R ELISA (quantitative method) before treatment and at 6 and 12 months post-therapy
5. Efficacy outcome variables during the study period including the trend of the magnitude of proteinuria, serum albumin, serum proteins, serum creatinine and estimated GFR (measured by the CKD-EPI formula) at baseline, 6 and 12 months
6. Proportion of patients developing the following adverse events during the study period, as determined by clinical evaluation including patient interviews and patient notes and lab investigations:
 - 6.1. Infections: symptoms such as fever, cough, dysuria; signs of infection such as fever, respiratory system signs including rales/crepitations, skin lesions of Tinea; lab investigations including total white blood cell (WBC) count (by automated cell counter machine), urine routine analysis (using Multistix and microscopic examination of urine sediment after centrifugation), chest X-ray
 - 6.2. Leucopenia determined by total WBC count (by automated cell counter machine)
 - 6.3. Infusion reaction determined by clinical evaluation
 - 6.4. Acute kidney injury after the start of therapy determined by measuring serum creatinine using the Jaffe kinetic reaction

Completion date

30/06/2021

Eligibility

Key inclusion criteria

1. Patients older than 18 years who provide written informed consent.
2. Biopsy-proven primary MN within 2 years of enrolment with nephrotic range proteinuria denoted by 24-hour urine protein ≥ 3.5 g or UPCR (urine protein:creatinine ratio) ≥ 3500 mg/g as well as the following:

- 2.1. Serology or biopsy positive for AntiPLA2R
- 2.2. Serology or biopsy negative for AntiPLA2R and secondary causes ruled out
- 2.3. Evaluation for secondary causes was done in all patients, even in patients who were positive for Anti-PLA2R antibodies on serology or PLA2R antigen on biopsy because these have also been found in some cases of secondary MN
3. Estimated GFR ≥ 30 ml/min/1.73m². The CKD-EPI creatinine equation was used for the calculation of the eGFR
4. Treatment with an ACEI or ARB for at least 3 months before enrolment

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

68

Key exclusion criteria

1. Secondary MN
2. Active serious infections
3. Pregnant women
4. Suspected or known hypersensitivity to either interventional drug
5. Patients with persistently low estimated GFR < 30 ml/min/1.73m² in the absence of acute causes such as acute tubular injury, renal vein thrombosis and others

Date of first enrolment

26/08/2018

Date of final enrolment

03/10/2020

Locations**Countries of recruitment**

India

Study participating centre

Muljibhai Patel Urological Hospital
Dr. Virendra Desai Road

Nadiad
India
387001

Sponsor information

Organisation

Muljibhai Patel Urological Hospital

ROR

<https://ror.org/059h1d250>

Funder(s)

Funder type

Research organisation

Funder Name

Muljibhai Patel Society for Research in Nephro-Urology

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study will be published as a supplement to the subsequent results publication

IPD sharing plan summary

Published as a supplement to the results publication

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet			05/04/2022	No	Yes
Protocol file			12/04/2022	No	No