

Peripheral arterial disease, high blood pressure and aneurysm screening trial

Submission date 24/02/2022	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 01/06/2022	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 18/03/2026	Condition category Circulatory System	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

An abdominal aortic aneurysm (AAA) is a swelling of the main blood vessel (aorta) in the abdomen. If the swelling gets too large the aorta can burst and this is usually fatal. In order to prevent bursting, AAAs can be repaired with surgery. This is usually carried out when the size of the AAA is more than 5.5 cm in diameter because below this size, the risk of it bursting is lower than the risks associated with surgery. AAAs are usually asymptomatic before bursting but can be easily and safely diagnosed by ultrasound scanning.

People with peripheral arterial disease (PAD) (furred up blood vessels in the legs) and high blood pressure (BP) are at high risk of heart attacks, strokes and other cardiovascular (CV) problems. 30 to 40% of people over the age of 65 years have PAD and/or high BP. At least a third of these people do not know why they have these problems.

Men are invited to an ultrasound scan when they are aged 65 years to check them for an abdominal aortic aneurysm (AAA). Attendance for AAA screening is good, with eight out of ten men attending. The AAA programme represents a good opportunity to identify and treat men with PAD and/or high BP with minimal extra effort. It is known from speaking to men who have been for AAA screening that they think it is a good idea to check for other conditions at the same time. Women are not invited for AAA screening because they rarely get AAAs. This means a different approach will be required to screen women.

The overall aim of the programme is to find out if screening for PAD and high BP at the same time as screening for AAA improves CV health. The main aim of this study is to assess the feasibility of adding PAD+BP screening to AAA screening by assessing attendance for screening. PHAST-F will also provide detailed data to 1) undertake a process evaluation; 2) help plan the delivery of the main trial; and 3) inform the clinical and economic evaluation of the main trial. Screening device(s) identified from an earlier study (as part of this programme) will be road-tested in 3-5 screening units. Men will be interviewed to find out what they think of the new screening and how their screening results might change their behaviour. Women will be included to find out if they would also benefit from PAD and BP screening. The information from the women will be used to apply for research funds to test screening for women as part of a larger, national screening programme for women.

Who can participate?

1. Men aged 65 years invited for AAA and AAA + PAD + BP screening

2. Women aged 60 to 69 years invited for PAD + BP screening
3. Screeners within the UK NHS AAA Screening Programmes and the Pilot PAD + BP Screening Programme

What does the study involve?

Participants complete a questionnaire before screening, a brief questionnaire after screening, a minimal set of questions at the 5-week phone call, and repeat questionnaires at 3, 6 and 12 months and annually up until 5 years. Data is collected from routine electronic health records and hospital/GP records. Participants are asked to wear a GENEActiv activity watch-like device at Leicester only.

Participants attend interviews where the questions will seek to find out: a) their experience and how acceptable they found the different parts of the screening process; b) the psychosocial impact of screening (i.e. the relationship between social factors and thoughts and behaviours); c) their thoughts and actions following a positive screening result for PAD and/or high blood pressure (where relevant) and the likelihood that they think they will engage with cardiovascular risk management strategies. Screeners will give their thoughts on the screening programme.

What are the possible benefits and risks of participating?

Giving up some time to complete the questionnaires and interviews (where relevant). Some people may find discussing thoughts and feelings as part of the interview distressing. The researchers cannot guarantee any direct benefit from participating, but those participating will be helping to make a significant contribution to the research of new NHS screening programmes.

Where is the study run from?

Leicester Clinical Trials Unit (UK)

When is the study starting and how long is it expected to run for?

January 2021 to September 2028

Who is funding the study?

National Institute for Health Research (NIHR) (UK)

Who is the main contact?

Dr Sian Baldock
phast@leicester.ac.uk

Contact information

Type(s)

Public

Contact name

Dr Sian Baldock

ORCID ID

<https://orcid.org/0000-0001-9231-6114>

Contact details

Leicester Clinical Trials Unit
College of Life Sciences
University of Leicester

Maurice Shock Building
University Road
Leicester
United Kingdom
LE1 7RH
+44 (0)116 229 7246
phast@leicester.ac.uk

Type(s)

Principal investigator

Contact name

Prof Matthew Bown

ORCID ID

<https://orcid.org/0000-0002-6180-3611>

Contact details

Department of Cardiovascular Sciences
University of Leicester
BHF Cardiovascular Research Centre
Glenfield General Hospital
Leicester
United Kingdom
LE2 7LX
+44 (0)116 252 3190
mjb42@leicester.ac.uk

Type(s)

Scientific

Contact name

Prof Matthew Bown

Contact details

Department of Cardiovascular Sciences
University of Leicester
BHF Cardiovascular Research Centre
Glenfield General Hospital
Leicester
United Kingdom
LE2 7LX
+44 (0)116 252 3190
mjb42@leicester.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

299928

Protocol serial number

IRAS 299928, CPMS 51100, UoL 0838

Study information

Scientific Title

Peripheral arterial disease, High blood pressure and Aneurysm Screening Trial: a Feasibility study to determine acceptability, uptake and effect of combined peripheral arterial disease, high blood pressure and abdominal aortic aneurysm screening (PHAST-F)

Acronym

PHAST-F

Study objectives

The primary hypothesis to be tested is that screening men for peripheral arterial disease (PAD), high blood pressure (BP) and abdominal aortic aneurysm (AAA) contemporaneously improves health, is acceptable and is cost-effective. WP2 will specifically look at the acceptability, uptake and effect of combined screening.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 09/12/2021, North of Scotland (2) Research Ethics Committee (Summerfield House, 2 Eday Road, Aberdeen, AB15 6RE, UK; +44 (0)1224 558458; gram.nosres@nhs.scot), REC ref: 202121/NS/0147

Study design

Observational cross-sectional multicentre feasibility study

Primary study design

Observational

Study type(s)

Screening

Health condition(s) or problem(s) studied

Observation of the impact of PAD and BP screening on CV risk uptake in men and women being screened

Interventions

Observation of a pilot PAD and BP screening programme vs AAA screening in men regarding the impact on attendance. Post-screening (i.e. a non-research activity), men and women in follow up (research activity) will complete questionnaires assessing quality of life, change in medical history, anxiety, activity, healthcare resource, and psychosocial impact.

At attendance for either AAA screening or AAA plus PAD and high BP screening, subjects will be invited to participate in the following studies (as applicable).

Healthcare professionals in primary care will also be interviewed as well as the screeners and screening staff managers.

Please note: PAD+BP screening is part of a Pilot Screening Programme not the research study, due to how screening programmes are run and the primary outcome being attendance for screening.

The follow-up study:

1. At enrolment to the follow-up study, participants will complete a baseline questionnaire
2. After screening, participants will complete a post-screening questionnaire
3. If screened for PAD and high BP and they screened positive for either, participants will be phoned at 5 weeks and asked a further set of questions on their engagement with cardiovascular risk management strategies and whether they have booked an appointment with their GP
4. Participants will be sent a follow-up questionnaire to complete at 3 months, 6 months, 12 months and annually until 5 years. This is the end of the follow-up.
5. Follow-up using electronic health records over 5 years. The end of follow-up is after 5 years.

The qualitative interview study (optional extra on AAA+PAD+BP screening days):

1. After screening, those enrolled in the optional extra qualitative interview study will be interviewed shortly after screening
2. For those that screened positive for PAD and/or high BP, this interview will be repeated at 6 months
3. After screening, the screeners and managers will be interviewed. This interview will be repeated at 3 months. Primary care staff will also be interviewed.

The GENEActiv sub-study (optional extra on AAA+PAD+BP screening days, UHL site only):

1. After screening, those enrolled in the optional extra GENEActiv sub-study and screened positive for PAD and/or high BP will be issued with a GENEActiv activity device monitor (if available) to wear for 7 days and complete an activity diary.
2. The device will be re-issued at 6 months for a repeat reading over a 7-day period.

Intervention Type

Other

Primary outcome(s)

1. Attendance measured as part of the Pilot Screening Programme using the UK AAA Screening Programme methods at baseline for all individuals invited, not just participants in the follow-up, interview, and GENEActiv studies
2. Quality of life measured using EQ-5D-5L at baseline, 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study

Key secondary outcome(s)

1. Prevalence of PAD and high BP measured using a questionnaire at baseline as part of the Pilot Screening Programme and baseline, 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study
2. Prevalence of undiagnosed PAD and high BP measured using a questionnaire at baseline, 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study

3. Prevalence of untreated PAD and high BP measured using a questionnaire at baseline, 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study.
4. Acceptability of screening test measured by a question on a scale of 1 to 10 after screening as part of follow-up study. Acceptability of the screening is also assessed by interview immediately after screening for those that screen positive for PAD and/or high BP. Acceptability is also assessed by interview after 3 months for the managers and again for those that screen positive for PAD and/or high BP at 6 months.
5. Training requirements for AAA+PAD+BP screening determined from the qualitative interviews with screening staff and managers after screening, 3 months, and throughout screening activities and following clinical training under the pilot screening programme
6. Additional time required for PAD+BP screening vs AAA screening in men measured by collecting the start and end times of appointments at the screening appointment as part of the service evaluation of the clinical pilot screening programme
7. Uptake of CV risk management in men and women that screen positive for PAD and/or high BP, measured by questionnaire (including IPAQ) at 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study and from activity data collected by wearing the GENEActiv activity device monitor at baseline and 6 months. Uptake of CV risk management will also be assessed by interviews with those that screen positive for PAD and/or high BP after 6 months following a positive screen result.
8. Major adverse CV events (MACE [death/admission from/for angina/myocardial infarction /ischaemic heart disease/stroke]) assessed by questionnaire at 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study and as part of routine data collection via NHS Digital or contacting the participant's GP at 1 year, 2 years, 3 years, 4 years, and 5 years.
9. Major adverse limb events (MALE [death/admission from/for lower limb revascularization /major limb amputation]) assessed by questionnaire at 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study and as part of routine data collection via NHS Digital or contacting the participant's GP at 1 year, 2 years, 3 years, 4 years, and 5 years
10. Psychosocial impact of screening assessed by interview immediately after screening and again at 6 months for those that screen positive for PAD and/or high BP. It will also be assessed by questionnaire (GAD-7) at 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study.
11. Health care resource use and cost effectiveness assessed by questionnaire at 3 months, 6 months, 12 months, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study and as part of routine data collection via NHS Digital or contacting the participant's GP at 1 year, 2 years, 3 years, 4 years, and 5 years as part of the follow-up study.

Completion date

01/09/2028

Eligibility

Key inclusion criteria

Men's study:

1. Willing and able to give informed consent
2. Male
3. Aged 65 years or in the year of their 65th birthday
4. In receipt of an invitation to primary routine AAA screening and have attended for screening
5. An ability to understand verbal and written English

Women's study:

1. Willing and able to give informed consent
2. Female
3. Aged 60 to 69 years
4. Registered at the Willows Health group of GP practices under University Hospitals of Leicester (UHL)
5. An ability to understand verbal and written English

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

60 years

Upper age limit

69 years

Sex

All

Total final enrolment

1224

Key exclusion criteria

Men's study:

1. Men being re-invited for a second appointment having not attended for their primary screening appointment (unless involvement with the study specifically requested upon re-invitation)

Date of first enrolment

17/02/2022

Date of final enrolment

30/09/2024

Locations

Countries of recruitment

United Kingdom

England

Scotland

Study participating centre
University Hospitals of Leicester NHS Trust
Leicester Royal Infirmary
Infirmary Square
Leicester
England
LE1 5WW

Study participating centre
University Hospitals Coventry and Warwickshire NHS Trust
Walsgrave General Hospital
Clifford Bridge Road
Coventry
England
CV2 2DX

Study participating centre
NHS Greater Glasgow and Clyde
J B Russell House
Gartnavel Royal Hospital
1055 Great Western Road Glasgow
Glasgow
Scotland
G12 0XH

Study participating centre
Pilgrim Hospital
Sibsey Road
Boston
England
PE21 9QS

Sponsor information

Organisation
University of Leicester

ROR
<https://ror.org/04h699437>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Men single screening version 1.4	09/12/2021	14/03/2022	No	Yes
Participant information sheet	Men triple screening version 1.4	09/12/2021	14/03/2022	No	Yes
Participant information sheet	Women version 1.4	09/12/2021	14/03/2022	No	Yes