

Effectiveness of pink noise on reducing pain and anxiety during primary molars pulpotomy

Submission date 02/08/2025	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 04/08/2025	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 04/08/2025	Condition category Oral Health	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Pink noise has more energy in lower frequencies, making it sound deeper and more balanced, similar to natural sounds. Examples include the sound of crashing waves, falling rain, and rustling leaves. Pink noise improves sleep quality, aids relaxation, treats tinnitus, reduces stress, enhances comfort in office environments, boosts productivity, and decreases anxiety during dental procedures.

Another type of auditory stimulus used in clinical settings is children's songs, which serve as preferred music. These songs modulate pain-related activity in the brain's anterior cingulate cortex, and effectively raise pain thresholds by increasing endorphin production. Listening to music for 10 minutes is a sufficient duration to allow its anxiety-reducing effects to take hold. Studies suggest that waiting in dental clinics while listening to music is equally effective in lowering anxiety levels compared to taking benzodiazepines. The current study aimed to evaluate and compare the effects of pink noise and children's songs on anxiety and pain in children aged 6–9 years during pulpotomy procedures. The null hypothesis is that there is no significant difference in the reduction of anxiety and pain levels during pulpotomy procedures in children aged 6–9 years when exposed to pink noise compared to children's songs.

Who can participate?

1. Children aged 6–9 years.
2. Classified as "definitely positive" or "positive" on Frankel's behavior rating scale.
3. Requiring pulpotomy treatment in mandibular primary molars.

What does the study involve?

Following screening against inclusion criteria, 75 eligible patients were enrolled and randomly allocated into three study groups:

- Group 1: Pink noise (n = 25).
- Group 2: Children's songs (n = 25).
- Group 3: Control group, tell-show-do (TSD) technique (n = 25).

What are the possible benefits and risks of participating?

Possible benefits are: Performing non-urgent dental treatment in the mandibular arch.

Possible risk is: Injection will be painful if the device is not effective.

Where is the study run from?
Damascus University (Syria)

When is the study starting and how long is it expected to run for?
October 2024 and December 2024

Who is funding the study?
Damascus University (Syria)

Who is the main contact?
Dr Mawia Karkoutly, mawia95.karkoutly@damascusuniversity.edu.sy,
mawiamaherkarkoutly@hotmail.com

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Dr Mawia Karkoutly

ORCID ID

<https://orcid.org/0000-0003-0227-1560>

Contact details

Mazzeh

Damascus

Syria

-

+963 992647528

Mawia95.karkoutly@damascusuniversity.edu.sy

Additional identifiers

Study information

Scientific Title

Effectiveness of pink noise on reducing pain and anxiety during primary molars pulpotomy compared with music: a randomized clinical trial

Study objectives

The current study aimed to evaluate and compare the effects of pink noise and children's songs on anxiety and pain in children aged 6–9 years during pulpotomy procedures.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 27/05/2024, The Biomedical Research Ethics Committee (BMREC) of Damascus University (Mazzeh, Damascus, -, Syria; +963 992647528; info@damascusuniversity.edu.sy), ref: N3551/2024

Study design

Randomized controlled trial with a double-blinded three-arm parallel-group

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Dental pain and anxiety

Interventions

Baseline anxiety levels were evaluated by measuring pulse rate and administering the FI scale. Children were shown two sets of five facial expressions and instructed to choose the image that best represented their current anxiety state. Children in the pink noise group listened using headphones. In the children's songs group, the child was asked to choose one of the songs from their favorite program and cartoon movies on YouTube, so that it would be familiar to them to listen to. The music or pink sound continued throughout the session through headphones. In the TSD technique group, the procedure was first explained verbally and then demonstrated in a child-friendly manner before treatment initiation. The "Do" phase was implemented during actual treatment without altering the initial explanation. Pulse rate was recorded after conditioning the child using the TSD technique or after listening to the songs or pink noise for 10 minutes.

The conventional pulpotomy procedure is based on AAPD guidelines. Following IANB administration, pulse rate, FLACC behavioral pain scores, and FI scale ratings were documented. Pulse rate and FLACC behavioral pain scores were repeated after rubber dam placement and again following cavity preparation and pulp amputation, including during hemorrhage control with a moist cotton pellet. Final pulse rate and FI scale assessments were conducted upon completion.

Intervention Type

Other

Primary outcome(s)

1. Pulse rate was objectively measured as an indicator of dental anxiety using a finger pulse oximeter (Alpha, Prolinx GmbH, Düsseldorf, Germany). Measurements were recorded at six critical time points: Baseline (t0). Following patient conditioning before treatment initiation (t1). Immediately after IANB administration (t2). Post rubber dam placement (t3). Following pulp amputation (t4). Upon treatment completion (t5)

2. The Face, Legs, Activity, Cry, and Consolability (FLACC) Behavioral Pain Assessment Scale (FLACC) behavioral pain scale, a nonverbal pain assessment tool, was recorded at three critical time points: Immediately after IANB administration (t2). Post rubber dam placement (t3). Following pulp amputation (t4)

Key secondary outcome(s)

Subjective anxiety levels were assessed using a 5-point pictorial scale (Facial Image (FI) Scale) with facial expressions ranging from very happy (score = 1) to very fearful (score = 5). The children were asked to choose the face that best matched their emotional state at three-time points: Baseline (t0), immediately after IANB administration (t2), and following pulp amputation (t4)

Completion date

17/12/2024

Eligibility

Key inclusion criteria

1. Children aged 6–9 years.
2. Classified as "definitely positive" or "positive" on Frankel's behavior rating scale.
3. Requiring pulpotomy treatment in mandibular primary molars.

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Child

Lower age limit

6 years

Upper age limit

9 years

Sex

All

Total final enrolment

75

Key exclusion criteria

1. Children with oral, mental, and/or systemic conditions that could affect treatment outcomes.
2. Children who had taken sedative or analgesic medications within the 24 hours preceding the dental examination.

Date of first enrolment

03/10/2024

Date of final enrolment

28/11/2024

Locations

Countries of recruitment

Syria

Study participating centre

Damascus University

Mazzeh

Damascus

Syria

-

Sponsor information

Organisation

Damascus University

ROR

<https://ror.org/03m098d13>

Funder(s)

Funder type

Not defined

Funder Name

Damascus University

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from mawia95.karkoutly@damascusuniversity.edu.sy, mawiamaherkarkoutly@hotmail.com

The type of data that will be shared: Not currently known

Timing for availability: Upon a reasonable request

Whether consent from participants was required and obtained: Informed consent was obtained

IPD sharing plan summary

Available on request