

A single-centre study looking at the effect of compression therapy in patients having foam sclerosant injection for varicose veins

Submission date 11/04/2016	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 20/04/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 07/05/2025	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Varicose veins are a common condition, in which the veins (usually on the legs) become swollen and enlarged. It happens when the small valves inside the veins which are designed to make sure blood is travelling in the right direction (back towards the heart) stop working properly, causing blood to leak backwards and collect in the vein. Over the past decade, new minimally invasive treatments ('key hole' procedures) have been introduced. A common type of procedure used is ultrasound guided foam sclerotherapy. This is where special foam is injected into the veins (a process guided using ultrasound scanning) which seal the damaged veins closed. This type of procedure is becoming more and more popular as they are associated with fewer complications and a quicker recovery than traditional surgery (to remove the veins). Despite advances in surgical techniques, there is still uncertainty as to the role of compression therapy (wearing tight stockings which squeeze the leg to improve circulation) in the management of varicose veins. The American Venous Forum (an American veins organisation) recommends the use of compression stockings to reduce the risk of clot (haematoma) formation, pain and swelling. The NICE Guidelines have recommended further research into the use of compression stockings as post-procedure intervention because of this lack of evidence. The aim of this study is to compare the effect of compression therapy following foam sclerotherapy.

Who can participate?

Adults with varicose veins suitable for foam sclerotherapy.

What does the study involve?

Participants in both groups undergo foam sclerotherapy of their varicose veins, before being randomly allocated to one of two groups. Those in the first group wear bandages for 24 hours and receive no compression therapy. Those in the second group are asked to wear compression stockings for 1 week following foam sclerotherapy. All participants are given a diary card to rate their pain levels every day for 10 days after their surgery. They are then received two weeks after the procedure in order to fill in questionnaires about their quality of life and undergo a

physical examination to see if there is there is any skin discolouration or inflammation present. Six months after the procedure, participants attend another clinic appointment where they have an ultrasound scan to find out if the surgery has worked and fill in further questionnaires.

What are the possible benefits and risks of participating?
Not provided at time of registration

Where is the study run from?
Charing Cross Hospital (UK)

When is the study starting and how long is it expected to run for?
June 2015 to December 2017

Who is funding the study?
Chemische Fabrik Kreussler & Co. GmbH (Germany)

Who is the main contact?
Mr Roshan Bootun
r.bootun@imperial.ac.uk

Contact information

Type(s)
Public

Contact name
Mr Roshan Bootun

Contact details
Vascular Research Office
Floor 4 North
Charing Cross Hospital
Imperial College London
Fulham Palace Road
London
United Kingdom
W6 8RF
+44 20 3311 7335
r.bootun@imperial.ac.uk

Type(s)
Scientific

Contact name
Dr Amjad Belramman

ORCID ID
<https://orcid.org/0000-0003-2057-6755>

Contact details

Four North,
Charing Cross Hospital
Fulham Palace Road
London
United Kingdom
W6 8RF
None provided
a.belramman17@imperial.ac.uk

Additional identifiers

ClinicalTrials.gov (NCT)
NCT02655406

Protocol serial number
20419

Study information

Scientific Title

A randomised controlled trial of compression therapy following foam sclerotherapy of varicose vei

Study objectives

Compression therapy following foam sclerotherapy more effective than foam sclerotherapy alone.

Ethics approval required

Old ethics approval format

Ethics approval(s)

National Research Ethics Service (NRES) Committee North East - Newcastle & North Tyneside 1, 09/09/2015, ref: 15/NE/0314

Study design

Randomized; Interventional; Design type: Treatment, Device

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Varicose veins

Interventions

Patients will be randomised to have compression (group A) or no compression (group B). This will be done by using an online randomisation software (Sealed Envelope Ltd).

The compression therapy used will be Class II compression stockings. The treatment offered will be foam sclerotherapy of varicose veins using either Polidocanol or Sodium Tetradecyl Sulphate (STS) with the concentration used left to the discretion of the clinician.

Group A: Participants will be asked to wear compression stockings for 1 week following foam sclerotherapy.

Group B: Participants will be provided with bandages to wear for 24 hours only, with no further compression afterwards.

Patients in both groups will be followed-up at 2 weeks and 6 months.

Intervention Type

Other

Primary outcome(s)

Pain is measured using the visual analogue scale (VAS) at baseline and daily for the first 10 days post procedure.

Key secondary outcome(s)

1. Quality of life is measured using EQ-5D, AVVQ and CIVIQ at baseline, 2 weeks and 6 months
2. Clinical change is assessed using the VCSS at baseline, 2 weeks and 6 months
3. Degree of phlebitis is assessed at 2 weeks and 6 months
4. Degree of skin discolouration and matting is assessed at 2 weeks and 6 months
5. Patient compliance with the intervention
6. Time taken to return to work and normal activities
7. Occlusion rates at 6 months
8. Cost effectiveness of the intervention

Completion date

31/01/2021

Eligibility

Key inclusion criteria

1. Adults over 18 years of age
2. Varicose veins suitable for foam sclerotherapy

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

139

Key exclusion criteria

1. Allergic to sclerosant
2. Current DVT
3. Arterial disease (ABPI<0.8)
4. Patients unable to wear compression stockings
5. Patient who are unwilling to participate
6. Inability or unwillingness to complete questionnaires
7. Inability to attend follow-up appointments

Date of first enrolment

10/01/2016

Date of final enrolment

31/01/2020

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Charing Cross Hospital

Imperial College London

Fulham Palace Road

London

United Kingdom

W6 8RF

Sponsor information

Organisation

Imperial College London

ROR

<https://ror.org/041kmwe10>

Funder(s)

Funder type

Industry

Funder Name

Chemische Fabrik Kreussler & Co. GmbH

Results and Publications

Individual participant data (IPD) sharing plan

Anonymised data is available on request by researchers after the publication of the manuscript from Prof. Alun Davies (a.h.davies@imperial.ac.uk) or Mr Tristan Lane (tristan.lane@imperial.ac.uk). It will be uploaded to a secure data repository for ease of access. Consent for anonymised analysis of data and consent for publication has been ethically approved and prospectively sought from participants. This data will be stored for 10 years in line with Imperial College London guidelines and will be available from that time. All data should be referenced and cited appropriately.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/03/2024	07/05/2025	Yes	No
Basic results	version 23	06/11/2023	06/11/2023	No	No
HRA research summary			28/06/2023	No	No
Protocol file	version 3.1	04/05/2017	18/10/2022	No	No