

Investigating tranexamic acid for bleeding control in facelift surgery

Submission date 01/11/2025	Recruitment status Recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 06/11/2025	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 06/11/2025	Condition category Surgery	<input type="checkbox"/> Individual participant data
		<input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

This randomized controlled trial investigates whether local tranexamic acid (TXA) can reduce intraoperative bleeding and postoperative complications (like hematoma, swelling, and bruising) in patients undergoing facelift (rhytidectomy) surgery at King Abdulaziz University Hospital. TXA is known for minimizing bleeding in surgery, but no RCT has evaluated its local use in facelifts.

Who can participate?

Healthy adults aged 35 years and older, or those with stable (compensated) medical conditions, who are undergoing facelift surgery.

People are excluded if they are under 35, have bleeding or clotting disorders, take anticoagulants or hormones, or have major organ failure.

What does the study involve?

Participants are randomly assigned to either:

Intervention group: receive local TXA under the skin flap, or

Control group: receive standard care (lidocaine with epinephrine only).

Bleeding is measured during surgery, and participants attend three follow-up visits (days 1, 6, and 9) and one remote check. Surgeons and patients rate bruising and swelling at each visit.

What are the possible benefits and risks of participating?

Benefits: TXA may reduce bleeding, bruising, swelling, and overall complications, leading to faster recovery.

Risks: Rare allergic or clotting reactions, nausea, low blood pressure, rash, or visual disturbances. Pregnant or breastfeeding individuals must not participate.

Where is the study run from?

Conducted at King Abdulaziz University Hospital (KAUH), Riyadh, Saudi Arabia, under King Saud University supervision.

When is the study starting and how long is it expected to run for?

February 2024 to June 2026

Who is funding the study?
Investigator initiated and funded

Who is the main contact?
Principal Investigator:
Prof. Ahmad Al-Arfaj, aalarrfaj@ksu.edu.sa

Contact information

Type(s)
Scientific

Contact name
Dr Abdulaziz AlEnazi

ORCID ID
<https://orcid.org/0000-0002-0270-1906>

Contact details
SB-ORLHNS, EBEORL-HNS, ABHS-ORLHNS, JMC-ORL,DOHNS
Otolaryngology Head Neck & Facial Plastic Surgery,
Imam Abdulrahman Bin Faisal University.
Al Khobar
Saudi Arabia
34212
-
asenazi@iau.edu.sa

Type(s)
Principal investigator

Contact name
Prof Ahmed Alarfaj

ORCID ID
<https://orcid.org/0009-0009-5436-3660>

Contact details
Facial Plastic Unit
King Saud University, Riyadh KAUH, ENT Department
KSU Facial Plastic Fellowship
Riyadh
Saudi Arabia
12629
-
aalarrfaj@ksu.edu.sa

Type(s)
Public

Contact name

Dr Abdulaziz Alderaywsh

ORCID ID

<https://orcid.org/0000-0001-7327-5708>

Contact details

Family and Community Medicine King Saud University

Riyadh

Saudi Arabia

12372

-

aalderaywsh@ksu.edu.sa

Additional identifiers**Clinical Trials Information System (CTIS)**

Nil known

Protocol serial number

E-23-8441

Study information**Scientific Title**

Effects of local tranexamic acid on bleeding during rhytidectomy and post operative complication: randomized, controlled, single-blind trial

Acronym

TXA-Rhytid Trial

Study objectives

Determine whether Local TXA has any effect on intraoperative bleeding or postoperative sequelae in patients undergoing a facelift.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 19/02/2024, College of Medicine Institutional Review Board (King Saud University – College of Medicine, Riyadh, 11472, Saudi Arabia; +966 114670011; irbresearch2@gmail.com), ref: E-23-8441

Study design

Interventional single blind randomized controlled trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Prevention of bleeding and postoperative complication in Rhytidectomy

Interventions

Control: Tumescent (0.5% lidocaine with 1:200,000 epinephrine).

Case: Tranexamic acid (TXA) +Tumescent (1 mg/ml TXA and 0.5% lidocaine with 1:200,000 epinephrine)

Administration Rout: Local Infiltration

Randomization of allocation: by using random numbers table generated in the Statistical Package for Social Studies (SPSS v. 21) (SPSS v. 21, IBM Corp., New York, NY, USA)

The total duration of treatment and follow-up will be 1 month

Intervention Type

Drug

Phase

Phase III/IV

Drug/device/biological/vaccine name(s)

Tranexamic acid

Primary outcome(s)

Certainly, Nicholas. Here's the revised list of outcome measures in the correct format:

1. Intraoperative blood loss is measured using the volume collected in the surgical suction container during surgery
2. Intraoperative blood loss is measured using gravimetric analysis by weighing surgical sponges and gauze before and after use during surgery
3. Postoperative edema is measured using a subjective 3-point scale (mild, moderate, severe) assessed by both patient and surgeon at postoperative visits 1, 6, and 9
4. Postoperative ecchymosis is measured using a subjective 3-point scale (mild, moderate, severe) assessed by both patient and surgeon at postoperative visits 1, 6, and 9
5. Postoperative edema is measured using objective volumetric assessment with Vectra imaging at postoperative visits 1, 6, and 9
6. Postoperative ecchymosis is measured using objective volumetric assessment with Vectra imaging at postoperative visits 1, 6, and 9

Key secondary outcome(s)

1. Patient satisfaction is measured using the FACE-Q questionnaire at baseline (before procedure) and postoperative day 9
2. Time to drain removal is measured using clinical records at end of study
3. Complications will be assessed at follow up visits:
 - Skin necrosis
 - Hematoma
 - Thromboembolic events
 - Infection

Completion date

01/06/2026

Eligibility

Key inclusion criteria

1. \geq 35-year-old
2. Capable for follow up
3. Healthy patients or with compensated comorbidities, after the agreement of the assigned health care team
4. Candidates for cosmetic face and neck surgery
5. Absence of previous surgeries on the face

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

35 years

Upper age limit

75 years

Sex

All

Key exclusion criteria

1. $<$ 35-year-old
2. History of thromboembolic events or seizure
3. History of bleeding disorder
4. Active use of oral contraceptive pills and/or hormone replacement therapy, Aspirin, anti-coagulant, and Omega 3
5. Patient preference
6. Any condition increase edema (Renal failure, heart failure, liver disease)

Date of first enrolment

01/05/2024

Date of final enrolment

01/05/2026

Locations

Countries of recruitment

Saudi Arabia

Study participating centre
King Abdulaziz University Hospital, King Saud University
King Abdulaziz Rd, Al Malaz
Riyadh
Saudi Arabia
12629

Sponsor information

Organisation
King Saud University

ROR
<https://ror.org/02f81g417>

Funder(s)

Funder type
Other

Funder Name
Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date