

A trial of suction for the early resolution of a collapsed lung

| | | |
|----------------------------------------|---------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------|
| Submission date 24/11/2022 | Recruitment status No longer recruiting | <input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol |
| Registration date 07/12/2022 | Overall study status Ongoing | <input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results |
| Last Edited 24/02/2026 | Condition category Respiratory | <input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year |

Plain English summary of protocol

Background and study aims

A pneumothorax occurs when air gets into the space between the lung and the chest wall, usually through a small hole in the lung. This causes the lung to collapse and can occur "spontaneously", meaning without an injury to the chest. Primary spontaneous pneumothorax (PSP) occurs in younger patients without known lung disease. Most patients with PSP need to have a tube (or drain) inserted into the chest to remove the air between the lung and chest wall. This allows the lung to re-inflate. The tube is attached to a bottle with water in it, creating an underwater seal, and air bubbles out through the water.

Although some patients with PSP can be treated at home (either by not draining the chest or with a home drainage device), at least 50% of patients stay in the hospital waiting for the lung to re-inflate for 4 to 8 days. In these patients, it is possible to provide suction (negative pressure) to the drain with the aim of expanding the lung more quickly and reducing time in the hospital. However, we do not know if using suction is helpful, or if it has risks. There have been no large studies conducted to prove whether suction is effective in reducing treatment time. The current guidelines provide conflicting advice on the routine use of suction, but despite this, doctors often use it. We want to address this question directly. Reducing treatment time is important because interviews and questionnaires conducted with patients who have had a pneumothorax have told us that their top priorities include reducing the amount of time that they have a chest tube and the length of their hospital stay. In addition, treating these patients in hospital costs the NHS around £7.2m per year, so safely reducing the time spent in the hospital will be cost-saving for the NHS.

Who can participate?

Patients aged 16* to 50 years old with PSP being treated at 36 centres in hospitals around the UK. *Common law presumes that young people aged between 16 and 18 years old are usually competent to give consent to treatment and consent from those with parental responsibility is not legally necessary. Eligible young persons believed to be competent by the PI or delegate should be approached about the study. The involvement of parents in decision-making should be encouraged unless the young person objects.

What does the study involve?

Patients will be randomly assigned to either have suction applied to their chest tube or treated with usual care (no suction) and followed up for 6 months in total.

What are the possible benefits and risks of participating?

In terms of benefits, those randomly assigned to suction may resolve their pneumothorax more quickly and hence a shorter hospital stay. When the study is complete, we hope the information collected will help to improve the treatment of patients in the future. Once admitted to the hospital with a chest drain, management of ongoing pneumothorax remains contentious. The rationale behind the use of suction (the application of negative pressure to the pleural space via the chest tube) is that the lung will expand more quickly and potentially heal more quickly once the lung is re-expanded and in contact with the parietal pleura. However, suction is not without risk, as it may precipitate injury to the lung (by too rapidly expanding the lung, or may result in delayed healing of the pneumothorax if the application of suction maintains flow through the hole in the lung. Patients will have daily chest x-rays (CXRs) during their admission as per standard care. There are theoretical health risks from excessive radiation exposure, but CXRs are the safest tests (the radiation dose is only equivalent to around four days' worth of normal background radiation). No additional radiological investigations will be undertaken for research purposes. All patients will be carefully monitored for safety outcomes in both arms of the trial, with adverse events reported and reviewed according to the Oxford Respiratory Trials Units (ORTU) standard operating procedures.

Where is the study run from?

Oxford Respiratory Trials Unit, University of Oxford (UK)

When is the study starting and how long is it expected to run for?

May 2022 to February 2027

Who is funding the study?

National Institute for Health and Care Research - Health Technology Assessment (NIHR-HTA) (UK)

Who is the main contact?

Dr Rob Hallifax, Robert.Hallifax@ndm.ox.ac.uk (UK)

Contact information

Type(s)

Public

Contact name

Mrs Emma Hedley

Contact details

Senior Clinical Trial Manager

Churchill Hospital

Old Road

Headington

Oxford

United Kingdom

OX3 7LE

+44 (0)1865 282952
emma.hedley@ndm.ox.ac.uk

Type(s)

Principal investigator

Contact name

Dr Robert Hallifax

ORCID ID

<https://orcid.org/0000-0002-9467-668X>

Contact details

Academic Clinical Lecturer
University of Oxford
Oxford Respiratory Trials Unit
Oxford Centre for Respiratory Medicine
Churchill Hospital
Old Road
Oxford
United Kingdom
OX3 7LE
+44 (0)1865857104
Robert.Hallifax@ndm.ox.ac.uk

Type(s)

Scientific

Contact name

Dr Robert Hallifax

Contact details

Academic Clinical Lecturer
University of Oxford
Oxford Respiratory Trials Unit
Oxford Centre for Respiratory Medicine
Churchill Hospital
Old Road
Oxford
United Kingdom
OX3 7LE
+44 (0)1865857104
Robert.Hallifax@ndm.ox.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

316434

Protocol serial number

CPMS 54383, IRAS 316434

Study information

Scientific Title

Randomised trial of suction for primary pneumothorax early resolution

Acronym

RASPER

Study objectives

It is hypothesised that using suction to treat people in the hospital with a lung collapse is safe and can shorten the time that people need to have a chest tube in place.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 30/11/2022, West Midlands, Solihull (Meeting held by video-conference via Zoom; +44 (0)207 104 8191, (0)207 104 8269; solihull.rec@hra.nhs.uk) ref: 22/WM/0253

Study design

Multi-centre open-label randomized controlled study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Primary spontaneous pneumothorax

Interventions

RASPER is a multi-centre open-label randomised controlled trial of patients with primary spontaneous pneumothorax (PSP) requiring treatment in the hospital. The aim of the study is to assess the superiority of suction versus standard care with respect to total treatment duration. The main hypothesis is that treatment duration will be significantly reduced by the use of suction, as compared to standard care. Suction devices are currently in clinical use in the UK but are used inconsistently without any clear evidence of benefit. The use of suction has not been robustly tested in a large clinical trial, compared to standard management, as we propose here. 450 patients with PSP requiring chest drainage in the hospital will be enrolled from 36 UK centres, including our established pneumothorax network. Patients will be screened from respiratory and general medical wards. Data will be collected from routinely recorded clinical data and patient-completed questions on pain and breathlessness scores and overall health (EQ5D). Other important factors being assessed include safety, the level of comfort of the device, and the cost-effectiveness of the use of suction.

Patients requiring treatment in a hospital with a chest drain in place (to re-inflate the lung) will be invited to take part in the randomised control trial: and randomly assigned to one of two groups: suction arm or standard care (no suction).

Baseline.

At baseline patients will have a chest-x ray or CT (rarely) to confirm their diagnosis. As part of the study, patients will have their demographics/medical history taken, they will be asked to complete a questionnaire (EQ-5D-5L) and another one about pain and breathlessness (VAS assessment).

1. Intervention arm: Suction

The suction will be incrementally increased as per a trial-specific procedure: starting at -1.0kPa (-10cm H₂O, -7.5mmHg), increasing to -1.5kPa and then -2.0kPa as tolerated, every 2-4 hours). Suction should be reduced on the basis of specific criteria (pain, complications). Adherence will be documented on daily clinical review CRFs.

2. Control arm: Usual care

Patients in the usual care will be managed as standard (as per current national guidelines) without the use of suction (i.e. connected to an underwater seal bottle) unless suction is clinically needed for safety reasons. For example, rarely the air leak from the lung is too large to be drained by the chest drain and the pneumothorax continues to get larger despite the drain working – in this situation suction will be allowed for safety reasons (and recorded in the trial records).

Follow-up

Patients will be followed up 14 days (+/- 3 days) in person after completion of treatment, at 30 days (+ up to 7 days) in person/over the phone and 6 months (+/- 2 weeks) after enrolment by telephone. Patients will be asked to complete an online questionnaire, about pain and breathlessness, and tell us if they had any contact with healthcare providers since their last visit (for health economic analysis).

Minimising Bias

Due to the nature of the interventions, patients and clinicians cannot be blinded to allocation and therefore code-breaking is not needed for this trial. However, criteria will be specified for “chest drain removal” and “surgical referral/failure of medical treatment” to provide objective data that will be blind reviewed after the trial by an independent assessor blind to the treatment arm (i.e. objective blind outcome assessment) and compared to actual treatment duration at study end.

Interim Monitoring and Analyses

A blinded interim analysis of the primary outcome (hospital stay) will be undertaken after approximately 50% of patients have been recruited in order to assess the assumptions made in the sample size calculation. This analysis will be reviewed by the DSMC which will make recommendations regarding any necessary changes to the sample size required. No correction of the significance level of the final analysis is planned on this single assessment of the early event rate by the DSMC.

Patient and public involvement

We conducted a survey of twelve patients to find out what was most important to them and designed this trial based on the results. The survey showed that patients that their top priorities are: 1. To reduce the amount of time that they have a chest tube, and 2. To reduce the length of their hospital stay. A representative from our patient group who previously had a PSP has reviewed the patient-facing documents (including the patient information sheet) and is a member of the trial committee and will represent our Patient Advisory Group.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Total treatment duration, defined as the time from randomisation to completion of pleural treatment (including surgery, if required), measured using data recorded in patient medical notes at the completion of treatment (discharge home from hospital with no drain in place)

Key secondary outcome(s)

Outcomes measured using data recorded in patient medical notes:

- 1.1. In-patient surgical rates at the completion of treatment
 - 1.2. Length of hospital stay over first the 30 days post-randomisation (including readmissions) at 30 days post-randomisation
 - 1.3. Pain and breathless scores (100mm Visual Analogue Scale (VAS)) at baseline, daily until completion of treatment, and at follow-14 days after the completion of treatment and 30 days post-randomisation
 - 1.4. EQ5D at baseline, Completion of treatment, 30 days and 6 months post-randomisation
 - 1.5. Rate of recurrence of pneumothorax at 6 months post-randomisation
2. Incremental cost per quality-adjusted life years (QALYs) gained at randomisation to 6 months measured using data gathered between those timepoints
- 3.1. Complication rates at the completion of treatment
 - 3.2. Overall number of in-patient pleural procedures at the completion of treatment
 - 3.3. Adverse events related to the use of suction at the completion of treatment

Completion date

27/02/2027

Eligibility

Key inclusion criteria

Current inclusion criteria as of 21/05/2025:

1. Participants with primary spontaneous pneumothorax (PSP) (either first or recurrent episode)
2. Male and Female aged 16* to 50 years old OR aged 51 to 60 years old without significant smoking history
3. Pneumothorax requiring chest drain in hospital (ideally within 24 hours of drain insertion, but up to 72 hours)
4. Willing and able to give written consent
5. Access to an electronic device for questionnaire completion

*Common law presumes that young people aged between 16 and 18 years old are usually competent to give consent to treatment and consent from those with parental responsibility is not legally necessary. Eligible young persons believed to be competent by the PI or delegate should be approached about the study. The involvement of parents in decision-making should be encouraged unless the young person objects.

Previous inclusion criteria:

1. Participants with primary spontaneous pneumothorax (PSP) (either first or recurrent episode)
2. Male and Female aged 16* to 50 years old

3. Pneumothorax requiring chest drain in hospital (ideally within 24 hours of drain insertion, but up to 72 hours))
4. Willing and able to give written consent
5. Access to an electronic device for questionnaire completion

*Common law presumes that young people aged between 16 and 18 years old are usually competent to give consent to treatment and consent from those with parental responsibility is not legally necessary. Eligible young persons believed to be competent by the PI or delegate should be approached about the study. The involvement of parents in decision-making should be encouraged unless the young person objects.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

16 years

Upper age limit

60 years

Sex

All

Total final enrolment

0

Key exclusion criteria

1. Known or suspected underlying lung disease**. This does not include the presence of blebs /bullae on CT chest in the absence of another specific respiratory diagnosis
2. Inability to consent or comply with trial requirements
3. Any other significant disease or disorder which, in the opinion of the Investigator, may either put the participants at risk because of participation in the study, or may influence the result of the trial, or the participant's ability to participate in the study.

***"Childhood asthma" is not considered an exclusion criterion. Patients with a diagnosis of asthma in childhood/young adulthood who do not require the use of a regular "preventer" inhaler (i.e. inhaler containing a steroid or long-acting beta-agonist), and only occasionally use a "reliever" inhaler (short-acting beta-agonist) and have never been hospitalised due to asthma remain eligible for participation in this study.

Date of first enrolment

27/02/2023

Date of final enrolment

27/02/2026

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Addenbrookes

Addenbrookes Hospital

Hills Road

Cambridge

England

CB2 0QQ

Study participating centre

John Radcliffe Hospital

Headley Way

Headington

Oxford

England

OX3 9DU

Study participating centre

Plymouth Hospital

Derriford Hospital

Derriford Road

Plymouth

England

PL6 8DH

Study participating centre

Imperial College Healthcare NHS Trust

The Bays

St Marys Hospital

South Wharf Road

London

England

W2 1BL

Study participating centre
St Thomas' Hospital
Westminster Bridge Road
London
England
SE1 7EH

Study participating centre
The Royal Victoria Infirmary
Queen Victoria Road
Newcastle upon Tyne
England
TS1 4LP

Study participating centre
Furness Hospitals NHS Trust
Furness General Hospital
Dalton Lane
Barrow-in-furness
England
LA14 4LF

Study participating centre
Princess Royal University Hospital
Farnborough Common
Orpington
England
BR6 8ND

Study participating centre
Norfolk & Norwich University Hospital
Colney Lane
Colney
Norwich
England
NR4 7UY

Study participating centre
Royal Stoke University Hospital
Newcastle Road
Stoke-on-trent

England
ST4 6QG

Study participating centre

City Hospital NHS Trust
City Hospital N H S Trust
Dudley Road
Birmingham
England
B18 7QH

Study participating centre

Queen Elizabeth Hospital
Queen Elizabeth Medical Centre
Edgbaston
Birmingham
England
B15 2TH

Study participating centre

Watford General Hospital
60 Vicarage Road
Watford
England
WD18 0HB

Study participating centre

Aneurin Bevan University Lhb
Headquarters - St Cadoc's Hospital
Lodge Road
Caerleon
Newport
Wales
NP18 3XQ

Study participating centre

Northern General Hospital
Northern General Hospital NHS Trust
C Floor, Huntsman Building
Herries Road
Sheffield

England
S5 7AU

Study participating centre

Chelsea and Westminster Hospital NHS Foundation Trust
Chelsea & Westminster Hospital
369 Fulham Road
London
England
SW10 9NH

Study participating centre

Macclesfield District General Hospital
Macclesfield District Hospital
Victoria Road
Macclesfield
England
SK10 3BL

Study participating centre

Royal Lancaster Infirmary
Ashton Road
Lancaster
England
LA1 4RP

Study participating centre

St James's University Hospital NHS Trust
St James's University Hospital
Gledow Wing
Beckett Street
Leeds
England
LS9 7TF

Study participating centre

Pinderfields General Hospital
Aberford Road
Wakefield
England
WF1 4DG

Study participating centre
Queen Elizabeth Hospital
Woolwich Stadium Road
Woolwich
London
England
SE18 4QH

Study participating centre
West Suffolk Hospital
Hardwick Lane
Bury St. Edmunds
England
IP33 2QZ

Study participating centre
The Worcestershire Royal Hospital
Newtown Road
Worcester
England
WR5 1ZL

Study participating centre
Glenfield General Hospital
Grobby Road
Leicester
England
LE3 9QP

Study participating centre
Chesterfield Royal Hospital NHS Foundation Trust
Chesterfield Road
Calow
Chesterfield
England
S44 5BL

Study participating centre

Queen Elizabeth University Hospital

1345 Govan Road
Glasgow
Scotland
G51 4TF

Study participating centre

Arrowe Park Hospital (site)

Arrowe Park Hospital
Arrowe Park Road
Wirral
England
CH49 5PE

Study participating centre

Lincoln County Hospital

Greetwell Road
Lincoln
England
LN2 5QY

Study participating centre

Ninewells Hospital

Ninewells Avenue
Dundee
Scotland
DD1 9SY

Study participating centre

Mersey and West Lancashire Teaching Hospitals NHS Trust

Whiston Hospital
Warrington Road
Prescot
England
L35 5DR

Study participating centre

The Hillingdon Hospital

Field Heath Road

Uxbridge
England
UB8 3NN

Study participating centre

Lister Hospital
Coreys Mill Lane
Stevenage
England
SG1 4AB

Study participating centre

Northwick Park Hospital
Watford Road
Harrow
England
HA1 3UJ

Study participating centre

South Tyneside District Hospital
Harton Lane
South Shields
England
NE34 0PL

Sponsor information

Organisation

University of Oxford

ROR

<https://ror.org/052gg0110>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health and Care Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Individual participant data will be available for sharing after de-identification to researchers who provide a methodologically sound proposal. The study protocol, consent forms, and participant information leaflet will also be available on request immediately after publication. Data may be shared to allow researchers to achieve the aims of their approved proposal. There is no planned end date for data sharing. Data requests should be submitted to ORTU@ndm.ox.ac.uk

IPD sharing plan summary

Available on request

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|--------------------------------------|---------|--------------|------------|----------------|-----------------|
| HRA research summary | | | 26/07/2023 | No | No |