

Acupressure combined with a flexible pressure sensor to treat coughing in patients undergoing bronchoalveolar lavage

Submission date 21/10/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 25/10/2024	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 29/10/2025	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

The study focuses on improving the quality of anesthesia recovery in patients undergoing bronchoalveolar lavage (BAL) using flexible bronchoscopy. Traditional recovery methods often involve routine monitoring, but this study explores the potential benefits of combining acupressure with flexible pressure sensor manometry. The primary aim is to investigate whether acupressure, when combined with flexible pressure sensor manometry, can enhance the recovery process post-anesthesia. Specifically, the study examines the impact on postoperative cough frequency, length of stay in the post-anesthesia care unit (PACU), incidence of stress urinary incontinence, and heart rate during the recovery period.

Who can participate?

Adult patients aged 18-64 years old who underwent BAL using flexible bronchoscopy at the First Affiliated Hospital of Zhengzhou University.

What does the study involve?

The patients were randomly divided into two groups: the acupressure group (Group A) and the control group (Group C). After the operation, all patients were transferred to the post-anesthesia care unit (PACU). In Group A, acupressure was applied to specific points (Tiantu [RN22], Chize [LU5], and Taiyuan [LU9]) after the removal of the laryngeal mask. The pressure and duration of the acupressure were measured using flexible pressure sensors. In contrast, Group C received routine PACU monitoring without acupressure.

During the recovery period in the PACU, several parameters were observed and analyzed, including:

Postoperative cough frequency (number of coughs)

Length of stay in PACU

Occurrence of stress urinary incontinence

Heart rate (HR)

These observations aimed to assess the effectiveness of acupressure combined with flexible pressure sensor manometry in improving the quality of anesthesia recovery.

What are the possible benefits and risks of participating?

Benefits: Reduces postoperative cough;Painless treatment of cough

Risks: Local skin redness phenomenon

Where is the study run from?

Department of Anesthesiology and Perioperative Medicine, First Affiliated Hospital of Zhengzhou University, China.

When is the study starting and how long is it expected to run for?

November 2023 to May 2024

Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Dr Lijun Jia, fccjialj@zzu.edu.cn

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Dr Lijun Jia

ORCID ID

<https://orcid.org/0000-0002-4650-666X>

Contact details

Department of Anesthesiology and Perioperative Medicine, First Affiliated Hospital of Zhengzhou University

Zhengzhou

China

450002

+8618737197159

fccjialj@zzu.edu.cn

Additional identifiers

Study information

Scientific Title

Effect of acupressure combined with flexible pressure sensor manometry on the quality of anesthesia recovery in patients undergoing bronchoalveolar lavage-A randomized controlled trial

Acronym

BAL

Study objectives

Acupressure combined with flexible pressure sensor manometry improves the quality of anesthesia recovery

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 26/12/2023, Research and Clinical Trial Ethics Committee of the First Affiliated Hospital of Zhengzhou University (First Affiliated Hospital of Zhengzhou University, Zhengzhou, 450052, China; +86 13783593652; 13783593652@163.com), ref: 2023-KY-0912-003

Study design

Single-center interventional single-blinded randomized controlled trial

Primary study design

Interventional

Study type(s)

Efficacy, Prevention, Safety, Treatment

Health condition(s) or problem(s) studied

Acupressure combined with flexible pressure sensor manometry to improve the quality of anesthesia recovery

Interventions

This study is a two-group parallel randomized trial. All the patients were divided into two groups (1:1): the acupressure group (group A) and the control group (group C) using the random number table method.

Group A acupressure method: The patient was asked to inhale, with the thumb acupoint pressed for 2.0s to reach the peak, and the pressure gradually decreased (0~3.5s), and each acupoint was pressed for 3 min. All three acupoints were pressed using the same method. The total intervention time was 30 min. The pressure of acupuncture points (N) was observed and the time of acupoint pressure was recorded (s).

Group C was not given acupressure after surgery, and the rest of the post-anesthesia care unit (PACU) care was the same as group A's.

Follow-up for both study arms:

During the recovery period of anesthesia:

PACU dwell time; Mean cough during the recovery period from anesthesia (times/min); Mean pulse during anesthesia recovery (times/min); Mild Cough; Severe cough; HR (>100 times/min); BP (>140mmHg); Stress Urinary Incontinence (Female)

After 6 hours:

Cough with bloody sputum; the throat is swollen and sore

Intervention Type

Procedure/Surgery

Primary outcome(s)

Cough Grading: The intensity and duration of coughing were recorded. According to the outbreak phase of the cough, a series of coughs in each time course is recorded as 1 cough, and the intensity of the cough is divided into three levels:
mild (1~2 times/min),
moderate (3~4 times/min),
and severe (>5 times/min).

The evaluation was done immediately after extubation, 30 minutes after removal of the laryngeal mask, and on the transfer out of the PACU.

Key secondary outcome(s)

The following secondary outcome measures data were collected throughout the procedure:

1. Duration of anesthesia
2. Remifentanil dosage
3. Micuronium chloride dose
4. Betamethasone dose
5. Duration of surgery
6. Endotracheal lavage
7. Endotracheal lavage recovery fluid

Completion date

30/05/2024

Eligibility

Key inclusion criteria

1. Age between 18-64 years old
2. ASA Class I or II
3. No cough or mild cough

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

64 years

Sex

All

Total final enrolment

132

Key exclusion criteria

1. Skin lesions around Tiantu (RN22), Chize (LU5), and Taiyuan (LU9) points
2. Recent massive hemoptysis
3. Diffuse alveolar hemorrhage
4. Untreated active tuberculosis
5. Obvious electrolyte imbalance
6. Inability to understand communication-related trials or informed consent

Date of first enrolment

26/12/2023

Date of final enrolment

30/05/2024

Locations

Countries of recruitment

China

Study participating centre

Department of Anesthesiology and Perioperative Medicine

First Affiliated Hospital of Zhengzhou University

Zhengzhou

China

450002

Sponsor information

Organisation

First Affiliated Hospital of Zhengzhou University

ROR

<https://ror.org/056swr059>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

First Affiliated Hospital of Zhengzhou University

Alternative Name(s)

The First Affiliated Hospital of Zhengzhou University,

Funding Body Type

Government organisation

Funding Body Subtype

Other non-profit organizations

Location

China

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Lijun Jia, fccjialj@zzu.edu.cn, after publication.

IPD sharing plan summary

Available on request, Published as a supplement to the results publication, Stored in publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		28/10/2025	29/10/2025	Yes	No