

Does rituximab added to usual treatment reduce kidney transplant failure following rejection?

Submission date 12/11/2018	Recruitment status Stopped	☐ Prospectively registered
		☐ Protocol
Registration date 11/03/2019	Overall study status Stopped	☐ Statistical analysis plan
		☐ Results
Last Edited 12/09/2023	Condition category Injury, Occupational Diseases, Poisoning	☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol

Background and study aims

Antibody-mediated rejection (AMR) is when the immune system of the person who has received a transplanted organ, known as the recipient, produces antibodies against the organ. This can lead to damage to the organ so that it does not function well. AMR is the leading cause of kidney transplant failure. How to best treat AMR is not known. Different treatment combinations are used across the world. This study aims to investigate whether a drug called rituximab, when added to the usual treatment for AMR, can improve transplant survival. Rituximab works by reducing the numbers of B cells, the white blood cells that produce antibodies. This drug is widely used in transplantation and in other diseases involving B cells.

Who can participate?

Patients aged 5 years or older with AMR of their kidney transplant

What does the study involve?

All patients will receive treatment accepted as standard of care for AMR. This involves the use of plasma exchange (a process that removes antibodies from a patient's body), together with steroids (to suppress immune system activity) and immunoglobulins (which inactivate antibodies). This treatment is given over approximately 2-3 weeks. One half of the patients will be allocated at random to receive rituximab. Rituximab is given through a patient's vein, and is administered in 2 separate doses, 2 weeks apart. After treatment, all patients will be followed up for a period of 4 years and have their transplant function monitored. Study visits will coincide with the usual hospital appointments.

What are the possible benefits and risks of participating?

There are no guaranteed benefits of taking part in the study. Use of immunosuppressants (drugs that reduce the activity of the immune system) leads to an increased risk of infection and cancer compared with the general population. All patients, regardless of whether they participate in the trial or not, will require immunosuppression to treat AMR. The side effects of rituximab include that in patients who have previously had hepatitis B, the infection might be reactivated. Patients with active hepatitis will be excluded and patients who have previously had hepatitis B

will be included or excluded from the trial on the decision of the researchers. There is also a very small chance of the brain condition progressive multifocal leukoencephalopathy (PML) with rituximab, but research suggests that this occurs in less than one in 25,000 people taking rituximab.

Where is the study run from?

The study is planned to take place across approximately 24 different hospitals, with Imperial College Healthcare NHS Trust being the lead centre.

When is the study starting and how long is it expected to run for?

November 2018 to February 2026

Who is funding the study?

The study is funded jointly by Kidney Research UK and the National Institute for Health Research.

Who is the main contact?

The chief investigator is Dr Michelle Willicombe

The trial contact is Dr Graham Armstrong, graham.armstrong@addenbrookes.nhs.uk

Contact information

Type(s)

Public

Contact name

Dr Graham Armstrong

Contact details

Cambridge Clinical Trials Unit,
Coton House, Level 6, Box 401,
Cambridge University Hospitals NHS Foundation Trust,
Addenbrooke's Hospital,
Hill's Road,
Cambridge
Cambridge
United Kingdom
CB2 0QQ
+44 01223 254918
graham.armstrong@addenbrookes.nhs.uk

Additional identifiers

ClinicalTrials.gov (NCT)

NCT03994783

Clinical Trials Information System (CTIS)

2018-002882-20

Central Portfolio Management System (CPMS)

40785

Protocol serial number

1.0

Study information

Scientific Title

Transplant Antibody Mediated Rejection: Guiding Effective Treatments (TAR:GET-1): A multicentre randomised controlled trial to assess the safety and efficacy of rituximab compared with control in treating acute antibody-mediated rejection in kidney transplantation

Acronym

TAR:GET-1

Study objectives

The study aims to test the hypothesis of whether rituximab in addition to standard of care treatment (plasma exchange, IVIG and corticosteroids) is more effective than standard of care alone in treating antibody-mediated rejection of kidney transplants.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 17/04/2019, London – West London & GTAC REC (The Old Chapel, Royal Standard Place, Nottingham, NG1 6FS, United Kingdom; +44 (0)2071048007; NRESCommittee.London-WestLondon@nhs.net), ref: 19/LO/0180

Primary study design

Interventional

Study design

Multicentre phase 3 open-label randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Antibody-mediated kidney transplant rejection

Interventions

Patients will be stratified by age and kidney transplant function and randomised at a 1:1 ratio to receive standard of care treatment alone or standard of care plus rituximab. All patients will receive standard of care treatment, which involves 7 plasma exchanges, intravenous immunoglobulins and corticosteroids (3 doses intravenous methylprednisolone followed by oral corticosteroids). This treatment will occur over a 2- to 3-week period. Those in the rituximab group will also receive rituximab dosed at 2 x 375 mg/m², given 2 weeks apart. Each patient will be followed up for a period of 48 months following randomisation.

Intervention Type

Drug

Phase

Phase III

Drug/device/biological/vaccine name(s)

Rituximab

Primary outcome(s)

Allograft survival, defined as the duration from the date of randomisation to the date of starting dialysis dependency or date of eGFR <15 mL/min/1.73 m², with follow-up of 48 months.

Key secondary outcome(s)

1. Serum creatinine) at 1, 3, 6, and 12 months post-randomisation and then annually until 4 years
2. Estimated GFR (CKD-EPI)) at 1, 3, 6, and 12 months post-randomisation and then annually until 4 years
3. Proteinuria (urinary protein:creatinine ratio)) at 1, 3, 6, and 12 months post-randomisation and then annually until 4 years
4. Donor-specific antibody (DSA) positivity at 3 and 12 months
5. Number of different DSA at at 3 and 12 months
6. Level of the immunodominant DSA assessed using mean fluorescence index at 3 and 12 months
7. Adverse event rate assessed by reviewing patient medical records
8. Patient-reported health-related quality of life , assessed using EQ-5D-5L/EQ-5D-Y questionnaires at baseline, 3 months, and 1, 2, 3 and 4 years following transplantation
9. Economic analysis of cost per quality-adjusted life year (QALY) gained from the perspective of the NHS

Completion date

01/02/2026

Reason abandoned (if study stopped)

Objectives no longer viable

Eligibility

Key inclusion criteria

1. Informed consent provided by patient or by a parent or legal guardian for patients aged <16 years
2. Aged 5 years or older
3. Diagnosis of acute antibody-mediated rejection (AMR) as defined by:
 - 3.1. The presence of ≥1 donor-specific antibodies (DSA)
 - 3.2. An adequate transplant biopsy (≥7 glomeruli and ≥1 artery) with histological features consistent with active AMR with no evidence of chronicity as defined by the Banff histological classification of allograft pathology:
 - 3.2.1. If C4d positive (2 or 3): v score (for arteritis) ≥1 and/or thrombotic microangiopathy and/or g score (for glomerulitis) ≥1 and/or ptc score (for peritubular capillaritis) ≥1, or if co-existing cellular rejection, a g score ≥1
 - 3.2.2. If C4d negative (0 or 1): microcirculation inflammatory score (g + ptc) ≥2, or if co-existing cellular rejection, a g score ≥1 and (g + ptc) ≥2 plus chronic glomerulopathy (cg) score 0 or 1a, or tubulo-interstitial fibrosis <50% and glomerular obsolescence <50%

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Sex

All

Key exclusion criteria

1. ABO-incompatible transplant
2. Received rituximab as part of induction or post-transplant for any other indications within the preceding 12 months (eg. recurrent focal and segmental glomerular sclerosis)
3. Received complete plasma exchange (PEX) treatment prior to the index biopsy on the suspicion of acute AMR in the absence of histology
4. Active infection including bacterial, viral (including CMV and EBV) or fungal infection or tuberculosis, which in the investigator's opinion could affect the conduct of the study
5. Co-existing BK nephropathy
6. Active hepatitis B or hepatitis C (patients with prior exposure to hepatitis B may be enrolled at the discretion of the PI; patients may be included if a negative hepatitis C recombinant immunoblot assay is confirmed or have a negative hepatitis C virus RNA [qualitative] test), or subjects with suspected human immunodeficiency virus (HIV) infection
7. Active malignancy
8. Known allergy, intolerance or contraindication to the treatments in the standard of care arm or rituximab as outlined in the Summaries of Product Characteristics (SmPCs)
9. Clinically significant comorbidity
10. Females must be either post-menopausal for at least 1 year, surgically sterile or, if of child-bearing potential, must not be pregnant or lactating. If sexually active, must agree to use an acceptable method of birth control for the first year post-randomisation.

Date of first enrolment

01/02/2019

Date of final enrolment

01/08/2022

Locations

Countries of recruitment

United Kingdom

England

Northern Ireland

Scotland

Wales

Study participating centre

Imperial College Healthcare NHS Trust

Hammersmith Hospital, Du Cane Road, Acton. W12 0HS
London
United Kingdom
W12 0HS

Study participating centre

Cambridge University Hospitals NHS Foundation Trust

Addenbrookes Hospital, Hills Road,
Cambridge
United Kingdom
CB2 0QQ

Study participating centre

OXFORD UNIVERSITY HOSPITALS NHS FOUNDATION TRUST

JOHN RADCLIFFE HOSPITAL
HEADLEY WAY
HEADINGTON OXFORD OXFORDSHIRE
Oxford
United Kingdom
OX3 9DU

Study participating centre

Leeds Teaching Hospitals NHS Trust

ST. JAMES'S UNIVERSITY HOSPITAL
BECKETT STREET
LEEDS WEST YORKSHIRE
Leeds
United Kingdom
LS9 7TF

Study participating centre

Great Ormond Street Hospital for Children NHS Foundation Trust

Great Ormond Street,
London
United Kingdom
WC1N 3JH

Study participating centre
GUY'S AND ST THOMAS' NHS FOUNDATION TRUST
GUY'S HOSPITAL
GREAT MAZE POND LONDON GREATER LONDON
London
United Kingdom
SE1 9RT

Study participating centre
Royal Infirmary of Edinburgh
51 Little France Crescent,
Edinburgh
Edinburgh
United Kingdom
EH16 4SA

Study participating centre
University Hospital of Wales
Cardiff & Vale University Health Board
Heath Park Way
Cardiff
United Kingdom
CF14 4XW

Study participating centre
NORTH BRISTOL NHS TRUST
SOUTHMEAD HOSPITAL, SOUTHMEAD ROAD
Bristol
United Kingdom
BS10 5NB

Study participating centre
UNIVERSITY HOSPITALS COVENTRY AND WARWICKSHIRE NHS TRUST
CLIFFORD BRIDGE ROAD
Coventry
United Kingdom
CV2 2DX

Study participating centre
NHS Greater Glasgow and Clyde
Queen Elizabeth University Hospital 1345 Govan Road

Glasgow
United Kingdom
G51 4TF

Study participating centre

UNIVERSITY HOSPITALS OF LEICESTER NHS TRUST
LEICESTER ROYAL INFIRMARY, INFIRMARY SQUARE
Leicester
United Kingdom
LE1 5WW

Study participating centre

ROYAL LIVERPOOL AND BROADGREEN UNIVERSITY HOSPITALS NHS TRUST
ROYAL LIVERPOOL UNIVERSITY HOSPITAL, PRESCOT STREET
Liverpool
United Kingdom
L7 8XP

Study participating centre

ST GEORGE'S UNIVERSITY HOSPITALS NHS FOUNDATION TRUST
St George's Hospital, Blackshaw Road,
London
United Kingdom
SW17 0QT

Study participating centre

ROYAL FREE LONDON NHS FOUNDATION TRUST
ROYAL FREE HOSPITAL, POND STREET
London
United Kingdom
NW3 2QG

Study participating centre

BARTS HEALTH NHS TRUST
THE ROYAL LONDON HOSPITAL, WHITECHAPEL
London
United Kingdom
E1 1BB

Study participating centre

MANCHESTER UNIVERSITY NHS FOUNDATION TRUST

Manchester Royal Infirmary, Oxford Road
Manchester
United Kingdom
M13 9WL

Study participating centre

The Newcastle Upon Tyne Hospitals NHS Foundation Trust

Freeman Hospital
Freeman Road
Newcastle
United Kingdom
NE7 7DN

Study participating centre

NOTTINGHAM UNIVERSITY HOSPITALS NHS TRUST

City Hospital
Hucknall Road
Nottingham
United Kingdom
NG5 1PB

Study participating centre

PLYMOUTH HOSPITALS NHS TRUST

DERRIFORD HOSPITAL, DERRIFORD ROAD
Plymouth
United Kingdom
PL6 8DH

Study participating centre

Portsmouth Hospitals NHS Trust

Queen Alexandra Hospital
Portsmouth
United Kingdom
PO6 3LY

Study participating centre

SHEFFIELD TEACHING HOSPITALS NHS FOUNDATION TRUST

NORTHERN GENERAL HOSPITAL, HERRIES ROAD
Sheffield

United Kingdom
S5 7AU

Study participating centre
Belfast City Hospital
Belfast
N Ireland
Belfast
United Kingdom
BT9 7AB

Sponsor information

Organisation
Imperial College London

ROR
<https://ror.org/041kmwe10>

Funder(s)

Funder type
Charity

Funder Name
Kidney Research UK

Alternative Name(s)

Funding Body Type
Private sector organisation

Funding Body Subtype
Trusts, charities, foundations (both public and private)

Location
United Kingdom

Funder Name
National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date