

# Understanding the impact of different doses of Reducose® mulberry leaf extract on blood glucose and insulin responses after eating a complex meal

<b>Submission date</b> 30/05/2022	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 08/06/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 19/07/2024	<b>Condition category</b> Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Reducose mulberry leaf extract can be used to lower the body's glucose (sugar) response after eating foods containing carbohydrates. It has been developed for use as both a food ingredient and for use in dietary supplements. The sponsor has successfully completed four clinical trials with Reducose which evaluated the effects of Reducose® in lowering the glycaemic index of common carbohydrates in a healthy Indian population; the effects of Reducose® in lowering the glycaemic index of common carbohydrates in a healthy Chinese population; the effects of three different doses of Reducose® versus placebo on the blood glucose and insulin response in healthy individuals following a starch challenge; and the effects of Reducose® versus placebo on the blood glucose and insulin response in healthy individuals following a sugar challenge. This study aims to evaluate the effects of three different doses of a dietary supplement (capsule) containing Reducose® on the blood glucose and insulin response of healthy volunteers after a complex meal. As proteins and fats are known to impact gastric (stomach) emptying and blood glucose response, a meal containing fats, proteins and carbohydrates will be used. Three dose levels are included to better understand the dose-response of Reducose®.

### Who can participate?

Healthy volunteers aged 18-60 years with a body mass index (BMI) of 20 - 29.9 kg/m<sup>2</sup>

### What does the study involve?

Participants will be recruited to attend four study visits, at least 48 hours apart, in order to test either three different doses of Reducose dietary supplement capsules or a matching placebo (dummy supplement). The test product or placebo will be taken before eating a meal that matches the composition of a typical Western diet. This process will be replicated in each of the four study visits. The test products are packaged in opaque capsules that have been matched for size, weight, and colour, and neither participants nor the study team will be aware of the order

that the test products are given until all of the tests have been completed. Blood will be drawn from finger pricks at regular intervals over 180 minutes. Changes in blood glucose and insulin levels will be measured in the blood samples.

What are the possible benefits and risks of participating?

Mulberry extract has an excellent safety profile and is widely consumed in Asia and is available as a dietary supplement around the world. Previous clinical trials have shown that Reducose does not have any reported adverse reactions. Participants will have multiple finger pricks to test for blood glucose and insulin levels, therefore there will be a risk of discomfort and bruising and a small risk of inflammation/infection at the site. Local infection control/procedures will be followed during each study visit.

Where is the study run from?

Oxford Brookes Centre for Nutrition and Health, Oxford Brookes University (UK)

When is the study starting and how long is it expected to run for?

August 2019 to May 2021

Who is funding the study?

Phynova Group Ltd (UK)

Who is the main contact?

Andrew Gallagher

agallagher@phynova.com

## Contact information

### Type(s)

Scientific

### Contact name

Mr Andrew Gallagher

### ORCID ID

<https://orcid.org/0000-0002-5974-4093>

### Contact details

Office 3 at Magenta

2 Brookhill Way

Banbury

United Kingdom

OX16 3ED

+44 (0)1993880700

agallagher@phynova.com

## Additional identifiers

### Protocol serial number

REDUCE 1.1

# Study information

## Scientific Title

A double-blind, placebo-controlled, four-arm, single-dose, cross over, dose-ranging trial to investigate the effects of Reducose mulberry leaf extract on postprandial glucose and insulin after consuming a complex meal

## Acronym

REDUCE

## Study objectives

The primary hypothesis is that a dietary supplement tablet containing mulberry leaf extract (Reducose®) when co-administered with mixed macronutrient meal will reduce the incremental area under the curve for plasma glucose concentration over 120 minutes in normoglycaemic adults when compared to co-administration with a placebo in a dose-dependent manner. The secondary hypothesis is that the dietary supplement when co-administered with a meal will not disproportionately increase the incremental area under the curve for plasma insulin concentration over 120 minutes in normoglycaemic adults compared to co-administration with placebo.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

1. Approved 07/03/2014, University Research Ethics Committee (UREC) at Oxford Brookes University (Research & Business Development Office, Directorate of Finance & Legal Services, Oxford Brookes University, Headington Campus, Gypsy Lane, Oxford, OX3 0BP, UK; +44 (0) 1865 48 4445; ethics@brookes.ac.uk), ref: 140806 (for glycaemic response),
2. Approved 24/01/2012, University Research Ethics Committee (UREC) at Oxford Brookes University (Research & Business Development Office, Directorate of Finance & Legal Services, Oxford Brookes University, Headington Campus, Gypsy Lane, Oxford, OX3 0BP, UK; +44 (0) 1865 48 4445; ethics@brookes.ac.uk), ref: 110594 (for insulinaemic response)

## Study design

Double-blind placebo-controlled randomized repeat measure cross over dose-ranging trial

## Primary study design

Interventional

## Study type(s)

Prevention

## Health condition(s) or problem(s) studied

Postprandial glycaemic control

## Interventions

The trial has a randomised cross over design and so each subject would act as their own control. Subjects are randomized to the trial using a 4x4 Latin Square block randomization, randomizing participants to one of three active doses or placebo in a block-wise manner. Researchers recruiting the participants are unaware of the allocation sequence (concealed allocation).

Participants are assigned to a participant number according to their chronological order of enrolment in the study. The allocated participant number is used to identify the participants and their corresponding intervention sequence.

Participants are recruited to attend four study visits, with at least 48-hour washout period between visits. The study will evaluate three different doses of Reducose mulberry leaf extract (200 mg, 225 mg or 250 mg) against matched placebo. Reducose mulberry leaf extract is standardized to contain 5% 1-deoxynojorimycin (w/w). The test product is individually encapsulated in identical, opaque capsules (CapsuGel V-Caps Plus, Swedish Orange colour), and individually blistered and packed into cardboard boxes. The placebo contains microcrystalline cellulose and is matched for size, weight and colour.

At T-10 minutes subjects will take either the test product or placebo product with a little water.

At T=0 minutes subjects should consume the entire test meal within 10 minutes. The test meal consisted of 150 g Hovis White Bread Thick Slice and 50g Tesco Egg Mayo filler providing a macronutrient composition of 60.9% carbohydrate, 16.6% protein and 20.4% fat.

Blood samples are taken at -5 min, 0 min (baseline taken as mean of -5 min and 0 min), 15, 30, 45, 60, 90, 120, 150 and 180 min. Glycaemic response is measured using the method described by Brous et al. and carried out in accordance with ISO 26642 (2010). Blood drawn at the above time points is used to measure blood glucose and blood insulin. 300µL of capillary blood is required for insulin analysis. Sufficient blood should be drawn to enable two sets of analysis at every time point to reduce the likelihood of missing data.

## **Intervention Type**

Supplement

## **Primary outcome(s)**

1. Blood glucose concentrations measured from capillary blood using HemoGue Glucose 201 DM analyser (units = mmol/L), measured at -5 min, 0 min, 15 min, 30 min, 45 min, 60 min, 90 min, 120 min, 150 min, 180 min
2. Blood glucose incremental area under curve (iAUC) calculated from blood glucose concentrations using the trapezoid rule (units = mmol/L/min), measured at 60, 90, 120, 150 and 180 min
3. Peak blood glucose calculated from blood glucose curves (see above) (units = mmol/L)
4. Time of peak blood glucose calculated from blood glucose curves (see above) (units = minutes)

## **Key secondary outcome(s)**

1. Plasma insulin concentration measured from capillary blood using electrochemiluminescence immunoassay using automated analyzer (Cobas E411) (units = µU/mL) at -5 min, 0 min, 15 min, 30 min, 45 min, 60 min, 90 min, 120 min, 150 min, 180 min
2. Plasma insulin iAUC calculated from plasma insulin concentrations using the trapezoid rule (units = µU/mL/min), measured at 60, 90, 120, 150, and 180 min

## **Completion date**

17/05/2021

## **Eligibility**

### **Key inclusion criteria**

1. Consent to study protocol
2. Aged 18-60 years
3. Body mass index (BMI) 20-29.9 kg/m<sup>2</sup>
4. Fasting blood glucose <6.1 mmol/l

**Participant type(s)**

Healthy volunteer

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Upper age limit**

60 years

**Sex**

All

**Total final enrolment**

43

**Key exclusion criteria**

1. Aged < 18 or > 60 years
2. Pregnant or lactating
3. Body mass index (BMI) <18.5 or >30 kg/m<sup>2</sup>
4. Fasting blood glucose value >6.1 mmol/l
5. Any known food allergy or intolerance
6. Medical condition(s) or medication(s) known to affect glucose regulation or appetite and/or which influence digestion and absorption of nutrients (stable doses of oral contraceptives, acetylsalicylic acid, thyroxin, vitamins and mineral supplements or drugs to treat hypertension or osteoporosis are acceptable)
7. Known history of diabetes mellitus or the use of antihyperglycaemic drugs or insulin to treat diabetes and related conditions
8. Known IBS, severe liver, heart or kidney problems or blood clotting disorders
9. Major medical or surgical event requiring hospitalization within the preceding 3 months
10. Use of steroids, protease inhibitors or antipsychotics (all of which have major effects on glucose metabolism and body fat distribution)
11. High vulnerability level for COVID-19 (assessed using this tool - <https://alama.org.uk/covid-19-medical-risk-assessment/>)
12. Displayed any COVID-19 symptoms 14 days prior to any test session (high temperature, new continuous cough, loss or change to sense of smell or taste)

**Date of first enrolment**

03/01/2020

**Date of final enrolment**

07/05/2021

## Locations

### Countries of recruitment

United Kingdom

England

### Study participating centre

#### Oxford Brookes Centre for Nutrition and Health

Oxford Brookes University

Headington Campus

Oxford

United Kingdom

OX3 0BP

## Sponsor information

### Organisation

Phynova Group Ltd

## Funder(s)

### Funder type

Industry

### Funder Name

Phynova Group Ltd

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated and analysed during the current study will be available to researchers upon request from Phynova Group Limited. Anonymised patient level data will be available for research purposes within six months of publishing the results of the primary endpoints, key secondary endpoints and safety data in a peer-reviewed journal. As participants only consented for their data to be used to investigate a particular health benefit of the test product, access to data will only be provided for researchers that are investigating the same or an equivalent product for the same health benefit. Data will be made available on request provided Phynova

has the legal authority to provide the data; for example, if Phynova has out-licensed the data to another company then it would no longer have the legal authority to provide the data. Data will only be made available if Phynova is able to anonymise the data without compromising the confidentiality and privacy of research participants. Data from clinical trials where the data labels and/or supporting documents are not in English will only be made available in their original format and no translations will be provided. The following information will be made available: raw dataset excluding any patient images, protocol with any amendments, analysis and reporting plan. Researchers requesting access to data should email their requests to [info@phynova.com](mailto:info@phynova.com) and should include a description of the intended research.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		29/05/2024	19/07/2024	Yes	No