

How does hyalobarrier gel influence ovulation and pregnancy outcomes in women with scarring around the tubes and ovaries?

Submission date 27/07/2016	Recruitment status No longer recruiting	☐ Prospectively registered
		☐ Protocol
Registration date 14/09/2016	Overall study status Completed	☐ Statistical analysis plan
		☒ Results
Last Edited 05/12/2016	Condition category Pregnancy and Childbirth	☐ Individual participant data

Plain English summary of protocol

Background and study aims

An adhesion is a type of scar tissue that can make the tissues or organs inside the body stick together. This type of scarring is often caused by undergoing a laparoscopy (a type of operation in which the surgeon accesses the inside of the abdomen (tummy) or pelvis through a small cut. If the adhesions are around the fallopian tubes and ovaries, it can reduce fertility, as it can interfere with the monthly release of an egg necessary for conception. The removal of scarring using an anti-scarring gel is potentially useful for stopping scars from reforming after surgery. But it is not known if the gel will influence ovulation (egg production) monthly. The aim of this study is to find out whether using an anti-scarring gel called Hyalobarrier after a laparoscopy has an effect on the function of ovaries.

Who can participate?

Women aged between 18 and 28 who are undergoing a laparoscopy for the treatment of a condition relating to the female reproductive system, who have possible

What does the study involve?

Participants are randomly allocated to one of two groups. Both groups undergo a laparoscopy according to standard procedure, but those in the first group have 10ml hyalobarrier applied around the operative site. Participants have levels of reproductive hormones measured at the start of the study and then again after three and six months, as well as undergoing ultrasound scanning at the same times to monitor their egg production.

What are the possible benefits and risks of participating?

There are no direct benefits or risks involved for participants taking part in this study.

Where is the study run from?

Princess Anne Hospital, Southampton (UK)

When is the study starting and how long is it expected to run for?

November 2011 to November 2014

Who is funding the study?
Nordic Pharma Ltd. (UK)

Who is the main contact?
Professor Ying Cheong
y.cheong@soton.ac.uk

Contact information

Type(s)
Scientific

Contact name
Prof Ying Cheong

ORCID ID
<https://orcid.org/0000-0001-7687-4597>

Contact details
Human Development and Health
University of Southampton Faculty of Medicine
Level F
Princess Anne Hospital
Coxford Road
Southampton
United Kingdom
SO16 5YA
+44 (0)2380 216033
y.cheong@soton.ac.uk

Additional identifiers

Protocol serial number
11/H0504/6

Study information

Scientific Title
Randomised controlled trial of hyalobarrier versus no hyalobarrier on the ovulatory status of women with peri-ovarian adhesions: a pilot study

Study objectives
The aim of this study is to compare the ovarian function of women who have peri-ovarian adhesiolysis with and without Hyalobarrier as an anti-adhesion barrier.

Ethics approval required
Old ethics approval format

Ethics approval(s)

Primary study design

Interventional

Study design

Randomised controlled pilot study

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Adnexal adhesions

Interventions

Participants are randomised to one of two groups using a computer generated random numbers table and opaque envelopes.

Intervention group: Participants receive hyalobarrier gel at laparoscopy. This involves having 10ml hyalobarrier applied around the operative site as per manufacturer's instruction.

Control group: Participants receive no hyalobarrier gel at laparoscopy.

Follow up for all participants involves hormonal profiling (Day 2/3 FSH and LH, and Day 21 progesterone) plus follicular tracking using ultrasonography at month 3 and month 6 post operatively.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Ovulatory status is measured using serum hormonal status (Day 2 FSH, LH and Day 21 progesterone) prior to the surgery and at 3 and 6 months after the surgery. Post-operatively, the participant will also undergo a follicular tracking cycle at 3 and 6 months.

Key secondary outcome(s)

Clinical pregnancy is measured by the presence of a fetal heart at 5-6 weeks gestation.

Completion date

01/11/2014

Eligibility**Key inclusion criteria**

1. Aged between 18-38
2. Women were undergoing operative laparoscopy for gynaecological pathology, with possible peri-ovarian adhesions

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Upper age limit

38 Years

Sex

Female

Key exclusion criteria

1. Presence of malignancies or a history of malignancies
2. Women on medications that affected ovulation
3. Women with known conditions that resulted in anovulation (PCOS, Pituitary causes)

Date of first enrolment

01/02/2012

Date of final enrolment

01/02/2014

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

Princess Anne Hospital

Coxford Road

Southampton

United Kingdom

So16 5YA

Sponsor information**Organisation**

Princess Anne Hospital Southampton

ROR

<https://ror.org/02yjksy18>

Funder(s)

Funder type

Industry

Funder Name

Nordic Pharma Ltd.

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/01/2017		Yes	No