

Study of a milk protein concentrate supplement to help recovery after hip or knee replacement surgery

Submission date 22/03/2026	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 23/03/2026	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 31/03/2026	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

This study looked at whether a milk protein supplement could help people recover better after hip or knee replacement surgery. The supplement, called milk protein concentrate, is designed to support muscle strength and reduce inflammation. Researchers wanted to find out if adding this supplement to a person's usual diet could improve their mobility, muscle mass, and general recovery after surgery.

Who can participate?

Adults aged 45 to 85 years who were scheduled for a planned hip or knee replacement at the 424 General Military Training Hospital in Greece could take part. They had to be able to give informed consent and follow the study instructions. People could not join if they had allergies to milk or soy proteins, severe kidney problems, certain autoimmune or neurological conditions, active infections, dementia, or if they were having revision surgery.

What does the study involve?

Participants were placed into one of two groups. Both groups received the same amount of daily protein, but one group received part of their protein intake in the form of the milk protein supplement. This was taken twice a day by mixing the powder with drinks such as milk, kefir, or juice. All participants followed a standard physiotherapy programme after surgery. Researchers measured strength, walking speed, balance, muscle mass, and markers of inflammation before surgery and again 15 weeks later.

What are the possible benefits and risks of participating?

Taking part might help improve recovery after surgery by supporting muscle strength and reducing inflammation. However, this could not be guaranteed. Risks included possible digestive discomfort or allergic reactions, especially for people sensitive to milk or soy proteins. The study excluded anyone with known allergies to reduce this risk.

Where is the study run from?

The study was carried out at the 424 General Military Training Hospital in Thessaloniki, Greece.

When is the study starting and how long is it expected to run for?
Participants were first enrolled on 9 May 2023, and the final participant was enrolled on 18 November 2024. The study was completed on 20 April 2025.

Who is funding the study?
The study is funded by the International Hellenic University, Greece.

Who is the main contact?
Mrs Maria Spanoudaki, maryspan1@gmail.com
Professor Sousana K. Papadopoulou, souzpapa@gmail.com

Contact information

Type(s)
Public

Contact name
Mrs Maria Spanoudaki

Contact details
International Hellenic University/Department of Nutritional Sciences and Dietetics
Sindos
Greece
57001
+30 6936 853 437
maryspan1@gmail.com

Type(s)
Scientific, Principal investigator

Contact name
Prof Sousana K. Papadopoulou

Contact details
International Hellenic University/Department of Nutritional Sciences and Dietetics
Sindos
Greece
57001
+30 694 479 8916
souzpapa@gmail.com

Additional identifiers

Study information

Scientific Title
The effect of Milk Protein Concentrate (MPC) supplement on functional recovery, muscle strength, inflammatory and oxidative markers in elderly patients following total hip or knee arthroplasty: A randomized controlled trial

Study objectives

To evaluate the role of a combined Milk Protein Concentrate (MPC) with essential minerals (Calcium and Phosphorus),(Protifar, Nutricia), in supporting muscle and overall recovery in orthopaedic patients.

Ethics approval required

Ethics approval required

Ethics approval(s)

Submitted 31/07/2023, Scientific Council of the 424 General Military Training Hospital (424 GSNE). (Ring Road, Nea Efkarpia, Thessaloniki, 56429, Greece; +2310381000; 424-gsne@army.gr), ref: 6th Topic, Scientific Council Session July 2023/ 424 Military Hospital of Thessaloniki

Primary study design

Interventional

Allocation

Randomized controlled trial

Masking

Blinded (masking used)

Control

Active

Assignment

Parallel

Purpose

Supportive care

Study type(s)

Health condition(s) or problem(s) studied

Primary knee or hip arthroplasty due to end-stage osteoarthritis of the lower limbs

Interventions

The intervention consists of oral nutritional supplementation with Protifar (Nutricia), a high-protein Milk Protein Concentrate (MPC) powder. The formula features a low sodium and potassium profile and a specific Calcium-to-Phosphorus (Ca/P) ratio of 1.9:1, optimized for bone and muscle recovery. The omega-6 to omega-3 (n-6/n-3) ratio is 3.3:1, with trace amounts of total fat to avoid interference with systemic inflammation markers. The solution exhibits low osmolality (25 mOsm/L), ensuring high gastrointestinal tolerance. The product is officially notified to the Greek National Organization for Medicines (EOF, Prot. No. 81732/13-11-2009). Dietary caloric intake is standardized for both groups. Total protein intake is set at 1.3 g/kg/day for all participants.

Participants were allocated to the intervention and control groups using a computer-generated random sequence (1:1 ratio) created via Microsoft Excel.

Intervention Group (IG): 30g of the daily protein requirement is provided via the MPC formula, administered in two equal doses of 15g (at breakfast and 30 minutes post-physiotherapy). Each dose is diluted in 50 ml of water and added to milk, kefir, or juice to ensure compliance.

Control Group (CG): Receives the same total protein intake (1.3 g/kg/day) solely through standard hospital meals, without the specific MPC supplementation."

Dietary caloric intake is standardized for both groups, with total protein intake set at 1.3 g/kg /day. For the Intervention Group (IG), 30g of this daily requirement is administered in the form of a milk protein concentrate (MPC) formula, divided into two equal doses: 15g with breakfast and 15g approximately 30 minutes after the physiotherapy session. Each dose is diluted in 50 ml of water and subsequently added to milk, kefir, or juice to ensure compliance and palatability. The Control Group receives the same total protein intake (1.3 g/kg/day) through standard meals without the specific MPC supplementation.

IG and CG followed a standardized early physical therapy program, which focused on isometric exercises and progressive resistance training. The CG and IG were recruited during different, non-overlapping time periods. This design ensured that the groups remained independent.

The total duration of the intervention, including the follow-up period, was 15 weeks for both study arms.

Intervention Type

Supplement

Primary outcome(s)

1. Functional performance measured using handgrip strength(kgr); endurance (4 meters gait speed (m/sec); static balance (Single leg stance test(sec); time for 5 repetitions sit to stand test (sec) at 24 hours before surgery and 15 weeks post-operatively
2. Skeletal Muscle Mass(kgr); Lean Mass (kgr) measured using Bioelectrical Impedance Analysis (kgr) at 24 hours before surgery and 15 weeks post-operatively

Key secondary outcome(s)

1. Inflammation measured using C Reactive Protein in serum(mg/dL) at 24 hours before surgery and 15 weeks post-operatively
2. Oxidative Stress measured using 8-Isoprostane in plasma and urine(pg/ml) at 24 hours before surgery and 15 weeks post-operatively
3. Clinical and biochemical data, including blood and urine analysis (CRP and 8-isoprostane levels), as recorded in the study's structured assessment forms. These measures are part of the protocol approved by the Scientific Council of the 424 General Military Training Hospital (Session July 2023, 6th Topic), evaluating the impact of the MPC nutritional intervention (notified to EOF, Prot. No. 81732/13-11-2009)." measured using 8-Isoprostane in plasma and urine(pg/ml) and CRP (mg/dl) at 24 hours before surgery and 15 weeks post-operatively

Completion date

20/04/2025

Eligibility

Key inclusion criteria

1. Adults aged 18 years and over
2. Scheduled for elective primary total hip or knee arthroplasty at 424 General Military Hospital
3. Ability to provide written informed consent and willingness to comply with the nutritional supplementation protocol and follow-up assessments

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

45 Years

Upper age limit

85 Years

Sex

All

Total final enrolment

88

Key exclusion criteria

Current key exclusion criteria 31/03/2026:

1. Known allergy to milk and soy proteins or severe lactose intolerance
2. Severe renal impairment or kidney disease
3. Autoimmune diseases, Cognitive impairment, dementia, or psychiatric disorders, neuro-muscular disorders
4. Rheumatoid arthritis or active infection
5. Revision surgery
6. Individuals with active implantable medical devices.

Previous key exclusion criteria:

1. Known allergy to milk and soy proteins or severe lactose intolerance
2. Severe renal impairment or kidney disease
3. Autoimmune diseases, Cognitive impairment, dementia, or psychiatric disorders, neuro-muscular disorders
4. Rheumatoid arthritis or active infection
5. Revision surgery

Date of first enrolment

09/05/2023

Date of final enrolment

18/11/2024

Locations

Countries of recruitment

Greece

Sponsor information

Organisation

International Hellenic University

ROR

<https://ror.org/00708jp83>

Funder(s)

Funder type

Funder Name

International Hellenic University

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Other files	Participant flow diagram		31/03/2026	No	No