

# Safety and performance of a novel device for the administration of regional anaesthesia

<b>Submission date</b> 22/12/2021	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 27/01/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 27/01/2022	<b>Condition category</b> Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Regional anaesthesia procedures require two operators - an anaesthetist who holds an ultrasound scanner and uses this to guide the needle tip placement and a second operator to inject the anaesthetic solution at a required pressure.

It has been reported that anaesthetic solutions are often injected at high pressure.

High pressure can cause damage to nerve fascicles, with serious nerve damage occurring in up to 1% of procedures and transient nerve damage in up to 8% of cases.

SAFIRA® - SAFer Injection for Regional Anaesthesia - allows a single operator, an anaesthetist or anesthesiologist, to conduct the whole regional block at safer pressures.

The study will examine the following questions:

- Does the SAFIRA device perform as intended in routine clinical practice?
- Is the SAFIRA device safe to use as per intended use?
- What is the user feedback regarding acceptability of and confidence in SAFIRA?

### Who can participate?

Peripheral nerve block (PNB) procedures involving patients aged > 18 years referred for any type of elective surgery suitable for PNB and who are offered and agree to surgery that will be performed via any ultrasound guidance and/ or nerve stimulation single-injection PNB regional anaesthesia. Any type of PNB will be included in the study.

### What does the study involve?

This study will involve 43 regional anaesthesia procedures undertaken by six anaesthetists already trained in the use of the SAFIRA system in one UK hospital site, in order to ascertain the safety and performance of SAFIRA in a 'real-world' clinical setting. The study will be entirely observational. The investigation will not involve any clinical investigation or treatment additional to standard care and will not include any patient-orientated research instruments. All patients will undergo treatment according to normal clinical practice.

### What are the possible benefits and risks of participating?

Unintentional needle-to-nerve contact, intraneural penetration or intrafascicular penetration during PNB may cause nerve injury which, even if minor and transient, presents a sizeable

problem when the number of PNB procedures are scaled-up to a healthcare system or population level. Patients experiencing more severe and/or persistent nerve injury may experience significant negative consequences in terms of decreased health status, physical function, ability to work and quality of life. Meta-analyses have identified that Anaesthetists cannot rely on ultra sound guidance, nerve stimulation, 'syringe-feel' or injection pressure monitoring as indicators of unintentional needle-to-nerve contact, intraneural penetration or intrafascicular penetration. It is possible that the safest way of preventing nerve damage might be to automatically limit local anaesthetic (LA) opening and injection pressure to levels that are not associated with nerve damage. The SAFIRA system is a medical device that limits LA opening and injection pressures to a level of >20psi. This study will assess the safety and performance of SAFIRA in a 'real-world' clinical setting. It is important to note that the SAFIRA system has all necessary approvals and is currently in use within the NHS.

Where is the study run from?

Queen Elizabeth Hospital NHS Foundation Trust King's Lynn (UK)

When is the study starting and how long is it expected to run for?

March 2021 to March 2022

Who is funding the study?

Medovate Ltd (UK)

Who is the main contact?

Dr Benjamin Fox, benjamin.fox@qehkl.nhs.uk

## Contact information

### Type(s)

Principal investigator

### Contact name

Dr Benjamin Fox

### Contact details

Queen Elizabeth Hospital King's Lynn NHS Foundation Trust

King's Lynn

United Kingdom

CB1 1PT

+44 1553613583

benjamin.fox@qehkl.nhs.uk

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### Integrated Research Application System (IRAS)

293036

### Protocol serial number

## Study information

### Scientific Title

Prospective, open, non-controlled, single-arm post-market clinical follow-up investigation to confirm the safety and performance of the SAFIRA system in ultrasound guidance and/or nerve stimulation peripheral nerve block (PNB)

### Study objectives

The study will examine:

- Does the SAFIRA device perform as intended in routine clinical practice?
- Is the SAFIRA device safe to use as per intended use, are the known risks acceptable and do any new risks identified impact the benefit-risk ratio?
- What, if any, is the impact on the clinical workflow of PNB procedures when using SAFIRA compared with standard practice?
- What is the user feedback regarding acceptability of and confidence in SAFIRA for performing PNB procedures?

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Approved 22/03/2021, HRA and Health and Care Research Wales (HCRW) (HRA, NHSBT Newcastle Blood Donor Centre, Holland Dr, Newcastle upon Tyne NE2 4NQ, UK; +44 207 972545; HCRW.approvals@wales.nhs.uk), ref: 21/ES/0029

### Study design

Prospective open non-controlled single-arm post-market clinical follow-up investigation

### Primary study design

Observational

### Study type(s)

Treatment

### Health condition(s) or problem(s) studied

Safety and performance of the SAFIRA system in ultrasound guidance and/or nerve stimulation peripheral nerve block

### Interventions

This prospective, open, non-controlled, single-arm post-market clinical follow-up investigation will involve 43 PNB procedures undertaken by up to 6 anaesthetists trained in the use of the SAFIRA system in one UK hospital site; in order to ascertain the safety and performance of SAFIRA in a 'real-world' clinical setting. For the purposes of the study any operation that requires more than one simultaneous PNB (e.g. for complex lower limb surgery) will be considered a single PNB procedure. The study will be entirely observational. The investigation will not involve allocation (or withholding) of any aspect of clinical care, will not involve any clinical investigation or treatment additional to standard care and will not include any patient-orientated research instruments (e.g. multidimensional pain questionnaires) unless routinely used at participating

sites. All patients will undergo treatment according to normal clinical practice i.e. as determined by local institutional protocol and/or individual clinician discretion. The principal units of analysis will be PNB procedures undertaken with the SAFIRA system and various procedure-related, patient-related and operator-related variables will be assessed.

In summary, at the participating site a study assessor (SA) - a trained member of the care team not directly involved in the care of patients within the study - will record the following data:

- i) Procedural data related to anaesthetists use of the SAFIRA system in patients undergoing any type of elective surgery via any type of PNB involving ultrasound guidance or nerve stimulation;
- ii) Routine clinical data (or clinical data potentially available for collection without impacting on patient care) in patients undergoing SAFIRA PNB procedures
- iii) Anaesthetists appraisal of the SAFIRA system.

The study will not consider outcome measures of any of the surgeries undertaken using PNB nor will it consider the underlying effectiveness of ultrasound guidance or nerve stimulation.

## **Intervention Type**

Device

## **Phase**

Not Applicable

## **Drug/device/biological/vaccine name(s)**

SAFIRA system

## **Primary outcome(s)**

1. Time to perform block at 'Day 0'. This will be defined as time from the first application of the probe on skin for ultrasound guidance or first application of needle on the skin for nerve stimulation to final removal of the PNB local anaesthetic needle.
2. Composite measure of block success at 'Day 0'. This will be defined as adequate block (sufficient blocking of the transmission of nerve impulses as assessed by routine institutional sensory and/or motor testing protocol to allow surgery to proceed) AND no block conversion (avoidance of rescue block) NB: both composite and separate measures of block success will be reported.
3. Safety at 'Day 0' measured using a composite measure of evidence of potential acute nerve injury. This will be defined as any patient-reported severe and painful symptoms such as tingling, prickling, burning, uncomfortable pins and needles or any type of electric shock sensation during the PNB procedure AND suspected or confirmed needle-nerve contact, intraneural needle penetration or intrafascicular needle penetration as reported by the anaesthetist NB: both composite and separate measures of potential nerve injury will be reported.
4. Safety at 'Day 30' measured using evidence of persistent nerve injury. This will be defined as any documented evidence in the patient medical records of sensory and/or motor deficit including: i) patient-reported chronic, severe and painful symptoms such as tingling, prickling, burning, uncomfortable pins and needles or any type of electric shock sensation; ii) patient-reported limb or muscle weakness or iii) clinician-observed sensory and/or motor deficit lasting longer than 48 hours after surgery. Sensory and/or motor deficit in a different regional area to that of the PNB will be recorded but will be considered separately to sensory and/or motor deficit in the regional area of the PNB.

## **Key secondary outcome(s)**

Measured at 'Day 0' using patient records and questionnaire completed by anaesthetists:

1. Number of needle punctures (any new needle insertion through skin)
2. Number of needle redirects (any needle insertion-withdrawal-insertion of  $\geq 10$ mm)

3. Block onset time (interval between completion of LA injection and adequate sensory block to permit surgery in distribution of blocked nerve)
4. Event-free SAFIRA assembly/deployment (any suspected or reported issue in any of the steps of the SAFIRA Instruction for Use)
5. SAFIRA malfunction/failure (any event requiring conversion to manual injection)
6. Major complications (cardiac arrest, pneumothorax, death, other (specify))
7. Minor complications (vascular puncture, haematoma, local anaesthetic toxicity, cardiac arrhythmia, systemic hypotension, other (specify))
8. LA volume required
9. Anaesthetists appraisal of the SAFIRA system after completing all each PNB procedure
10. Injection Pressure (How important is limiting LA opening/injection pressure in your own clinical practice: Not at all important/Quite unimportant/Neither unimportant or important /Quite important/Very important)
11. Ease of Use (Compared to usual practice SAFIRA is easy to use: Strongly disagree/Disagree /Neither disagree or agree/Agree/Strongly agree)
12. Effectiveness (Compared to usual practice SAFIRA is effective: Strongly disagree/Disagree /Neither disagree or agree/Agree/Strongly agree)
13. Drawbacks (Compared to usual practice SAFIRA has few drawbacks: Strongly disagree /Disagree/Neither disagree or agree/Agree/Strongly agree)
14. Workload (Compared to usual practice SAFIRA increases workload: Strongly disagree /Disagree/Neither disagree or agree/Agree/Strongly agree)
15. Preference (Compared to usual practice I prefer SAFIRA: Strongly disagree/Disagree/Neither disagree or agree/Agree/Strongly agree)

Measured at 'Day 30' using patient records and questionnaire completed by anaesthetists::

16. Infection at PNB site
17. Falls (any documented inpatient or at home falls in 30 days following surgery)
18. PNB limb trauma (any documented inpatient or at home trauma (scalds, burns, severe pressure sore) in 30 days following surgery)
19. Postoperative 1hr pain score (assessed using standard institutional method)
20. Postoperative 24hr analgesic consumption
21. Length of hospital stay
22. Anaesthetists appraisal of the SAFIRA system after completing all PNB procedures
23. Injection Pressure (How important is limiting LA opening/injection pressure in your own clinical practice: Not at all important/Quite unimportant/Neither unimportant or important /Quite important/Very important)
24. Ease of Use (Compared to usual practice SAFIRA is easy to use: Strongly disagree/Disagree /Neither disagree or agree/Agree/Strongly agree)
25. Effectiveness (Compared to usual practice SAFIRA is effective: Strongly disagree/Disagree /Neither disagree or agree/Agree/Strongly agree)
26. Drawbacks (Compared to usual practice SAFIRA has few drawbacks: Strongly disagree /Disagree/Neither disagree or agree/Agree/Strongly agree)
27. Workload (Compared to usual practice SAFIRA increases workload: Strongly disagree /Disagree/Neither disagree or agree/Agree/Strongly agree)
28. Preference (Compared to usual practice I prefer SAFIRA: Strongly disagree/Disagree/Neither disagree or agree/Agree/Strongly agree)

**Completion date**

22/03/2022

**Eligibility**

## **Key inclusion criteria**

PNB procedures involving patients aged >18 years referred for any type of elective surgery suitable for PNB and who are offered and agree to surgery that will be performed via any ultrasound guidance and/ or nerve stimulation single-injection PNB regional anaesthesia. Any type of PNB will be included in the study.

## **Participant type(s)**

Patient

## **Healthy volunteers allowed**

No

## **Age group**

Adult

## **Lower age limit**

18 years

## **Sex**

All

## **Key exclusion criteria**

1. PNB procedures involving patients aged < 18 years
2. PNB procedures involving (pre-scheduled) continuous PNB

## **Date of first enrolment**

22/11/2021

## **Date of final enrolment**

01/03/2022

## **Locations**

### **Countries of recruitment**

United Kingdom

England

### **Study participating centre**

**Queen Elizabeth Hospital King's Lynn Foundation Trust**

Gayton Road

King's Lynn

United Kingdom

PE30 4ET

## **Sponsor information**

## Organisation

Queen Elizabeth Hospital King's Lynn NHS Foundation Trust

## ROR

<https://ror.org/01m6k8878>

## Funder(s)

### Funder type

Industry

### Funder Name

Medovate Ltd.

## Results and Publications

### Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date

### IPD sharing plan summary

Data sharing statement to be made available at a later date

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">HRA research summary</a>			28/06/2023	No	No
<a href="#">Protocol file</a>	version 2.0		04/01/2022	No	No