

To assess the efficacy and tolerability of three antimalarial treatments in The Democratic Republic of Congo (DRC)

Submission date 22/09/2011	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 06/10/2011	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 10/02/2016	Condition category Infections and Infestations	☐ Individual participant data

Plain English summary of protocol

Background and study aims

Malaria is a serious tropical disease spread by mosquitoes. Antimalarial medication is used to prevent and treat malaria, with the first-line treatment being the first treatment given. We are carrying out a detailed assessment of the effectiveness of the current first-line antimalarial treatment, 5 years after its introduction. We will also assess the effectiveness of two other antimalarial treatments which could replace the first line in the near future, or be proposed as first-line treatment as part of a multiple first-line treatment strategy.

Who can participate?

Children aged 6 to 59 months with uncomplicated malaria

What does the study involve?

Participants are randomly allocated to one of the three treatments. The three treatments are amodiaquine-artesunate, dihydroartemisinin-piperaquine and artemether-lumefantrine. The study is open label, so the doctor and the patients both know which treatment they have been assigned to. All children are hospitalized for four days and given supervised treatment. The effectiveness of the treatment is measured and participants are followed up for 42 days after treatment.

What are the possible benefits and risks of participating?

The results of the study will enable the Ministry of Health to make informed decisions about whether the current national antimalarial treatment guidelines should be updated, and offer possible alternatives.

Where is the study run from?

The study will be conducted in Kinshasa, Democratic Republic of Congo.

When is the study starting and how long is it expected to run for?

September 2011 to December 2012

Who is funding the study?
Oxford University (UK)

Who is the main contact?
Prof Nick PJ Day

Contact information

Type(s)
Scientific

Contact name
Prof Nick PJ Day

Contact details
Faculty of Tropical Medicine
Mahidol University
3rd Floor
60th Anniversary Chalermprakit Building
420/6 Rajwithi Road
Bangkok
Thailand
10400

Additional identifiers

Protocol serial number
Version # 1.01 30 March 2011

Study information

Scientific Title
A randomised study to assess the efficacy and tolerability of three Artemisinin-based combination therapy (ACT) in The Democratic Republic of Congo (DRC)

Study objectives
This study will assess the efficacy of amodiaquine-artesunate for the treatment of uncomplicated *P. falciparum* malaria in children in Kinshasa, DRC, five years after its introduction as a first line treatment, and compare this with the efficacies of dihydroartemisinin-piperaquine and artemether-lumefantrine.

Ethics approval required
Old ethics approval format

Ethics approval(s)
1. Oxford Tropical Research Ethics Committee (OXTREC), 06/04/2011
2. Ethics Committee of Kinshasa University, Ministry of Higher Education and University, Democratic Republic of Congo (Ministère de l'Enseignement Supérieur et Universitaire, République Démocratique du Congo), 21/04/2011

Study design

Randomised three-arm open study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Malaria

Interventions

This is a randomised, open study, comparing three ACTs, dihydroartemisinin-piperaquine (DHA-PQ), artemetherlumefantrine (ALN) and amodiaquine-artesunate coformulated (AQ-AS), for the treatment of symptomatic uncomplicated falciparum malaria. The study is post-marketing, all three study drugs are in fact commercialized and used worldwide. Our aim is to monitor their efficacy and tolerability in this particular geographical area. Dosage and length of treatment are according to manufacturer. Children aged 6 to 59 months with uncomplicated P. falciparum malaria are randomly assigned to one of the three arms. All children are hospitalized for 3 days and given supervised treatment. Treatment efficacy is measured according to WHO (Methods for surveillance of antimalarial drug, WHO 2009). Patients are followed-up actively weekly for 42 days after treatment. Treatment allocation is concealed until recruitment is confirmed and the laboratory technicians reading all malaria smears have no knowledge of the treatment received by individual patients. Randomisation is in blocks of 15.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Amodiaquine-artesunate, dihydroartemisinin-piperaquine, artemether-lumefantrine

Primary outcome(s)

1. Clinical and parasitological cure by day 42 post-treatment
2. Treatment tolerability

Key secondary outcome(s)

1. The median parasite clearance time (PCT)
3. Fever clearance time (FCT)

Completion date

31/12/2012

Eligibility**Key inclusion criteria**

1. Children aged 3 to 59 months
2. Weight ≥ 5 kg [the minimum weight for treatment with artemether-lumefantrine (ALN)]
3. Mono-infection with *P. falciparum*
4. Parasitaemia of $\geq 2,000$ and $\leq 200,000$ asexual parasites per μL
5. Axillary temperature 37.5°C or history of fever in the preceding 24 hrs
6. Ability to swallow oral medication
7. Haemoglobin ≥ 5.0 g/dL
8. Parents/guardians agree to hospitalize the child for the length of treatment (34 days) and bring the patient for planned follow-up visits at day 7, 14, 21, 28, 35, 42
9. Signed consent from guardian / parents

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Neonate

Sex

All

Key exclusion criteria

1. Danger signs of severe malaria or signs of severe malaria (WHO 2000)
2. Children with severe malnutrition, marasmus or oedematous malnutrition (WHO 2006)
3. Febrile condition due to diseases other than malaria [e.g. measles, acute lower respiratory tract infection, severe diarrhoea with dehydration) or other known underlying chronic or severe diseases (e.g. cardiac, renal or hepatic diseases, human immunodeficiency virus (HIV) / acquired immune deficiency syndrome (AIDS)]
4. History of hypersensitivity reactions or contraindication to any medicine being tested
5. A clear history of adequate antimalarial treatment in the preceding 72 hours with drugs expected to be effective
6. Ongoing prophylaxis with drugs having antimalarial activity such as cotrimoxazole for the prevention of *Pneumocystis carinii* pneumonia in children born to HIV positive women

Date of first enrolment

08/09/2011

Date of final enrolment

31/12/2012

Locations**Countries of recruitment**

Congo, Democratic Republic

Thailand

Study participating centre
Mahidol University
Bangkok
Thailand
10400

Sponsor information

Organisation
The Centre for Tropical Medicine (UK)

ROR
<https://ror.org/052gg0110>

Funder(s)

Funder type
University/education

Funder Name
Oxford University (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2014		Yes	No