

Study of whole blood in frontline trauma

Submission date 18/11/2022	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 07/12/2022	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 18/03/2026	Condition category Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Every year, uncontrolled bleeding due to major injury (major traumatic haemorrhage) accounts for more than 2 million deaths worldwide and 4,500 deaths in England. Blood transfusion is an essential part of the treatment for severe bleeding, and any delay in starting transfusion can reduce the chances of survival. In the UK patients are often transfused blood at the scene of an incident before they arrive at hospital. Transfusion may involve different blood components, red blood cells (important for carrying oxygen around the body), plasma (contains essential proteins to help blood clot) and platelets (small cells that are essential for blood clot formation). Most UK air ambulances treat bleeding patients with a combination of red blood cells and plasma, which come in separate bags. However, carrying separate blood component bags introduces logistical challenges due to the additional weight the team needs to carry; increased complexity as several bags may need to be given to each patient; and a potential delay in transferring patients to hospital. Whole blood contains red cells, plasma and platelets all in one bag, as taken from a blood donor. Giving a blood transfusion of all of the components in a single bag could overcome these challenges. The aim of this study is to assess the clinical and cost-effectiveness of pre-hospital whole blood administration versus standard care for traumatic haemorrhage.

Who can participate?

Patients of any age who have suffered a traumatic injury, attended by a participating Air Ambulance Service clinical team, who require pre-hospital blood transfusion to treat major traumatic haemorrhage.

What does the study involve?

In this study, one group of patients will be given transfusions of red blood cells and plasma. The other group of patients will receive transfusions of whole blood. The effects of the two different treatments will be compared by looking at survival in the two groups and the amount of blood needed over the first 24 hours after injury. At the end of the study the researchers will determine which of the transfusion types is better (or whether there is no difference between them), and the cost-effectiveness and safety of giving whole blood transfusions compared to red blood cells.

What are the possible benefits and risks of participating?

There are no known risks or benefits linked to/attributed to taking part in this study, and there are no known additional risks in participating in the study compared to the risk associated with

transfusing blood components. Information collected as part of this trial may benefit patients in the future.

Where is the study run from?

NHS Blood and Transplant Clinical Trials Unit (UK)

When is the study starting and how long is it expected to run for?

March 2020 to June 2025

Who is funding the study?

The study has been funded by NHS Blood and Transplant, the Ministry of Defence and the following Air Ambulance Services:

1. Air Ambulance Kent Surrey Sussex (AAKSS)
2. Dorset and Somerset Air Ambulance (DSAA)
3. Essex and Herts Air Ambulance (EHAAT)
4. Hampshire and Isle of Wight Air Ambulance (HIOWAA)
5. Great North Air Ambulance (GNAAS)
6. Great Western Air Ambulance (GWAAC)
7. London's Air Ambulance (LAA)
8. Magpas Air Ambulance (Magpas)
9. North West Air Ambulance (NWAA)
10. Thames Valley Air Ambulance (TVAA)

Who is the main contact?

NHS Blood and Transplant Clinical Trials Unit, swift@nhsbt.nhs.uk

Contact information

Type(s)

Public

Contact name

Mrs Viona Rundell

Contact details

Clinical Trial Coordinator
NHS Blood and Transplant Clinical Trials Unit
Long Road
Cambridge
United Kingdom
CB2 0PT
+44 (0)1223 588091
Viona.Rundell@nhsbt.nhs.uk

Type(s)

Public

Contact name

Ms Joanne Lucas

Contact details

Trial Manager, Long Road
Cambridge
United Kingdom
CB2 0PT
+44 (0)1223 588175
swift@nhsbt.nhs.uk

Additional identifiers

Clinical Trials Information System (CTIS)
2021-006876-18

Integrated Research Application System (IRAS)
300414

Central Portfolio Management System (CPMS)
52435

Study information

Scientific Title

A multi-centre randomised controlled trial of the clinical and cost-effectiveness of pre-hospital whole blood administration versus standard care for traumatic haemorrhage

Acronym

SWiFT

Study objectives

Pre-hospital leukocyte-depleted whole blood transfusion is better than standard care (component transfusion) in reducing the proportion of participants who experience death or massive transfusion at 24 hours.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 12/09/2022, South Central - Oxford C Research Ethics Committee (Health Research Authority (Bristol), Ground Floor, Temple Quay House, 2 The Square, BS1 6PN, UK; +44 (0)207 104 8241, oxfordc.rec@hra.nhs.uk), ref: 22/SC/0072

Study design

Randomized treatment process-of-care management-of-care health-economic study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Traumatic haemorrhage

Interventions

Study design

A randomised controlled trial of pre-hospital whole blood versus red blood cells and plasma (non-blinded), for the treatment of major traumatic haemorrhage.

Type of participant to be studied

Patients (of any age) who require a blood transfusion in the pre-hospital setting, for the treatment of major traumatic haemorrhage.

Setting

Pre-Hospital Emergency Medicine.

Randomisation

Randomised boxes containing the trial intervention (either two units of whole blood or two units of red blood cells and two units of plasma) will be prepared in advance by the Transfusion Laboratory Teams. The boxes will be supplied to the participating Air Ambulance Services. If they attend to a patient who has suffered major trauma and requires blood transfusion, the team will open the trial intervention box and administer the contents to the patient, in accordance with standard local blood transfusion protocols. The time that the box was opened will be recorded and referred to as the randomisation time for the purposes of follow-up data collection. Informed consent will not be obtained prior to the initiation of treatment, due to the life-threatening nature of the patient's condition. Patients will be enrolled under an emergency waiver of consent, and informed consent will be sought (either directly from the participant, if they have capacity, or via a representative) as soon as practically possible.

Treatment

The intervention arm will be up to two units of whole blood (www.transfusionguidelines.org/red-book/annex-3/a3-6-whole-blood-leucocyte-depleted-for-clinical-studies). The control arm will be up to two units of red blood cells and up to 2 units of plasma (this is the current standard of care for the participating Air Ambulance Services). The plasma used in the control arm will either be fresh-frozen plasma (FFP) or LyoPlas (freeze-dried plasma). LyoPlas is classified as an IMP as it involves a manufacturing process. All other products used in this trial are blood components and fall under The Blood Safety and Quality Regulations. If bleeding continues after the trial intervention(s) have been administered, participants will receive further treatment as per standard of care.

Follow-up of participants

Patients will be reviewed as per standard clinical care. Data will be collected for the trial, for the secondary outcome measures, up to 90 days post-randomisation.

Safety reporting

Serious adverse events will be documented and reported up to 14 days post-treatment. The protocol lists events which are excluded from reporting (i.e. those which are recognised complications and consequences of major trauma).

Qualitative research

Alongside the randomised controlled trial, an 'implementation study' will be conducted. This will assess the acceptability and implementation of the intervention (whole blood). In this sub-study, qualitative methods will be used, involving interviews and focus groups with operational staff, patient representatives and blood donors.

Intervention Type

Other

Primary outcome(s)

The proportion of participants with traumatic haemorrhage who have died (all-cause mortality) or received a total of 10 or more units of any blood components in the first 24 hours from randomisation

Key secondary outcome(s)

Clinical Outcomes:

1. Individual components of the primary outcome: Proportion of participants who:
 - 1.1. Experienced all-cause mortality at 24 hours from randomisation
 - 1.2. Received a total of 10 or more units of any blood components in the first 24 hours of randomisation IV
2. All-cause mortality within 6 hours and separately 30 and 90 days of randomisation IV
3. Number of organ failure free days up to 30 days after randomisation, defined as the number of days free of advanced cardiovascular, advanced respiratory and advanced renal support. Each component of organ failure-free days will also be reported separately:
 - 3.1. Number of days free of advanced respiratory support
 - 3.2. Number of days free of advanced cardiovascular support
 - 3.3. Number of days free of advanced renal support
4. Days in critical care and separately in an acute care hospital (up to 90 days)
5. Units of each blood component received in the 24 hours after randomisation IV (including prehospital transfusions): whole blood (WB) and red blood cells (RBC), plasma, platelets and cryoprecipitate
6. Amount of cell salvage received at 24 hours (in ml) after randomisation IV
7. Number of participants receiving additional haemostatic agents received at 24 hours after randomisation IV: recombinant Factor VIIa, fibrinogen concentrate, prothrombin complex concentrate (PCC), tranexamic acid (TXA)
8. Presence of coagulopathy (defined as prothrombin time above the limits of a normal range) in the first sample taken on arrival at an acute care hospital
9. Acid-base disturbance measured by lactate, base excess and pH level in the first sample taken on arrival at an acute care hospital

Cost-Effectiveness Analysis Outcomes:

1. Incremental cost of the whole blood intervention
2. Hospital resource use to discharge or death
3. Health, social and wider care resource use to 90 days after randomisation
4. Health-related quality of life measured by EQ-5D-5L at 90 days after randomisation

Safety Outcomes:

1. Thrombosis (arterial and venous thrombosis) up to 30 days after randomisation
2. All transfusion reactions/events relating to pre-hospital blood components which have been reported to SHOT (Serious Hazards of Transfusion) occurring in the first 14 days after randomisation

Completion date

01/06/2025

Eligibility

Key inclusion criteria

1. Patient (of any age) who has suffered a traumatic injury
2. Attended by a participating Air Ambulance Service (AAS) clinical team
3. Requires pre-hospital blood transfusion to treat major traumatic haemorrhage

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

All

Sex

All

Total final enrolment

942

Key exclusion criteria

1. No intravenous or intraosseous access
2. Knowledge that the patient will object to being given blood transfusion for any reasons
3. Blood already administered on-scene, prior to the arrival of the participating Air Ambulance team

Date of first enrolment

15/12/2022

Date of final enrolment

12/09/2024

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre**Addenbrookes**

Addenbrookes Hospital

Hills Road

Cambridge

England

CB2 0QQ

Study participating centre
Aintree University Hospital
Lower Lane
Liverpool
England
L9 7AL

Study participating centre
Kent, Surrey & Sussex Air Ambulance Trust
Rochester City Airport
Maidstone Road
Chatham
Kent
England
ME5 9SD

Study participating centre
Alder Hey Children's Hospital
Eaton Road
West Derby
Liverpool
England
L12 2AP

Study participating centre
Bristol Royal Hospital for Sick Children
St. Michaels Hill
Bristol
England
BS2 8BJ

Study participating centre
Bristol Royal Infirmary
Marlborough Street
Bristol
England
BS2 8HW

Study participating centre

Dorset and Somerset Air Ambulance

Henstridge Airfield, the Marsh
Henstridge
Templecombe
England
BA8 0TN

Study participating centre

Dorset County Hospital

Dorset County Hospital
Princes Street
Dorchester
England
DT1 1TS

Study participating centre

East Surrey Hospital

Canada Avenue
Redhill
England
RH1 5RH

Study participating centre

Essex & Herts Air Ambulance Trust

Flight House
Earls Colne Business Park
Colchester
England
CO6 2NS

Study participating centre

The Great North Air Ambulance Service

Progress House
Urlay Nook Road
Eaglescliffe
Stockton-on-tees
England
TS16 0QB

Study participating centre

Great Western Air Ambulance Charity

County Gates
Ashton Road
Bristol
England
BS3 2JH

Study participating centre

Hampshire & Isle of Wight Air Ambulance Air Base

Hangar 2
Thruxton Airfield
Thruxton
Andover
England
SP11 8PW

Study participating centre

James Cook University Hospital

Marton Road
Middlesbrough
England
TS4 3BW

Study participating centre

John Radcliffe Hospital

Headley Way
Headington
Oxford
England
OX3 9DU

Study participating centre

Kings College Hospital

Mapother House
De Crespigny Park
Denmark Hill
London
England
SE5 8AB

Study participating centre

London's Air Ambulance

5th Floor
77 Mansell Street
London
England
E1 8AN

Study participating centre

Magpas The Emergency Medical Charity

Centenary House
St. Marys Street
Huntingdon
England
PE29 3PE

Study participating centre

Manchester Royal Infirmary

Cobbett House
Oxford Road
Manchester
England
M13 9WL

Study participating centre

North West Air Ambulance

North Mersey Business Centre
Woodward Road
Knowsley
England
L33 7UY

Study participating centre

Princess Alexandra Hospital

Hamstel Road
Harlow
England
CM20 1QX

Study participating centre

The Royal London Hospital

Alexandra House

London
England
E1 1BB

Study participating centre
Royal Preston Hospital
Sharoe Green Lane
Fulwood
Preston
England
PR2 9HT

Study participating centre
Royal Victoria Infirmary
Queen Victoria Road
Newcastle upon Tyne
England
NE1 4LP

Study participating centre
Salford Royal Hospital
Stott Lane
Eccles
Salford
England
M6 8HD

Study participating centre
Southampton General Hospital
Tremona Road
Southampton
England
SO16 6YD

Study participating centre
Southmead Hospital
Southmead Road
Westbury-on-trym
Bristol
England
BS10 5NB

Study participating centre

St Georges Hospital

Blackshaw Road

London

England

SW17 0QT

Study participating centre

St Marys Hospital

The Bays

South Wharf Road

London

England

W2 1BL

Study participating centre

Thames Valley Air Ambulance (oxford Road)

Stokenchurch House

Oxford Road

Stokenchurch

High Wycombe

England

HP14 3SX

Study participating centre

University Hospital (coventry)

Clifford Bridge Road

Coventry

England

CV2 2DX

Study participating centre

Royal Sussex County Hospital

Eastern Road

Brighton

England

BN2 5BE

Sponsor information

Organisation

NHS Blood and Transplant

ROR

<https://ror.org/0227qpa16>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

NHS Blood and Transplant; Grant Codes: 21-102-GEN

Alternative Name(s)

National Health Service Blood and Transplant, UK National Health Service Blood and Transplant, NHSBT

Funding Body Type

Government organisation

Funding Body Subtype

Local government

Location

United Kingdom

Funder Name

Ministry of Defence

Alternative Name(s)

MOD

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Funder Name

Air Ambulance Kent Surrey Sussex (AAKSS)

Funder Name

Dorset and Somerset Air Ambulance (DSAA)

Funder Name

Essex and Herts Air Ambulance (EHAAT)

Funder Name

Hampshire and Isle of Wight Air Ambulance (HIOWAA)

Funder Name

Great North Air Ambulance (GNAAS)

Funder Name

Great Western Air Ambulance (GWAAC)

Funder Name

London's Air Ambulance (LAA)

Funder Name

Magpas Air Ambulance (Magpas)

Funder Name

North West Air Ambulance (NWAA)

Funder Name

Thames Valley Air Ambulance (TVAA)

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analyzed during the trial will be available upon request from the NHSBT Clinical Trials Unit after de-identification (text, tables, figures and appendices) 9 months after publication and ending 5 years following article publication. Data will be shared with investigators whose use of the data has been assessed and approved by the trial review committee as a methodologically sound proposal. Please contact ctu@nhsbt.nhs.uk to apply for data access. Data availability will begin at 9 months after publication and end at 5 years after publication.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		17/03/2026	18/03/2026	Yes	No
Protocol article	RCT protocol	14/11/2023	16/11/2023	Yes	No
Protocol article	Implementation study protocol	05/02/2024	06/02/2024	Yes	No
HRA research summary			20/09/2023	No	No
Protocol file	version 1.1	03/05/2022	02/12/2022	No	No
Protocol file	version 3.1	17/02/2025	18/03/2026	No	No
Study website		11/11/2025	11/11/2025	No	Yes