

Primary prophylaxis with ciprofloxacin improves survival and reduces bacterial infections in patients with cirrhosis and ascites

Submission date 19/01/2007	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 09/02/2007	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 09/05/2019	Condition category Digestive System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Ruben Terg

Contact details
Caseros 2061
Buenos Aires
Argentina
1264

Additional identifiers

Protocol serial number
01/3045

Study information

Scientific Title
Primary prophylaxis with ciprofloxacin improves survival and reduces bacterial infections in patients with cirrhosis and ascites

Acronym

SBP - Spontaneous Bacterial Peritonitis

Study objectives

The aim of the present study was to assess the efficacy of long-term administration of ciprofloxacin for primary prophylaxis of Spontaneous Bacterial Peritonitis (SBP) in patients with cirrhosis and low protein concentration in ascitic fluid.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The protocol was approved by local ethic committees (ethics committee of Hospital Bonorino Udaondo and Hospital Alejandro Posadas) on the 6th March 2000.

Study design

Multicentre, randomised, double blind, placebo-controlled study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Cirrhotic patients with ascites

Interventions

Ciprofloxacin 500 mg a day or placebo, oral, during 12 months for intervention group and the control group.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Ciprofloxacin

Primary outcome(s)

To assess the efficacy of long-term administration of ciprofloxacin for primary prophylaxis of SBP in patients with cirrhosis and low protein concentration in ascitic fluid.

Key secondary outcome(s)

To assess the efficacy of long-term administration of ciprofloxacin for primary prophylaxis of other infections and mortality.

Completion date

01/12/2005

Eligibility

Key inclusion criteria

Diagnosis of cirrhosis was based in clinical, biochemical, ultrasonographic and histological criteria.

Inclusion criteria were:

1. Aged more than 18 and less than 80 years
2. Total protein in ascitic fluid less than 1.5 g/dl
3. Ability to give written informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Not Specified

Total final enrolment

100

Key exclusion criteria

1. A previous episode of SBP
2. Antibiotics in the previous 30 days
3. Pregnancy
4. Active gastrointestinal bleeding
5. Encephalopathy more than grade two
6. Hepatocarcinoma or other malignancies
7. Allergy to quinolones
8. Serum creatinine more than 3 mg/dl
9. Bacterial infection

Date of first enrolment

01/03/2000

Date of final enrolment

01/12/2005

Locations

Countries of recruitment

Argentina

Study participating centre

Caseros 2061
Buenos Aires
Argentina
1264

Sponsor information

Organisation

The Health Council of Investigation of the Government of Buenos Aires (Argentina)

Funder(s)

Funder type

Government

Funder Name

The Health Council of Investigation of the Government of Buenos Aires (Argentina) (ref: 01/3045)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/05/2008	09/05/2019	Yes	No