

A pilot study of cognitive analytic therapy (CAT) in stressed pregnant women with underlying anxiety and depression

Submission date 28/09/2007	Recruitment status No longer recruiting	☐ Prospectively registered
Registration date 28/09/2007	Overall study status Completed	☐ Protocol
Last Edited 29/06/2016	Condition category Mental and Behavioural Disorders	☐ Statistical analysis plan
		☐ Results
		☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Ian B. Kerr

Contact details

Adult Mental Health/ Psychotherapy
Southern Acute Day Hospital
Sevenairst Rd
Sheffield
United Kingdom
S20 1 NZ

Additional identifiers

Protocol serial number

N0071183793

Study information

Scientific Title

A pilot study of cognitive analytic therapy (CAT) in stressed pregnant women with underlying anxiety and depression

Study objectives

It is hypothesised primarily that treatment with CAT for a group of pregnant women with stressful anxiety, either alone or in conjunction with, depressive disorders will:

1. Result in a reduction of psychological distress and disability as measured by standard psychometric instruments and also
2. In a biological marker of stress as measured by salivary cortisol levels when compared to an untreated control group with similar disorders

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Pragmatic randomised controlled pilot study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Mental and Behavioural Disorders: Depression

Interventions

We propose to conduct an essentially pragmatic, randomised, controlled pilot aiming to evaluate the treatment effects of a brief (16-session) treatment with CAT in addition to treatment as usual in two maternal mental health outpatient settings (Sheffield Care Trust and St Thomas Hospital, London). This intervention would be compared to treatment as usual provided by primary care (e.g. GPs, midwives, health visitors) and secondary mental health services (e.g. community mental health teams, peri-natal psychiatric services).

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

The principal screening and outcome measure will be the Spielberger State/Trait Anxiety Measure which has been used in previous studies in this patient group as indicator of stress (Glover 2002).

Key secondary outcome(s)

Secondary standard outcome measures will include the Edinburgh Post Natal Depression Questionnaire (EPDS - a measure specifically validated for use in pregnancy as well as post-partum), the CORE brief routine outcome battery (an increasingly widely used general baseline indicator of subjective well being, risk of self harm, symptoms and functioning (Barkham et al

1998)) and the SF 36 short form of the Duke social support questionnaire. The latter will also be used in the economic evaluation. Use of self-report measures will minimise interviewer bias. In addition, sequential salivary cortisol levels will be used as secondary biological outcome measure. These will be sampled at the time points described above. They will be collected by obtaining samples of saliva 4 times per day (waking, after 30 minutes, after 3 hours, after 12 hours), on 2 days running in order to minimise day to day and diurnal variation. These samples will be collected on Salivette dental rolls, stored in a fridge and posted on to the laboratory (VG) at Imperial College. It is well established that cortisol is stable under such conditions.

Data on the prevalence of mother-infant interaction difficulties using a 5 minute video recording of mother-infant interaction (Murray et al 1996) will also be collected as well as any reductions observed in such difficulties in the treatment group. This will generate data on their prevalence initially and also enable future more extended studies to be planned if indicated. Similarly data will be collected on the incidence of subsequent post-natal psychiatric morbidity in these patient groups along with any change observed in the treatment group. This would similarly inform for further extended study of interventions aimed at reducing such morbidity.

Completion date

30/06/2009

Eligibility

Key inclusion criteria

Patients between 18 and 65 being referred from adult health services to perinatal psychiatric services (e.g. by midwives, obstetricians, GPs or mental health professionals) who appear at clinical diagnostic interview to be suffering from stress in the context of underlying anxiety, with or without associated depressive disorders will be offered entry into the trial at initial assessment.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

65 years

Sex

Female

Key exclusion criteria

1. Serious active substance abuse
2. Active psychotic symptoms
3. Age under 18 years old

Date of first enrolment

01/07/2006

Date of final enrolment

30/06/2009

Locations

Countries of recruitment

United Kingdom

Study participating centre

Adult Mental Health/ Psychotherapy

Sheffield

United Kingdom

S20 1 NZ

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

Funder(s)

Funder type

Government

Funder Name

Sheffield Health and Social Research Consortium

Funder Name

NHS R&D Support Funding

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration