

# To compare the pain relief from combined intrathecal bupivacaine-morphine with bupivacaine alone in lower limb surgery

<b>Submission date</b> 25/04/2022	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 12/05/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 05/05/2022	<b>Condition category</b> Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Inadequate pain medication during and after surgery may lead to long term pain after surgery.

This study was designed to ascertain the effect of morphine addition to bupivacaine in the management of postoperative pain in patients undergoing lower limb surgeries at the Tamale Teaching Hospital, Ghana.

### Who can participate?

Patients aged 16 – 60 years, undergoing lower limb surgery.

### What does the study involve?

Participants will be randomly allocated to receive bupivacaine alone or bupivacaine plus morphine during surgery. Follow up lasts 72 hours.

### What are the possible benefits and risks of participating?

**Benefits:** There were no immediate benefits for participation in the study. No gifts or money were given in exchange for the information obtained. However, data generated from this study will help policy makers in the Ministry of Health to improve the management of postoperative pain after Orthopaedic and Trauma surgeries.

**Risks:** Participants were informed that they may experience slight pain during the sample collection process. Should they suffer any injury during the collection process, members of the research team will take full responsibility and cover all expenses pertaining to treatment.

### Where is the study run from?

Tamale Teaching Hospital, Ghana.

### When is the study starting and how long is it expected to run for?

April 2021 to November 2021

Who is funding the study?  
Investigator initiated and funded

Who is the main contact?  
Sylvanus Kampo, sylvanuskampo@yahoo.com  
Fidelis Bayor, bayorfidelis@gmail.com

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Mr Fidelis Bayor

**ORCID ID**  
<https://orcid.org/0000-0002-2703-8856>

**Contact details**  
University of Ghana Medical School  
Accra  
Ghana  
24  
+233 249229949  
bayorfidelis@gmail.com

**Type(s)**  
Principal investigator

**Contact name**  
Dr Sylvanus Kampo

**Contact details**  
CK Tedam University of Technology and Applied Sciences  
Navrongo  
Ghana  
19  
+233 559355229  
sylvanuskampo@yahoo.com

## Additional identifiers

## Study information

**Scientific Title**  
Analgesic efficiency of intrathecal bupivacaine with morphine and bupivacaine alone in lower limb orthopaedic surgery. A comparative study

**Acronym**

IBM

### **Study objectives**

Bupivacaine with morphine induces better postoperative pain relief and improve patient outcomes in lower limb orthopaedic surgery

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Approved 30/09/2021, Ethical review board of the Kwame Nkrumah University of Science and Technology (University Post Office, Kumasi, Ghana; +233(0)205453785; chrpe@knust.edu.gh), ref: CHRPE/AP/416/21

### **Study design**

Randomised comparative controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Postoperative pain management in trauma patients

### **Interventions**

After meeting the inclusion criteria, eligible patients received 15 mg preservative free hyperbaric bupivacaine and 0.2 mg morphine or 15 mg bupivacaine alone. Patients were followed up 72 hrs for postoperative complications and patient outcome.

Each recruited patient was randomly assigned to one of two groups using a computer-generated random number table. Group A (n=30) represented those who received bupivacaine alone and Group B (n=30) represented those who received bupivacaine with morphine.

### **Intervention Type**

Drug

### **Phase**

Not Applicable

### **Drug/device/biological/vaccine name(s)**

bupivacaine, morphine

### **Primary outcome(s)**

1. Vital signs (pulse rate, blood pressure, oxygen saturation, and respiratory rate) of the patient were monitored and recorded for every 5 minutes for the first 30 minutes and then for every 15 minutes during surgery
2. Pain severity was measured every 4 hours for 72 hours immediately after the surgery on a 100-mm VAS, 0 mm = no pain, and 100 mm = intolerable pain.

### **Key secondary outcome(s)**

1. The episodes of postoperative nausea and vomiting (PONV) were identified by direct scheduled assessments or by a spontaneous complaint by the patients after the surgery. The incidence of PONV was recorded hourly for the first 4 hours and then 4 hourly for the next 24 hours using a 3 point ordinal scale (0 = none, 1 = nausea, 2 = vomiting). The incidence of PONV was calculated and categorized as early (0 – 4 hours) or delayed (5 – 24hours).
2. The incidence of pruritus was recorded every 4 hours for 48 hours after surgery on a four-point categorical scale as; 0 = no pruritus, 1 = mild, 2 = moderate, 3 = severe pruritus.
3. Overall perioperative satisfaction was evaluated on the day of discharged during an interview as; 4 = excellent, 3 = good, 2 = satisfactory, 1 = poor.

**Completion date**

30/11/2021

## Eligibility

**Key inclusion criteria**

1. Patients undergoing elective lower limb orthopaedic surgery
2. ASA status class I and II
3. Patients between 16 - 60 years of age
4. Patients who were on admission for 72 hours or more postoperatively

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Total final enrolment**

60

**Key exclusion criteria**

1. Pregnant or breastfeeding mothers
2. Patients with polytrauma
3. Glasgow coma scale <15
4. Opioids abusers
5. Expected post-surgical stay <24 hours
6. ASA physical status class III and IV
7. Patients allergic to bupivacaine and/morphine
8. Contraindication to spinal anaesthesia

**Date of first enrolment**

10/09/2021

**Date of final enrolment**

29/11/2021

## Locations

### Countries of recruitment

Ghana

### Study participating centre

Tamale Teaching Hospital

Box 1350TL

Tamale

Ghana

-

## Sponsor information

### Organisation

Platinum Orthopaedic and Spine Clinic

## Funder(s)

### Funder type

Other

### Funder Name

Investigator initiated and funded

## Results and Publications

### Individual participant data (IPD) sharing plan

Availability of data and materials: The datasets generated and/or analysed during the current study are not publicly available due to patient confidentiality but are available from the corresponding author on reasonable request.

sylvanuskampo@yahoo.com

### IPD sharing plan summary

Available on request