

Assessment and management of children aged 1 - 59 months presenting with wheeze and fast breathing: multicenter study in Pakistan and Thailand

Submission date 27/07/2004	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 28/07/2004	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 11/10/2007	Condition category Respiratory	☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Shamim Qazi

Contact details

World Health Organization
20 Avenue Appia
Geneva-27
Switzerland
CH 1211
qazis@who.int

Additional identifiers

Protocol serial number

WHO/CAH ID 99025

Study information

Scientific Title

Study objectives

Using current World Health Organization (WHO) guidelines, children with wheezing are being over-prescribed antibiotics and bronchodilators are under utilised. To improve the WHO case management guidelines, more data is needed about the clinical outcome in children with wheezing/pneumonia overlap.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from:

1. Institutional Review Board (IRB) of Pakistan Institute of Medical Sciences, Islamabad, Pakistan
2. IRB of Queen Sirikit National Institute of Child Health, Bangkok, Thailand
3. World Health Organization (WHO) Ethical Review Committee

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Wheeze and fast breathing

Interventions

1. Inhaled Salbutamol and reassessment after up to three cycles of bronchodilator therapy repeated at 15 minute interval if necessary
2. Metered Dose Inhaler (MDI) with spacer device

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Proportions of children aged 1 - 59 months with auscultatory wheeze and fast breathing or lower chest indrawing:

1. Respond to up to three cycles of inhaled salbutamol: response defined as no fast breathing or lower chest indrawing present
2. Fail therapy at day 3 or days 5 - 7 of the initial successful bronchodilator therapy with inhaled salbutamol? Therapy failure defined as:
 - 2.1. Relapse: develop fast breathing or chest indrawing afresh, that does not respond to three cycles of inhaled salbutamol

2.2. Development of any danger sign (except wheezing and fever in young infant)

2.3. Death

2.4. Severe adverse reaction to salbutamol

Key secondary outcome(s)

In children aged 1 - 59 months with auscultatory wheeze and fast breathing or lower chest indrawing:

1. Proportion with audible wheeze at initial assessment

2. Proportion of responders to up to three cycles of inhaled salbutamol associated with:

2.1. Age

2.2. Respiratory Syncytial Virus (RSV) isolation

2.3. Season

2.4. Number of previous wheezing episodes

2.5. Audible versus auscultatory wheeze

2.6. Family history of asthma

3. Proportion of relapses in children who showed initial improvement associated with:

3.1. Age

3.2. RSV isolation

3.3. Season

3.4. Number of previous wheezing episodes

3.5. Audible versus auscultatory wheeze

3.6. Family history of asthma

4. Received any antibiotic for this or any concurrent illness before follow-up

5. Loss to follow-up

6. Withdrawal of consent

Completion date

01/04/2002

Eligibility

Key inclusion criteria

1. Age 1 to 59 months

2. Audible/auscultatory wheeze

3. Respiratory rate above age specific cut off point or lower chest indrawing

4. Classified as no pneumonia with wheeze

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

1 Months

Upper age limit

59 Months

Sex

All

Key exclusion criteria

1. Presence of danger sign:

1.1. Up to two months: stopped feeding well, drowsy, convulsions, stridor in a calm child, fever

1.2. 2 - 59 months: convulsions, clinically severe malnutrition, unable to drink

2. Antibiotics during last 48 hours

Date of first enrolment

01/05/2001

Date of final enrolment

01/04/2002

Locations

Countries of recruitment

Pakistan

Switzerland

Thailand

Study participating centre

World Health Organization

Geneva-27

Switzerland

CH 1211

Sponsor information

Organisation

The Department of Child and Adolescent Health (CAH)/World Health Organization (WHO)
(Switzerland)

ROR

<https://ror.org/01f80g185>

Funder(s)

Funder type

Research organisation

Funder Name

The Department of Child and Adolescent Health (CAH)/World Health Organization (WHO) (Switzerland)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results	01/11/2004		Yes	No