

Prospective randomised study of the influence of a lung recruitment strategy on oxygenation in the post-anaesthetic care unit (PACU)

Submission date 29/09/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 29/09/2006	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 01/04/2010	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Stuart Greenhill

Contact details
Department of Anaesthesia
Great George Street
Leeds
United Kingdom
LS1 3EX
+44 07766432470
greenhill@doctors.org.uk

Additional identifiers

Protocol serial number
N0436176302

Study information

Scientific Title

Study objectives

The aim of the study is to investigate whether lung recruitment manoeuvres performed near the end of surgery can have beneficial effects on oxygenation that persist into the early postoperative period.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Surgery: Anaesthesia

Interventions

Added 01/04/10 (see publication for further details):

Adult patients undergoing elective surgery requiring tracheal intubation and an arterial catheter were randomized to receive:

1. Lung recruitment manoeuvre of 40 cm H₂O for 15 s, 30 min before the end of anaesthesia, followed by 10 cm H₂O of PEEP and then 10 cm H₂O of CPAP from return of spontaneous breathing until extubation
2. No lung recruitment manoeuvre, ≤ 5 cm H₂O PEEP, and no CPAP.

Arterial blood gases taken at randomization and 1 h after extubation.

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome(s)

Difference between intra-operative and post-operative alveolar to arterial oxygen difference (A-aDO₂)

Key secondary outcome(s)

Not provided at time of registration

Completion date

27/02/2007

Eligibility

Key inclusion criteria

Elective adult patients (lower age limit 18 years, no upper age limit) undergoing operative procedures lasting longer than 45 minutes requiring tracheal intubation muscle relaxation and the insertion of an arterial catheter on clinical grounds who will postoperatively be admitted to the post anaesthetic care unit.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

Does not match inclusion criteria

Date of first enrolment

27/02/2006

Date of final enrolment

27/02/2007

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Department of Anaesthesia

Leeds

United Kingdom

LS1 3EX

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2006 Update - Department of Health

Funder(s)

Funder type

Government

Funder Name

Leeds Teaching Hospitals NHS Trust (UK)

Funder Name

NHS R&D Support Funding

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/05/2010		Yes	No