

Ultrasonographic evaluation of the submandibular space compliance to predict difficult airway in anaesthesia

Submission date 21/11/2016	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 25/11/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 09/08/2018	Condition category Other	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Oro-tracheal intubation is the placement of a flexible plastic tube through the mouth into the trachea (windpipe) to maintain an open airway. This is required during general anaesthesia, when the patient is connected to a breathing machine (ventilator). Managing the airway is an important skill for anaesthesiologists. It is important for the anaesthesiologist to assess the airway correctly in order to avoid hypoxemic events (low blood oxygen levels) and their consequences. The aim of this study is to find out whether an assessment of the anatomy of the airway can be used to predict the occurrence of airway difficulties in patients undergoing surgery.

Who can participate?

Patients aged over 18 undergoing elective general anaesthesia with oro-tracheal intubation

What does the study involve?

On the day before they undergo surgery, participants undergo an ultrasound scan of the anterior (front) region of the neck to look at the airway structures. Airway difficulties are assessed when the participants are intubated during general anaesthesia.

What are the possible benefits and risks of participating?

The results of this study could help us predict airway difficulties, thereby decreasing the risk of illness or death due to low blood oxygen levels. There are no risks for the participants as ultrasound is non-invasive and does not involve radiation.

Where is the study run from?

The Clinical Emergency County Hospital Cluj (Romania)

When is the study starting and how long is it expected to run for?

October 2016 to June 2017

Who is funding the study?
Investigator initiated and funded

Who is the main contact?
Dr Cristina Petrisor
petrisor.cristina@umfcluj.ro

Contact information

Type(s)
Scientific

Contact name
Dr Cristina Petrisor

Contact details
No. 14, Eftimie Murgu Street
Cluj-Napoca
Romania
400604
+40 (0)722 262 286
petrisor.cristina@umfcluj.ro

Additional identifiers

Protocol serial number
18637/20.09.2016

Study information

Scientific Title
Ultrasonographic evaluation of the submandibular space compliance to predict difficult airway in anaesthesia: a prospective diagnostic study

Acronym
HIO-MENTO US

Study objectives
To investigate the ability of ultrasound to evaluate the submandibular space compliance as a predictor for difficult airway in anaesthesia.

Ethics approval required
Old ethics approval format

Ethics approval(s)
The Ethics Committee of the Clinical Emergency County Hospital Cluj, 20/09/2016, ref: 18637

Primary study design
Observational

Study design

Prospective diagnostic study

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Management of the difficult airway

Interventions

After the Ethics Committee approval and signing the informed consent forms, adult patients scheduled to undergo general anaesthesia with oro-tracheal intubation are prospectively included. Exclusion criteria are morbid obesity and rapid sequence inductions. Each patient will undergo an ultrasound evaluation of the airway anatomy, focusing on submandibular space compliance. The HMDs (hyo-mandibular distances) are measured in neutral, sniffing, and maximal hyperextended positions, in mid-sagittal plane, using a curvilinear ultrasound transducer. The airway ultrasound is performed one day prior to the surgical intervention under general anaesthesia. All intubations are performed in sniffing position, with Macintosh curved laryngoscope blades, with no external laryngeal manipulation. The Cormack grade is registered on first attempt of laryngoscopy. Grades 3 and 4 Cormack-Lehane are considered difficult airway. ROC (receiver operating characteristics) curve analysis is used.

Intervention Type

Other

Primary outcome(s)

Submandibular space compliance (the ratio for the hyo-mental distance in maximal hyperextended position and sniffing position to the same distance in neutral position), measured by airway ultrasound one day before surgery

Key secondary outcome(s)

Laryngeal view, assessed using the Cormack grade at induction of anaesthesia during the first laryngoscopy attempt

Completion date

30/06/2017

Eligibility**Key inclusion criteria**

Patients aged over 18 undergoing elective general anaesthesia with oro-tracheal intubation

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. Rapid sequence induction
2. External laryngeal manipulation during laryngoscopy
3. Emergency surgery
4. Videolaryngoscopy

Date of first enrolment

01/10/2016

Date of final enrolment

30/06/2017

Locations**Countries of recruitment**

Romania

Study participating centre

The Clinical Emergency County Hospital Cluj

No.3-5, Clinicilor Street

Cluj-Napoca

Romania

400006

Sponsor information**Organisation**

The Clinical Emergency County Hospital Cluj

ROR

<https://ror.org/05j4kzc41>

Funder(s)**Funder type**

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

All data will be registered in a database and, provided patient confidentiality is protected, can be made available by contacting Cristina Petrisor (petrisor.cristina@umfcluj.ro).

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2018		Yes	No