

Novel approaches in in-vitro fertilisation (IVF)

Submission date 08/03/2006	Recruitment status No longer recruiting	☐ Prospectively registered
Registration date 08/03/2006	Overall study status Completed	☐ Protocol
Last Edited 27/10/2022	Condition category Pregnancy and Childbirth	☐ Statistical analysis plan
		☒ Results
		☐ Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Esther Heijnen

Contact details
Department of Reproductive Medicine
University Medical Center Utrecht
Heidelberglaan 100
Utrecht
Netherlands
3584 CX
-
esther.heijnen@organon.com

Additional identifiers

Protocol serial number
NTR587

Study information

Scientific Title
Novel approaches in in-vitro fertilisation (IVF)

Study objectives

Strategies involving shorter and milder ovarian stimulation protocols and single embryo transfer may allow for more in-vitro fertilisation (IVF) cycles in the same period of time, resulting in similar term live birth rate per treatment period despite a minor reduction in birth rate per treatment cycle.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Received from the local medical ethics committee

Primary study design

Interventional

Study design

Multicentre, randomised, active controlled, parallel group trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Infertility

Interventions

Patients were randomly assigned to undergo either:

1. Mild treatment (mild ovarian stimulation with gonadotropin-releasing hormone [GnRH] antagonist co-treatment combined with single embryo transfer), or
2. Standard treatment (stimulation with a GnRH agonist long-protocol and transfer of two embryos)

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Gonadotropin-releasing hormone (GnRH)

Primary outcome(s)

The primary outcome parameters chosen for this study were:

1. Pregnancy within one year of treatment after randomisation leading to term (greater than 37 weeks gestation) live birth
2. Total costs per couple and child up to six weeks after expected delivery
3. Patient discomfort/distress during IVF treatment

Key secondary outcome(s)

No secondary outcome measures

Completion date

01/03/2005

Eligibility

Key inclusion criteria

1. Patients with an indication for IVF or IVF/intracytoplasmic sperm injection (ICSI) based on tubal, male or unexplained infertility were recruited
2. Patients less than 38 years with a normal menstrual cycle (cycle length between period 25 - 35 days) and without severe obesity or underweight (body mass index [BMI] 18 - 28 kg/m²) were eligible for the study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

Female

Total final enrolment

404

Key exclusion criteria

Does not comply with the above inclusion criteria.

Date of first enrolment

01/02/2002

Date of final enrolment

01/03/2005

Locations

Countries of recruitment

Netherlands

Study participating centre

Department of Reproductive Medicine

Utrecht

Netherlands

3584 CX

Sponsor information

Organisation

University Medical Centre Utrecht (UMCU) (The Netherlands)

ROR

<https://ror.org/04pp8hn57>

Funder(s)

Funder type

Research organisation

Funder Name

The Netherlands Organisation for Health Research and Development (ZonMw) (The Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		03/03/2007		Yes	No
Results article	Patient discomfort, risks and costs	05/04/2008	27/10/2022	Yes	No
Other publications	Psychological impact	01/09/2007		Yes	No