

Can ambulance paramedics use FRAX (the WHO Fracture Risk Assessment Tool) to help GPs improve future fracture risk in patients that fall?

Submission date

13/03/2014

Recruitment status

No longer recruiting

☐ Prospectively registered

☒ Protocol

Registration date

13/03/2014

Overall study status

Completed

☐ Statistical analysis plan

☒ Results

Last Edited

12/10/2018

Condition category

Injury, Occupational Diseases, Poisoning

☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Shane Clarke

Contact details

Rheumatology Unit

The Courtyard

Old Building, Bristol Royal Infirmary

Marlborough Street

Bristol

United Kingdom

BS2 8HW

-

s.clarke@bristol.ac.uk

Additional identifiers

Protocol serial number

14057

Study information

Scientific Title

Can ambulance paramedics use FRAX (the WHO Fracture Risk Assessment Tool) to help GPs improve future fracture risk in patients that fall?

Study objectives

Broken bones are painful, often disabling, and cost the NHS £2 billion annually. Tablets reduce the risk of having a fracture in the most vulnerable but the number of people at high risk taking the right treatment (about 1 in 3) is unacceptably low.

Part of the problem is poor communication between healthcare providers; for example, GPs are seldom told about their patients who fall by the attending ambulance service. Falls and fracture are closely linked. Each fall that results in an ambulance call might represent an important opportunity to assess a patients fracture risk using a computer programme called FRAX®.

The proposed project is a feasibility study to help us design a full trial. The full trial will find out if ambulance crew can collect information from people that fall and help GPs target treatment for osteoporosis at those patients most likely to have a future fracture.

In this feasibility study we want to find out whether ambulance staff can obtain the necessary information to estimate a patients fracture risk, and if the GPs will act on the information given to them. We also require information to help us work out how many patients we will need to recruit, and might be at a high risk of fracture. It is also essential to ensure the study methods are acceptable to patients, ambulance staff and GPs before we plan the main trial.

Our study involves ambulance staff asking patients additional questions after they have fallen over, and the design includes a control group, whose members will not be informed of their fracture risk. These ethical issues were considered by members of the public.

The proposal was felt to be acceptable, with real potential to improve osteoporosis care. A large Medical Research Council study has already successfully used a similar approach.

More details can be found at: <http://public.ukcrn.org.uk/search/StudyDetail.aspx?StudyID=14057>

On 04/06/2014 the anticipated end date was changed from 28/03/2014 to 01/07/2014.

Ethics approval required

Old ethics approval format

Ethics approval(s)

12/SC/0604

Study design

Randomised; Interventional; Design type: Screening, Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Primary Care Research Network for England, Injuries and Emergencies; Subtopic: Not Assigned, Injuries and Emergencies (all Subtopics); Disease: Injuries and Emergencies, All Diseases

Interventions

Calculate future fracture risk: Paramedics will calculate future fracture risk in patients that fall; Transmit fracture risk GP, Paramedics will transmit calculated future fracture risk to patients GPs; Follow Up Length: 3 month(s); Study Entry : Single Randomisation only

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Patients on treatment; Timepoint(s): Proportion of patients taking new treatment to reduce future fracture risk

Key secondary outcome(s)

Appropriate investigations; Timepoint(s): The proportion of patients in control and intervention group that are referred for DXA or blood test

Completion date

01/07/2014

Eligibility

Key inclusion criteria

Men and women aged over 50 who fall (inside or outside their place of residence), call an ambulance, and are attended by study paramedics

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

All

Key exclusion criteria

1. Patients who, in the paramedics opinion, are medically unstable and for whom it would not be appropriate to delay treatment to ask study questions (for example where the paramedics suspect a cardiac incident).
2. Patients who have fallen in a public place and in the opinion of the treating paramedics should be conveyed without delay.

3. Patients who are deemed to lack capacity (according to the Mental Health Act [2005]) but for whom there is no available carer or consultee.
4. Patients who are admitted to hospital for 24 hours or more. These patients will need to be excluded by the RF prior to consent (pre-existing service arrangements cover inpatients and our principal target population is in primary care).

We will continue to monitor patients who fall twice or more during the study period, but count the first fall as the index event. Only the first eligible fall will trigger entry to the study.

Date of first enrolment

28/06/2013

Date of final enrolment

01/07/2014

Locations

Countries of recruitment

United Kingdom

England

Study participating centre**Rheumatology Unit**

Bristol

United Kingdom

BS2 8HW

Sponsor information

Organisation

University Hospitals Bristol NHS Foundation Trust (UK)

ROR

<https://ror.org/04nm1cv11>

Funder(s)

Funder type

Government

Funder Name

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	20/04/2015		Yes	No
Protocol article	protocol	03/09/2014		Yes	No