

Hormonal profiles modification, lower hunger scores, better anthropometric outcomes and reduced risk factors of the metabolic syndrome following a weight loss diet with carbohydrates eaten only at dinner

Submission date

23/11/2009

Recruitment status

No longer recruiting

Registration date

03/12/2009

Overall study status

Completed

Last Edited

06/02/2014

Condition category

Nutritional, Metabolic, Endocrine

Prospectively registered

Protocol

Statistical analysis plan

Results

Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Zecharia Madar

Contact details

P.O Box 12

Rehovot

Israel

76100

+972 (0)89489008

esofer1@012.net.il

Additional identifiers

Study information

Scientific Title

A randomised clinical trial examining leptin, ghrelin and adiponectin diurnal profiles modifications, hunger and satiety scores, anthropometric, biochemical and inflammation parameters following a weight loss diet with carbohydrates eaten only at dinner.

Study objectives

Obesity is often accompanied by uncontrolled hunger, insulin resistance and by the metabolic syndrome. The adipose tissue, "the energy storage site of the body", is also an endocrine organ that synthesizes and secretes a variety of adipocytokines. This includes hormones regulating hunger and satiety and associated with the development of insulin resistance, the metabolic syndrome and inflammation.

Hypotheses:

1. A weight loss diet with carbohydrates eaten only at dinner (the experimental diet) will lead to modifications of the typical diurnal pattern of leptin, and to higher relative concentrations throughout the day, helping experimental diet participants to experience satiety during the day and to better adhere to their diets.
2. Ghrelin's diurnal pattern will be inverted too, leading to the appearance of hunger sensations later in the day.
3. The experimental diet will increase adiponectin concentrations throughout the day, leading to improved insulin resistance, diminished symptoms of the metabolic syndrome and better inflammation profiles.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved by the Regional Committee for Human Experimentation, Kaplan Hospital, Israel in accordance with the Helsinki declaration of 1975 (revised in 1983). (ref: 024/2006)

Study design

Single centre randomised controlled parallel group trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Obesity; metabolic syndrome

Interventions

100 officers enrolled to a randomised single centre controlled interventional diet study. 78 Individuals met study criteria and were randomly allocated to the experimental/control groups.

A standard low calorie diet (20% protein, 30-35% fat, 45-50% carbohydrates, 1300-1500 kcal) providing carbohydrates only at dinner (Experimental diet) or a standard low calorie diet (20% protein, 30-35% fat, 45-50% carbohydrates, 1300-1500 kcal), providing carbohydrates

throughout the day (control diet) was consumed for 6 months
Blood samples were taken and the participants filled out hunger-satiety scales every 4 hours between 8:00-20:00 at day 0, 7, 90 and 180.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

1. Leptin, ghrelin (total) and adiponectin (High Molecular Weight) (Linco research sandwich ELISA kits)
2. Insulin (Abbot Microparticle Enzyme Immunoassay test kits)
3. Hunger-Satiety Score (H-SSc) a scale of descriptions from hunger to satiety (1= starving, 10= devastatingly full).
4. Glucose (Olympus enzymatic UV test kits)
5. Insulin resistance (Homeostasis Model Assessment)
6. Cholesterol, HDL-cholesterol, LDL-cholesterol and triglycerides (Olympus enzymatic colour test kits)
7. C-reactive protein (CRP) (Olympus Immunoturbidimetric test kits)
8. TNF- α (Human Serum [HS]) and IL-6 (HS) (R&D systems sandwich ELISA kits)

Key secondary outcome(s)

1. Weight, abdominal circumference, and percentage of body fat were measured regularly and on day 0, 7, 90,180.
2. "Urge to eat" and "preoccupation with thoughts about food".

Completion date

09/09/2007

Eligibility

Key inclusion criteria

1. Police officers from the Israeli Police Force (men and women)
2. Age 25-55
3. BMI >30

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Cardiovascular diseases
2. Hypertension
3. Diabetes mellitus or other primary diseases
4. Followed any type of diet regime within a year prior to the study
5. Pregnancy

Date of first enrolment

22/05/2006

Date of final enrolment

09/09/2007

Locations

Countries of recruitment

Israel

Study participating centre

P.O Box 12

Rehovot

Israel

76100

Sponsor information

Organisation

The Hebrew University of Jerusalem (Israel)

ROR

<https://ror.org/03qxff017>

Funder(s)

Funder type

Other

Funder Name

Meuhedet Medical Services (Israel)

Funder Name

Kaplan Medical Center, Rehovot (Israel)

Funder Name

Israeli Police Force (Israel)

Funder Name

Israel Diabetes Association (Israel)

Funder Name

Israel Lung and Tuberculosis Association (Israel)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2013		Yes	No