

Pain relief after instrumented spinal surgery trial

Submission date 02/09/2023	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 03/11/2023	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 05/03/2026	Condition category Surgery	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Spinal surgery is common in the NHS and can result in severe postoperative pain limiting recovery and rehabilitation. Pain after instrumented lumbar spinal surgery is severe and can persist for many weeks, with a mean length of hospital stay of 4.7 days. Severe postoperative pain can delay early mobilisation, with potential complications such as venous thromboembolism and infection, all of which carry costs for the NHS. The treatments under evaluation each carry a different harm:benefit profile. Multimodal analgesia with strong opioids (Usual Care) is standard treatment but there is some evidence from systematic reviews that the alternatives offer superior pain relief. Both intrathecal opioids and ESB are currently used in some centres, but without a rational basis for treatment selection in scientific evidence.

The PRAISE trial aims to investigate the clinical and cost effectiveness of three approaches to postoperative pain relief following lumbar spine surgery: Control (Usual care), Intervention 1 (Intrathecal Opioid injection) or Intervention 2 (Erector Spinae plane Block).

Who can participate?

Participants aged over 16 years scheduled for elective posterior lumbar-instrumented spinal surgery who are able to give informed consent

What does the study involve?

Patients will be randomised 1:1:1 to one of the three approaches above.

The primary outcome is back pain on moving around bed (sitting up and/or turning) on a 0-100 VAS at 24 hours post-surgery. Secondary outcomes include EQ-5D-5L, back pain at rest, leg pain, QoR-15, cumulative postoperative opioid consumption, quality of life, time to mobilisation, length of stay, adverse events and healthcare resource use. Parallel economic evaluation with the trial will estimate incremental cost-effectiveness ratios. Outcomes will be collected at baseline, on admission to theatre recovery, at 6, 24 and 72 hours after surgery, and at routine 6-8 week follow up.

What are the possible benefits and risks of participating?

Benefits:

Not provided at time of registration

Risks:

In addition to surgical complications attendant with spinal procedures, potential harms differ somewhat for each group allocation:

ESP Block complications - ESPB has very rare but potentially serious complications of delivery of local anaesthetic including intravenous injection, local anaesthetic toxicity, intraneural injection, infection at needle insertion site and pneumothorax. SAEs will be continually monitored throughout the trial accordingly and patients will undergo a detailed risk-benefit conversation with a consultant anaesthetist prior to consent and randomisation.

Usual Care UC also carries some risk of inadequate analgesia (severe, acute postoperative pain) postoperative nausea and vomiting (PONV), sedation. Intrathecal Opioid (ITO) also carries risk of Respiratory depression, pruritus (itching) and urinary retention - but both these procedures are standard of care treatments.

Where is the study run from?

Sheffield Teaching Hospitals NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for?

August 2023 to June 2026

Who is funding the study?

National Institute for Health and Care Research (NIHR) (UK)

Who is the main contact?

Marie Hyslop, m.c.hyslop@sheffield.ac.uk

Contact information

Type(s)

Scientific

Contact name

Ms Marie Hyslop

Contact details

Clinical Trials Research Unit The University of Sheffield

Regent Court

30 Regent Street

Sheffield

United Kingdom

S1 4DA

-

m.c.hyslop@sheffield.ac.uk

Type(s)

Principal investigator

Contact name

Dr Matthew Wilson

Contact details

ScHARR
The University of Sheffield
Regent Court
30 Regent Street
Sheffield
United Kingdom
S1 4DA
+44 114 222 0836
m.j.wilson@sheffield.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

1008666

Protocol serial number

STH22261

Study information

Scientific Title

Pain Relief After Instrumented Spinal surgery trial

Acronym

PRAISE

Study objectives

Primary objective:

A randomised controlled trial powered to test the hypothesis that enhanced anaesthetic techniques, intrathecal opioids or Erector Spinae plane Block, improve postoperative back pain on moving around the bed (sitting up and/or turning) at 24 hours by at least 10 points (0-100 Visual Analogue Scale) compared to usual care in patients undergoing spinal surgery +/- decompression.

Secondary objectives:

To determine:

1. if there is a difference in cumulative opioid consumption after surgery, at 24 hours and 72 hours (or discharge) between UC, ITO and ESB;
2. if there is a difference in quality of recovery between the three intervention groups using the Quality of Recovery Scale;
3. if there is a difference in the need for further clinical intervention between UC, ITO and ESB;
4. if there is a difference in time to mobilisation after surgery and length of hospital stay (readiness for discharge) between UC, ITO and ESB;
5. if there is a difference in readmissions between discharge and routine 6-8 week postoperative follow up, between UC, ITO and ESB;

6. if there is a difference in quality of life between the three intervention groups;
7. if there is a difference in healthcare resource use between three groups and to compare the adverse events between UC, ITO and ESB. Also feasibility and Health Economic objectives.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 02/11/2023, London - Dulwich Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 (0)2071048236; westminster.rec@hra.nhs.uk), ref: 23/LO/0811

Study design

Multicentre parallel-group superiority patient-blind individual participant-randomized controlled trial

Primary study design

Interventional

Study type(s)

Efficacy, Safety

Health condition(s) or problem(s) studied

Patients scheduled for elective, posterior lumbar-instrumented spinal surgery +/- decompression.

Interventions

Patients will be identified and screened for eligibility by their surgical clinical team when the decision for lumbar spinal surgery has been taken. Patients will be provided with written information materials and referred to an anaesthetist for an in-depth trial discussion. There will be a clear explanation of study procedures including the interventions and associated risks, and the opportunity for patients to have any questions. Consent will only be sought after the discussion with the anaesthetist has taken place. Recruitment will be aligned around the standard care pre-operative assessment appointment, which will take place sometime after the patient has been listed for surgery, allowing sufficient time for patients to consider the information and decide whether they would like to take part.

Once eligibility has been confirmed and consent acquired, the participant will be randomly allocated to one of the following groups: (1) Usual Care; (2) Intrathecal Opioids (ITO); (3) Erector Spinae plane Block (ESB) using the Sheffield CTRU online randomisation system (SCRAM). Participants will be allocated to their intervention using minimisation with a random element and the following factors ensuring baseline balance: site; levels of fusion (1 level vs. >1 level (2-3)), receiving Step 3 opioid therapy at randomisation (yes vs no). Trial participants can only be randomised if staff trained in delivery of all interventions are available.

The trial interventions are as follows:

1. Usual Care (with local wound infiltration):

Involves analgesia administered at the time of surgery whilst under General Anaesthesia and in the immediate post-operative period. Intravenous opioid, Paracetamol, NSAID (if not contraindicated). This will be administered by the anaesthetist providing clinical care. Trial

participants will also receive local infiltration of surgical incision with Local Anaesthetic (Levo-Bupivacaine) administered by the Operating Surgeon

How: Intravenous injection during surgery in Operating Theatre. Local anaesthesia administered at surgical closure.

Timing: Intravenous drugs administered intra-operatively. Local anaesthetic infiltration at completion of surgery.

Dose: Intravenous analgesia administered at the discretion of the Consultant anaesthetist: Local Anaesthetic Infiltration: up to maximum dose Levo-Bupivacaine 100mg

Tailoring: Opioid Analgesia Drug Type and Dose administered according to Anaesthetist's clinical opinion and local practice. Paracetamol as per local standard of care, NSAID if not contraindicated.

Modifications: Local anaesthetic infiltration volume adjusted to maximum dose of Levo-Bupivacaine 2 mg/kg to avoid toxicity.

Accountability: All drugs administered during surgery are routinely recorded in anaesthetic record.

2. Usual Care plus Intrathecal Opioid (with local wound infiltration)

Involves pain relief with central (neuraxial) opioid pain relief, reducing onward transmission of nociceptive stimulus to reduce patient pain sensation for sustained period post operatively.

Intrathecal Opioid Analgesia administered by operating surgeon at the time of surgery whilst under General Anaesthesia. Morphine (Preservative Free); diamorphine according to local clinical practice guidelines. ITO plus Usual Care as above.

How: Intrathecal Opioid administered by Operating Surgeon in Operating Theatre. Intrathecal injection under direct vision, via narrow gauge "pencil point" spinal needle via Ligamentum Flavum. ITO is permitted to be administered by an appropriately trained surgeon who does this procedure as part of their clinical care. The usual care and intrathecal opioid interventions are routinely used in the NHS and require no specific training.

Timing: At the time of surgery. Local wound infiltration at completion of surgery.

Dose: Preservative Free Morphine: 0.1 mg (minimum dose); or Diamorphine (dose range 0.2mg to 0.4mg).

Modifications: Choice of drug (morphine vs diamorphine) dictated by local practice and requirements for post-operative care. Paracetamol as per local standard of care. Omit NSAID if contraindication.

Accountability: Morphine or Diamorphine must be dispensed from research pharmacy.

Confirmation of intrathecal administration by cerebrospinal fluid aspiration prior to injection. All drugs administered during surgery are routinely recorded in anaesthetic records, which will inform the per protocol analysis.

Intrathecal Opioid Group will receive Usual Care, as described above plus the Intervention described above.

3. Usual Care plus bilateral Erector Spinae Plane Block (no local wound infiltration)

Interrupts pain signal transmission from the area of surgery to the Central Nervous System with local anaesthetic nerve blockade. Regional "Field" Block with local anaesthetic, given by Anaesthetist having completed ESPB training.

How: Ultrasound-guided injection of local anaesthetic, to produce a bilateral "Fascial Plane" block. Given in Operating Theatre at time of surgery, prior to emergence from anaesthesia.

Dose: 2 x 20 mls Levo-Bupivacaine 0.25% (2.5 mg/ml), 40 mls in total. 0.25% Levo-Bupivacaine is widely available "off the shelf" in all operating theatres. The volume and dose chosen will be effective in the majority of patients who will be substantially heavier than 50kg. Adjusting the volume down, rather than altering concentration, in the small number of patients below 50kg, likewise reduces the chance of drug error or local anaesthetic toxicity. The concentration and volume of Levo-Bupivacaine for local infiltration in Usual Care was chosen to emulate routine

clinical practice and deliver the same dose of drug as local infiltration (20 ml 0.5% = 100 mg).
Modifications: If patient weight < 50kg: Reduced Volume of Levo-Bupivacaine to maximum dose 2 mg/kg (e.g. 45 kg patient would receive 90 mg Levo-Bupivacaine or 36 mls as 2 x 18 mls injections) to avoid the potential for local anaesthetic toxicity. Paracetamol as per local standard of care: Omit NSAID if contraindication.

Accountability: Drug delivered to correct "Plane" by direct, real time ultrasound visualisation.
Confirmation of drug delivery by ultrasound "screen capture" after bilateral injection.

Erector Spinae Block group will receive Intervention described above plus Usual care, without Local Wound Infiltration.

Participants will be followed up for pain and other measures post-surgery whilst in hospital (in recovery, 6 hours post-surgery, 24 hours post-surgery and at discharge/72 hours post-surgery). Participants will also be followed up again 6-8 weeks post-surgery at their standard of care follow up visit.

Intervention Type

Drug

Phase

Phase III

Drug/device/biological/vaccine name(s)

Morphine, diamorphine, levobupivacaine

Primary outcome(s)

Back pain on moving around the bed (sitting up and/or turning) from Visual Analog Scale (VAS) recorded at 24 hours after surgery, reported using a 10 cm line, 0-100 score. Patients unable to sit up and/or turn as a result of pain will be assigned the highest pain score (100).

Key secondary outcome(s)

Patient-reported outcomes

1. Pain scores (VAS) 13 back & leg pain, at rest and movement, on a 0-100 scale;
2. EQ-5D-5L14: Health status questionnaire used to derive quality adjusted life year (QALYs) and used in the cost-effectiveness analysis;
3. Quality of Recovery Questionnaire (QoR-15) 15 to measure the quality of recovery after surgery and analgesia;
4. Healthcare Resource Use: measured using a bespoke questionnaire.
5. Oswestry Disability Index¹⁶ to measure functional recovery after surgery

Clinical

1. Cumulative opioid consumption after surgery, at 24 hours and 72 hours (or discharge) (converted to oral morphine equivalent)
2. Adverse events
3. Intervention-related side-effects (including: Postoperative Nausea and Vomiting (PONV), pruritus, respiratory depression (respiratory rate <8 breaths/minute)
4. Further clinical intervention such as: antiemetic administration, urinary catheterisation, High Dependency Care/Intensive Care admission
5. Time to mobilisation after surgery
6. Length of hospital stay (readiness for discharge)
7. Readmissions between discharge and routine 6-8 week postoperative follow up.

Completion date

30/06/2026

Eligibility

Key inclusion criteria

Current inclusion criteria as of 26/02/2025:

1. People aged 16 years or over.
2. Scheduled for elective, posterior lumbar-instrumented spinal surgery +/- decompression (including patients undergoing this surgery following a previous lumbar discectomy or decompression).
3. Able to give informed consent, with interpreters provided where necessary.

Previous inclusion criteria:

1. People aged 16 years or over
2. Scheduled for elective, posterior lumbar-instrumented spinal surgery +/- decompression
3. Able to give informed consent, with interpreters provided where necessary

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

16 years

Upper age limit

75 years

Sex

All

Total final enrolment

37

Key exclusion criteria

Current exclusion criteria as of 26/02/2025:

1. Patients with drug sensitivity or allergy to any of the trial agents i.e. intrathecal opioid or local anaesthetic.
2. Patients undergoing fusion at more than three vertebral levels.
3. Patients with an infection or tumour at the block site or surgical site.

4. Patients meeting criteria for American Society of Anaesthesiologists Physical Status Classification Grade 4-5.
5. Patients undergoing surgery during an emergency admission (this would preclude a detailed risk-benefit conversation with a consultant anaesthetist, which our PPI group told us was vital pre-consent).
6. Patients scheduled for single-level microdiscectomy and decompression only.
7. Patients undergoing anterior surgery.
8. Patients who have previously had posterior lumbar instrumentation. Current pregnancy: a pregnancy test, in the female patients of childbearing age is routine immediately prior to surgery.

Previous exclusion criteria:

1. Patients with drug sensitivity or allergy to any of the trial agents i.e. intrathecal opioid or local anaesthetic
2. Patients undergoing fusion at more than three vertebral levels
3. Patients with an infection or tumour at the block site or surgical site
4. Patients meeting criteria for American Society of Anaesthesiologists Physical Status Classification Grade 4-5
5. Patients undergoing surgery during an emergency admission (this would preclude a detailed risk-benefit conversation with a consultant anaesthetist, which our PPI group told us was vital pre-consent).
6. Patients scheduled for single-level microdiscectomy and decompression.
7. Current pregnancy

Date of first enrolment

30/09/2024

Date of final enrolment

26/09/2025

Locations

Countries of recruitment

United Kingdom

England

Scotland

Wales

Study participating centre

Northern General Hospital

Northern General Hospital NHS Trust

Herries Road

Sheffield

England
S5 7AU

Study participating centre
Queens Medical Centre
Nottingham University Hospital
Derby Road
Nottingham
England
NG7 2UH

Study participating centre
Ninewells Hospital
Ninewells Avenue
Dundee
Scotland
DD1 9SY

Study participating centre
Queen Elizabeth University Hospital
1345 Govan Road
Glasgow
Scotland
G51 4TF

Study participating centre
The Walton Centre NHS Foundation Trust
Lower Lane
Fazakerley
Liverpool
England
L9 7LJ

Study participating centre
Royal Preston Hospital
Sharoe Green Lane North
Fulwood
Preston
England
PR2 9HT

Study participating centre

Northumbria Healthcare NHS Foundation Trust

North Tyneside General Hospital
Rake Lane
North Shields
England
NE29 8NH

Study participating centre

Hull University Teaching Hospitals NHS Trust

Hull Royal Infirmary
Anlaby Road
Hull
England
HU3 2JZ

Study participating centre

South Tyneside and Sunderland NHS Foundation Trust

Sunderland Royal Hospital
Kayll Road
Sunderland
England
SR4 7TP

Study participating centre

Royal National Orthopaedic Hospital

Brockley Hill
Stanmore
England
HA7 4LP

Study participating centre

University Hospitals of Derby and Burton NHS Foundation Trust

Royal Derby Hospital
Uttoxeter Road
Derby
England
DE22 3NE

Study participating centre

University Hospitals Coventry and Warwickshire NHS Trust

Walsgrave General Hospital

Clifford Bridge Road

Coventry

England

CV2 2DX

Study participating centre

King's College Hospital

Denmark Hill

London

England

SE5 9RS

Study participating centre

Leeds Teaching Hospitals NHS Trust

St. James's University Hospital

Beckett Street

Leeds

England

LS9 7TF

Study participating centre

Whittington Health NHS Trust

The Whittington Hospital

Magdala Avenue

London

England

N19 5NF

Study participating centre

The Robert Jones and Agnes Hunt Orthopaedic Hospital NHS Foundation Trust

Gobowen

Oswestry

England

SY10 7AG

Sponsor information

Organisation

Sheffield Teaching Hospitals NHS Foundation Trust

ROR

<https://ror.org/018hjpz25>

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Anonymised datasets can be made available to other researchers upon request and appropriate ethical review. Participants are also asked to consent to future research (using their anonymised data) and that their anonymised datasets will be made publically available at the end of the study.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol file	version 3.1	12/02/2025	26/02/2025	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes