ISRCTN38773821 https://doi.org/10.1186/ISRCTN38773821

Excimer laser versus clobetasone propionate in prurigo form of atopic dermatitis

Submission date	Recruitment status
28/12/2006	No longer recruiting
Registration date	**Overall study status**
28/12/2006	Completed
Last Edited	**Condition category**
14/01/2021	Skin and Connective Tissue Diseases

[_] Prospectively registered
[_] Protocol
[_] Statistical analysis plan
[X] Results
[_] Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr E E A Brenninkmeijer

Contact details
Academic Medical Center Amsterdam
Department of Dermatology, A0-223
P.O. Box 22660
Amsterdam
Netherlands
1105 AZ
+31(0)20 566 2581
E.E.Brenninkmeijer@amc.uva.nl

Additional identifiers

Protocol serial number
NL785, NTR797

Study information

Scientific Title
Excimer laser versus clobetasone propionate in prurigo form of atopic dermatitis

Study objectives
As Narrow Band UltraViolet B (NB-UVB) is known to be effective in Atopic Dermatitis (AD), the excimer laser appears to be a promising treatment for localised AD. We designed a randomised single blind within-patient controlled trial to investigate the efficacy of the excimer laser versus routine topical corticosteroid, clobetason propionate.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Approval recieved from the Medical Ethical Committee (Medisch Ethische Commissie) on the 19th October 2006 (ref: MEC 06/239).

Primary study design
Interventional

Study design
Randomised, controlled, parallel group, single blind, trial

Study type(s)
Treatment

Health condition(s) or problem(s) studied
Atopic dermatitis

Interventions
All patients will be randomised to a within-patient left-right comparison study of excimer laser versus topical clobetason propionate. Treatment with the ecximer laser will be performed twice a week, during a treatment period of ten weeks. Clobetason propionate will be applied by the patients themselves once a day , according standardised instructions, during a treartment period of ten weeks.

Intervention Type
Drug

Phase
Not Specified

Drug/device/biological/vaccine name(s)
Clobetasol propionate

Primary outcome(s)
Clinical responses will be evaluated using physician assessment of individual signs (number of nodules, elevation of nodules, excoriation, erythema and pruritus).

Key secondary outcome(s)
1. Photographic documentation
2. Physician Global Assessment (PGA)

3. Patient Global Assessment (PaGA)
4. Besides the clinical responses, the patient and physician satisfaction/preference and duration of remission will be evaluated.

Completion date
01/08/2007

Eligibility

Key inclusion criteria
1. Adult patients (over 18 years old)
2. Prurigo form of atopic dermatitisbased on:
a. Hanifin and Rajka criteria fullfilled
b. presence of allergen specific Immunoglobulin E (IgE)
c. lasting for at least six months
d. refractory to the standard therapy
e. at least four symmetrical nodules
3. Upper or lower extremities affected
4. Written informed consent provided

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group
Adult

Lower age limit
18 Years

Sex
Not Specified

Total final enrolment
13

Key exclusion criteria
1. Patients unable to comply with the requirements of the study
2. Female patients who are pregnant or breastfeeding
3. Patients treated with sedating antihistamines within 24 hours before start of study treatment
4. Patients treated with topical steroids within one week before start of study treatment
5. Patients treated with phototherapy or Psoralen UltraViolet A (PUVA) therapy within one week before start of study treatment
6. Patients treated with systemic therapy that might have an effect on the prurigo form of AD within four weeks before start of study treatment
7. Patients with hypersensitivity to the study treatment or sunlight
8. Patients receiving drugs known to cause photosensitivity and/or photo toxicity
9. Patients with any other interfering skin diseases, which jeopardize the study

Date of first enrolment
01/11/2006

Date of final enrolment
01/08/2007

Locations

Countries of recruitment
Netherlands

Study participating centre
Academic Medical Center Amsterdam
Amsterdam
Netherlands
1105 AZ

Sponsor information

Organisation
Academic Medical Center (AMC) (The Netherlands)

ROR
https://ror.org/03t4gr691

Funder(s)

Funder type
Hospital/treatment centre

Funder Name
Academic Medical Center (AMC) (The Netherlands)

Alternative Name(s)
Academic Medical Center, Centre Médical Académique, ACADEMISCH MEDISCH CENTRUM AMSTERDAM, Academic Medical Center (Amsterdam), AMC

Funding Body Type
Government organisation

Funding Body Subtype
Universities (academic only)

Location
Netherlands

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2010	14/01/2021	Yes	No