

# Management of metformin-induced gastrointestinal problems

<b>Submission date</b> 22/06/2018	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 10/07/2018	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 16/08/2022	<b>Condition category</b> Other	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Metformin is a medicine used to treat type 2 diabetes that causes gastrointestinal (digestive) problems. The aim of this study is to find out whether the adverse effects of metformin can be treated with proton pump inhibitor drugs such as omeprazole and pantoprazole.

### Who can participate?

Patients aged 26-85 with type 2 diabetes who are using metformin

### What does the study involve?

This study consists of two phases. In the first phase patients with type 2 diabetes using metformin of different doses (500 mg, 850 mg and 100 mg) are randomly selected. Patients with symptoms of gastrointestinal problems are noted, along with dose, age, gender, frequency and way of taking metformin, all collected from the patient and their records. In the second phase of the study patients with gastrointestinal problems are divided into three groups. The first group receive omeprazole 40 mg, the second group receive pantoperazole 40 mg, and the third group receive non-drug treatment (i.e., take metformin during meal) for one month. Gastrointestinal symptoms are recorded before and after the treatment by interviewing the participants. Their blood sugar levels are also noted.

### What are the possible benefits and risks of participating?

The benefit to patients is that their tolerance to metformin could be increased and their quality of life could be better. There are no possible risks to the patients.

### Where is the study run from?

1. Allied Hospital
2. Faisalabad Diabetic Center
3. Diabetic Institute Pakistan

### When is the study starting and how long is it expected to run for?

May 2017 to December 2017

Who is funding the study?  
Investigator initiated and funded

Who is the main contact?  
1. Miss Madeeha Fatima  
2. Miss Saleha Sadeeqa

## Contact information

**Type(s)**  
Public

**Contact name**  
Miss Madeeha Fatima

**Contact details**  
Lahore College for Women University  
Jail Road  
Lahore  
Pakistan  
54000

**Type(s)**  
Public

**Contact name**  
Miss Saleha Sadeeqa

**Contact details**  
Lahore College for Women University  
Jail Road  
Lahore  
Pakistan  
54000

## Additional identifiers

**Protocol serial number**  
49165218

## Study information

**Scientific Title**  
Management of metformin-induced gastrointestinal problems by pharmacological and non-pharmacological treatment

**Study objectives**

Metformin is a biguanide that causes gastrointestinal problems. To manage the adverse effects of metformin proton pump inhibitors were used to investigate the outcome on the adverse effects.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

1. Board of Studies (BOS) and Advance Study Research Board (ASRB), Lahore College for Women University (LCWU), 26/05/2017
2. Hospital ethics committee Punjab Medical College Faisalabad, 19/07/2017, ref: PMC/PHRC /ERC/2017/11

### **Primary study design**

Interventional

### **Study design**

The study consisted of two parts: in the first phase a cross-sectional research design was used and in the second phase an experimental research (interventional) design was used

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Metformin gastrointestinal adverse effects

### **Interventions**

The first phase is a quantitative study in which a cross sectional research design was used. In this study patients having type 2 diabetes using metformin of different doses (500 mg, 850 mg and 100 mg) were considered. Patients having symptoms of metformin induced gastrointestinal problems were noted. The gastrointestinal symptoms induced by metformin were linked with dose, age, gender, frequency and way of intake of metformin. Data was collected from the patients by visiting the outpatient departments of different government and private hospitals.

In the second phase of the study an experimental design was used. In this phase patients having metformin induced gastrointestinal problems were divided into three groups (convenience sampling was used). Each group consisted of 20 participants. The participants were subjected to PPIs in order to see the effectiveness of proton pump inhibitors such as omeprazole and pantoprazole. The first group of participants were subjected to omeprazole 40 mg. Similarly the second group was subjected to pantoperazole 40 mg. In this phase the initial symptoms of the patient were recorded and then after intervention the level and severity of the symptoms was again checked by directly interviewing the participants on their follow-up and recorded. Their blood sugar levels were also noted. In this way the effectiveness of the medicine is checked. The third group of the patients were subjected to non-pharmacological treatment i.e. take metformin during meals and the outcomes are recorded. The total duration of treatment with PPIs of each patient was one month and after one month patients were followed up.

### **Intervention Type**

Drug

**Phase**

Not Applicable

**Drug/device/biological/vaccine name(s)**

Omeprazole, pantoperazole, metformin

**Primary outcome(s)**

GI symptoms measured by interview at baseline and 1 month

**Key secondary outcome(s)**

Blood sugar level measured at baseline and 1 month

**Completion date**

30/12/2017

**Eligibility****Key inclusion criteria**

1. Male and female aged 26-85 years
2. Patients diagnosed with Type 2 diabetes
3. Patients using metformin

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Key exclusion criteria**

1. Type 1 diabetes patients
2. Patients under 26 and older than 85 years
3. Patients who were not taking metformin
4. Patients on insulin therapy alone

**Date of first enrolment**

01/06/2017

**Date of final enrolment**

30/11/2017

**Locations****Countries of recruitment**

Pakistan

**Study participating centre**

**Allied Hospital**

Faisalabad

Pakistan

-

**Study participating centre**

**Faisalabad Diabetic Center**

Faisalabad

Pakistan

-

**Study participating centre**

**Diabetic Institute Pakistan**

Lahore

Pakistan

-

## **Sponsor information**

**Organisation**

Lahore College for Women University

**ROR**

<https://ror.org/02bf6br77>

## **Funder(s)**

**Funder type**

Other

**Funder Name**

Investigator initiated and funded

# Results and Publications

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request after the publication of the results of the study upon request.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>			16/08/2022	Yes	No