

The effectiveness of screening for postpartum depression in child health care

Submission date 28/10/2016	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 09/11/2016	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 09/11/2016	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Although a depression in the period after giving birth (a postpartum depression) is not a rare condition, many cases go unnoticed. Many women do not recognize the symptoms of postpartum depression (PPD) and fail to seek help and health care professionals who see the mother often do not spot the signs either. When PPD is not identified, this can have a serious impact on the functioning of the mother, the mother-child relationship, the mother's ability to care for their baby and on the family as a whole. One way to better identify PPD in an early stage is by screening women for signs of depression using a questionnaire. There are well developed screening instruments for PPD, including the Edinburg Postnatal Depression Scale (EPDS). In the Netherlands almost all mothers (around 95%) of newborn children visit the Child Health Care Centre (CHC) routinely during the first year after giving birth. Incorporating PPD screening into these visits could therefore provide a unique opportunity to improve the detection and treatment of PPD. The aim of this study is to find out whether screening by CHC with the EPDS is a good way of identifying women with PPD and speeding up the recovery of those with PPD.

Who can participate?

All mothers of children born between the first of December 2012 and the 31st of March 2014, using the offer of Child Health Care in the regions of the participating organizations.

What does the study involve?

Mothers in the screening region are asked to complete the a questionnaire about depression before their planned visit to the CHC center at one, three and six months. The physician discusses the outcome with the mother during the visit. Depending on the score, the mother is referred for further help or treatment or an extra home visit by the CHC nurse. In the control region the mothers and their infants also visit the CHC at one, three and six months, but without completing a depression questionnaire. The physician may ask the mother about depressive symptoms and give advice for further help, but this is not a required part of the consultation. The number of mothers suffering from depression after nine months is then measured.

What are the possible benefits and risks of participating?

There are no direct benefits or risks involved with participating in this study.

Where is the study run from?

Mothers in the screening region attend CHC centres in the region of Twente (GGD Twente), mothers for the control region attend CHC centres in the regions of Apeldoorn (Vérian) and Deventer (GGD IJsselland) (Netherlands)

When is the study starting and how long is it expected to run for?

October 2011 to March 2014

Who is funding the study?

Netherlands Organization for Health Research and Development (ZonMw) (Netherlands)

Who is the main contact?

Dr Angarath van der Zee – van den Berg

a.i.vandenberg@utwente.nl

Dr Magda Boere-Boonekamp

m.m.boere-boonekamp@utwente.nl

Contact information

Type(s)

Public

Contact name

Dr Magda Boere - Boonekamp

ORCID ID

<https://orcid.org/0000-0001-9707-1056>

Contact details

Institute for Innovation and Governance Studies

Dept. Health Technology & Services Research

Ravelijn Building room RA 5254 | PO Box 217

Enschede

Netherlands

7500 AE

Type(s)

Scientific

Contact name

Mrs Angarath van der Zee - van den Berg

ORCID ID

<https://orcid.org/0000-0001-9673-6677>

Contact details

Institute for Innovation and Governance Studies

Dep. Health Technology & Services Research

Ravelijn Building room RA 5260 | PO Box 217

Enschede
Netherlands
7500AE

Additional identifiers

Protocol serial number

AWJT project 15901.0003 (ZonMw)

Study information

Scientific Title

The Effectiveness of Screening for Postpartum Depression in Child Health Care: does Screening Improve Maternal Mental Health and the Child's Socioemotional Development when Compared to Care as Usual?

Acronym

Post-Up study

Study objectives

Screening for postpartum depression (PPD) by Child Health Care improves the early detection and results in a more rapid mental recovery of the mothers identified with PPD, better quality of parenting, and less problems in social-emotional development of the children of these mothers.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The protocol for the study was submitted to the Medical Ethics Committee Twente (MECT) for adjudication. The METC exempted the study from review as the nature of the study was found beyond the remit of the Medical Sciences Research with Human Subjects Act (Wet medisch-wetenschappelijk onderzoek met mensen) and the measures pertaining to confidentiality and informed consent were found appropriate.

Study design

Prospective non-randomised study

Primary study design

Interventional

Study type(s)

Screening

Health condition(s) or problem(s) studied

Postpartum depression

Interventions

Participating mothers will be asked to fill in the 10-item EPDS questionnaire prior to their planned visit to the Child Health Care (CHC) center at 1, 3 and 6 months. The CHC physician scores the EPDS during the visit and discusses the outcome with the mother. A score ≥ 13 is

interpreted as a high risk to be diagnosed with major depression. If the clinical impression of the WBC physician corresponds to the score, the mother will be advised to consult a family practitioner or mental health professional. In the case of EPDS scores from 9 to 12, the mother is offered a home visit by the CHC nurse. In all cases, the decision on the follow up step after screening is in dialogue with the mother.

In the control region the mothers and their infants visit the CHC at 1, 3 and 6 months. At these visits, the physician has the usual conversation with the mother and can ask about their psychological well-being if they want (this is not a standardized part of the consultation). If postpartum depression is suspected, the mother can be offered a home visit or referred for further help as per standard practice.

Intervention Type

Other

Primary outcome(s)

Presence of depression (major and minor) at 9 months postpartum, instrument: Dutch translation of the depression subscale of the Mini International Neuropsychiatric Interview at 9 months postpartum.

Key secondary outcome(s)

Mother:

1. Health-related quality of life is measured using the Short-Form 12-Item Health Survey (SF-12) at 12 months postpartum
2. Maternal anxiety is measured using the 6-item short form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI) at 12 months postpartum
3. Quality of parenting is measured using the Dutch translation of the Maternal Self-Efficacy in the nurturing role questionnaire (SENR) at 12 months postpartum

Child:

Socioemotional development is measured using Ages and Stages Questionnaires: Social Emotional (ASQ:SE), 12 months version, at 12 months postpartum.

Cost-effectiveness:

1. Health care use of the mother is measured using self-report questions at 12 months postpartum
2. Health care use of the infant is measured using self-report questions at 12 months postpartum
3. Proportion of mothers who receive treatment for postpartum depression is measured using self-report questions at 12 months postpartum
4. Sick leave of the mother is measured using self-report questions at 12 months postpartum

Completion date

30/11/2016

Eligibility

Key inclusion criteria

1. All mothers of children born between the first of December 2012 and the 31st of March 2014
2. Visiting Child Health Care (CHC) centers in the eastern part of the Netherlands

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

1. Insufficient mastery of the Dutch language to fill out questionnaires
2. Not utilizing the offer of Well Baby Care
3. First contact with Well Baby Care after the child was 3 months of age

Date of first enrolment

01/12/2012

Date of final enrolment

31/03/2014

Locations

Countries of recruitment

Netherlands

Study participating centre

GGD Regio Twente

Nijverheidstraat 30

Enschede

Netherlands

7511JM

Study participating centre

Vérian

Postbus 1032

Apeldoorn

Netherlands

7301 BG

Study participating centre

GGD IJsselland

Schurenstraat 8a

Deventer

Netherlands
7413 RA

Sponsor information

Organisation

University of Twente - Institute for Innovation and Governance Studies

ROR

<https://ror.org/006hf6230>

Funder(s)

Funder type

Not defined

Funder Name

Netherlands Organization for Health Research and Development (ZonMw)

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Study website	Study website	11/11/2025	11/11/2025	No	Yes