

Use of a nitric oxide (ISMN) for the PREVENTION and MANAGEMENT of pre-eclampsia (pilot study)

Submission date 29/11/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 21/12/2007	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 03/03/2009	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Graeme Smith

Contact details
Clinical Research Centre
Queen's University
Kingston General Hospital
76 Stuart Street, Angada 4, Room 5-415
Kingston
Canada
K7L 2V7
+1 613 549 6666 ext. 3936
gns@post.queensu.ca

Additional identifiers

Protocol serial number
FMI-63194

Study information

Scientific Title

A pilot study to evaluate glyceryl trinitrate (GTN) as a novel therapeutic for the prevention and treatment of pre-eclampsia

Study objectives

CIHR Grant Submission Title: Pre-eclampsia: Fetal and Maternal Outcomes and Innovative Therapies

To determine if exogenous glyceryl trinitrate (GTN), compared to placebo, will be effective at preventing the development and/or progression of clinical pre-eclampsia.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Queen's University Health Sciences and Affiliated Teaching Hospitals Research Ethics Board, Kingston, Ontario (Canada) approved on the 20th September 2002 (ref: ANAT-017-02)

Primary study design

Interventional

Study design

Randomised, multicentre, blinded, placebo-controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Pre-eclampsia

Interventions

The study is a randomised blinded drug/placebo trial. The randomisation scheme was prepared by an independent statistician prior to initiation of the pilot study, and prepared in blocks for if and when Ottawa Hospital comes on board. The study investigators, associated staff, outcome assessor, data analyst and the study participants will all be blinded to the treatment allocation. ISMN and placebo capsules will be prepared to have identical shape, size, color, smell and feel. No form of identification labeling will be visible on either intervention. When a suitable participant is identified, the research nurse coordinator will explain the details and potential risks and benefits of the study. If consent is granted, the research nurse coordinator will determine the treatment assignment for that subject by calling the research pharmacist who will provide the next code indicating the treatment for a given patient. Patients will then be provided with an appropriately labeled package of pills prepared by the hospital pharmacy.

Prevention Arm:

Experimental intervention: Daily dose of low dose Isosorbide-5-mononitrate (ISMN) (30 mg) beginning after 20 weeks gestation till delivery.

Control intervention: Matching placebo containing lactose. Patients randomly assigned to either receive low dose ISMN (30 mg) as stated above or placebo.

Management Arm:

Experimental intervention: Daily dose of low dose Isosorbide-5-mononitrate (ISMN) (30 mg)

following diagnosis of pre-eclampsia after 24 weeks gestation till delivery.
Control intervention: Matching placebo containing lactose. Patients randomly assigned to either receive low dose ISMN (30 mg) as stated above or placebo.

In both arms of the study, patients will receive standard clinical care. Total duration of treatment in each arm is flexible and based on each individual participant. There is no follow-up after delivery.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Glyceryl trinitrate

Primary outcome(s)

Prevention arm: incidence of pre-eclampsia in the ISMN/placebo groups, measured at delivery
Treatment arm: randomisation-to-delivery interval between ISMN/placebo groups, measured at delivery

Key secondary outcome(s)

1. Serial change in biochemical markers in treatment/no treatment groups in each of the studies, measured at routine obstetrical visits until delivery (generally every 2 weeks)
2. Incidence of any side effects (major or minor), measured at routine obstetrical visits until delivery (generally every 2 weeks)
3. Neonatal outcomes (composite of neonatal morbidity), measured at delivery

Completion date

31/12/2009

Eligibility

Key inclusion criteria

Women of childbearing years (approximately 18 - 42 years).

Prevention arm:

All women with a past obstetrical history of one or more cases of severe early onset pre-eclampsia or later onset severe pre-eclampsia associated with haemolysis, elevated liver enzymes, low blood levels of platelets (HELLP) syndrome.

Treatment arm:

All women that have been diagnosed with pre-eclampsia that are being followed clinically and that provide informed consent. For a diagnosis of pre-eclampsia a patient must meet all three criteria:

1. Systolic blood pressure greater than 140 mmHg or an increase of 30 mmHg from the participants baseline (with that increase present at two measurements taken 6 hours apart)
2. Diastolic blood pressure greater than 90 mmHg or an increase of 15 mmHg from the participants baseline (with that increase present at two measurements taken 6 hours apart)
3. Proteinuria greater than 0.3 g in 24 hour urine or 2+ on dipstick

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

Female

Key exclusion criteria

Potential women excluded are those:

1. That have a contraindication to use of isosorbide mononitrate (ISMN)
2. That have either a maternal or foetal indication for delivery
3. That have a diagnosis of severe pre-eclampsia (diastolic blood pressure greater than 100 mmHg; proteinuria greater than 1 g/d), eclampsia, or HELLP syndrome at time of recruitment

Date of first enrolment

01/01/2007

Date of final enrolment

31/12/2009

Locations**Countries of recruitment**

Canada

Study participating centre**Clinical Research Centre**

Kingston

Canada

K7L 2V7

Sponsor information**Organisation**

Queen's University (Canada)

ROR

<https://ror.org/02y72wh86>

Funder(s)

Funder type

Research organisation

Funder Name

Canadian Institutes of Health Research (CIHR) (Canada) - <http://www.cihr-irsc.gc.ca> (ref: FMI-63194)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration