

# Intravesical Ialuril® plus oral combination of curcumine, quercetine, hyluronic acid (HA) and chondroitin sulphate (CS) for prophylaxis of recurrent non-muscle invasive transitional cell carcinoma (TCC) of the bladder

<b>Submission date</b> 10/03/2013	<b>Recruitment status</b> Stopped	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 20/03/2013	<b>Overall study status</b> Stopped	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 21/01/2019	<b>Condition category</b> Cancer	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Transitional cell carcinoma (TCC) of the bladder is one of the most common cancers. In 75-85% of patients the cancer is contained inside the lining of the bladder (non-muscle-invasive bladder cancer). A surgical technique called complete transurethral resection of visible tumour (TURBT) is the gold standard treatment for non-invasive bladder tumours. A single dose of chemotherapy, usually mitomycin C, is delivered directly into the bladder using a catheter (intravesical therapy) to destroy any remaining microscopic tumour and prevent the cancer from coming back. Intravesical therapy is also used in a maintenance (long term) fashion to prolong the beneficial effects of the chemotherapy. Although immediate intravesical mitomycin C or epirubicin is highly recommended, it should not be used in patients with a bladder perforation as a small number of serious complications related to mitomycin C extravasation (leakage) have been reported. The bladder lining (urothelium) prevents undesirable substances penetrating the deeper layers of the bladder wall and neutralises toxic substances. Glycosaminoglycans (GAGs) make the inner bladder wall impervious to the contents of the urine. Two key components of the layer are hyaluronic acid (HA) and chondroitin sulphate (CS). The aim of this study is to find out whether intravesical HA and CS (Ialuril®) followed by oral HA and CS plus curcumine and quercetine are safe and effective at preventing the recurrence of TCC after TURBT.

### Who can participate?

Patients aged over 18 with TCC

### What does the study involve?

Patients receive intravesical HA and CS (Ialuril®) immediately after TURBT, followed by oral treatment with pills (one a day for 90 days) containing CS, HA, quercetine and curcumine. TCC recurrence is assessed using ultrasound, urine tests and endoscopic assessment after 3 months.

What are the possible benefits and risks of participating?  
Not provided at time of registration

Where is the study run from?  
Institut Biochimique SA (IBSA) (Italy)

When is the study starting and how long is it expected to run for?  
January 2012 to December 2013

Who is funding the study?  
Institut Biochimique SA (IBSA) (Italy)

Who is the main contact?  
Dr Massimo Lazzeri

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Massimo Lazzeri

**Contact details**  
via Stamira d'Ancona 20  
Milan  
Italy  
20127

## Additional identifiers

**Protocol serial number**  
001/laluril\_Oral

## Study information

**Scientific Title**  
Intravesical laluril® plus oral combination of curcumine, quercetine, hyaluronic acid (HA) and chondroitin sulphate (CS) for prophylaxis of recurrent non-muscle invasive transitional cell carcinoma (TCC) of the bladder: a prospective longitudinal investigative pilot non-randomised study

**Study objectives**  
To test the hypothesis whether or not immediate post-TURBT intravesical instillation of a combination of HA and CS (laluril®) repairing coating - followed by oral administration of a combination of HA and CS plus curcumine and quercetine maintaining coating - is safe, tolerated and effective in preventing the recurrence of TCC after TURBT in patients with superficial low-intermediate risk pTa-1 TCC.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Not provided at time of registration

**Study design**

Prospective longitudinal investigative pilot non-randomised (phase 1b) study

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Superficial (non muscle invasive) transitional cell carcinoma

**Interventions**

All the patients underwent standard transurethral resection of the visible lesions. Immediate post-TURBT intravesical instillation of a combination of HA and CS (laluril®: 50 ml/vial solution containing HA 800 mg and CS 1 mg) followed by oral treatment with pills (one a day for 90 days) containing CS 200 mg, HA 20 mg, quercitine 200 mg and curcumine 100 mg.

**Intervention Type**

Drug

**Phase**

Phase I

**Drug/device/biological/vaccine name(s)**

Laluril®

**Primary outcome(s)**

1. Tolerability, reported as discontinuation or deviation of the protocol and safety as general or local side effects
2. Efficacy, defined as the absence of recurrence as defined by negative ultrasound evaluation, negative urine cytology and negative endoscopic assessment after 3 months

**Key secondary outcome(s)**

Morphological immunohistochemical assessment, comparing samples at entry and after 3 months for transient receptor protein vanilloids 1 (TRPV1) expression and GAGs

**Completion date**

31/12/2013

**Reason abandoned (if study stopped)**

Local ethical committees requested several changes

**Eligibility**

## **Key inclusion criteria**

Patients (male and female) with single, ultrasound detected < 3 cm tumor diameter, or multiple 3 or less, each < 1 cm, low- and intermediate-grade (Grade II in old nomenclature) superficial TCC at the first diagnosis, confirmed by flexible cystoscopy and cold cup biopsy

## **Participant type(s)**

Patient

## **Healthy volunteers allowed**

No

## **Age group**

Adult

## **Sex**

All

## **Key exclusion criteria**

1. Concomitant CIS
2. Involvement of the prostatic urethra
3. Neurogenic bladder
4. Diabetes mellitus type I and II
5. Previous bladder or prostate surgery (any kind)
6. Previous treatment with drugs (capsaicin, vanilloids, ovanil, botulinum toxin etc)
7. Further exclusion criteria were previous chemotherapies and/or pelvic radiotherapy

## **Date of first enrolment**

01/01/2012

## **Date of final enrolment**

31/12/2013

## **Locations**

### **Countries of recruitment**

Italy

### **Study participating centre**

via Stamira d'Ancona 20

Milan

Italy

20127

## **Sponsor information**

**Organisation**

Institut Biochimique SA (IBSA) (Italy)

**ROR**

<https://ror.org/02cf8gj49>

**Funder(s)****Funder type**

Industry

**Funder Name**

Institut Biochimique SA (IBSA) (Italy)

**Results and Publications****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration