ISRCTN45979711 https://doi.org/10.1186/ISRCTN45979711

REVascularisation for Ischaemic VEntricular Dysfunction

Submission date	Recruitment status	[X] Prospectively registered
20/11/2012	No longer recruiting	[X] Protocol
Registration date	**Overall study status**	[X] Statistical analysis plan
20/11/2012	Completed	[X] Results
Last Edited	**Condition category**	[_] Individual participant data
26/11/2026	Circulatory System	

Plain English summary of protocol

Current plain English summary as of 14/09/2018:
Background and study aims
In 2002, it was estimated that approximately 900,000 individuals in the United Kingdom had a diagnosis of heart failure and at least 1 in 20 of all deaths here were related to this condition. There is evidence of an increase in heart failure in the population, with the number of associated hospital admissions expected to increase by around 50% in the next 25 years. This is the likely consequence of a progressively aging population and improved survival from acute coronary syndromes, partly due to more efficient and timely revascularisation techniques. Patients with heart failure are traditionally treated with a combination of tablets and (in some cases) by insertion of a special pacemaker. Together these treatments are called Optimal Medical Therapy (OMT). In patients who have heart failure as well as narrowed heart arteries, several recent studies have suggested that treatment of the narrowed arteries by Percutaneous Coronary Intervention (PCI) or Coronary Artery Bypass Grafting (CABG) may improve heart muscle pumping strength and heart failure symptoms. However, most of these studies have been too small or have not been scientific enough to allow widespread use of PCI or CABG as a treatment for heart failure. The purpose of this study is to assess whether treatment of heart arteries by angioplasty and stenting (PCI) in combination with OMT can improve heart muscle function, quality of life and life expectancy of patients, compared to OMT alone.

Who can participate?
Patients at least 18 years of age with poor heart pumping function and diseased arteries of the heart.

What does the study involve?
Patients will be randomly allocated to two treatment groups - either Percutaneous Coronary Intervention (PCI) and Optimal Medical Therapy (OMT), or to OMT alone.

What are the possible benefits and risks of participating?
As the benefit of treating narrowed arteries has not been clearly established yet, patients should assume that there would be no direct benefit to them. There is a very small risk of major

complications during or shortly after the PCI procedure (including damage to an artery, heart attack, stroke or death). PCI procedures involve exposure to radiation in the form of X-rays, which can potentially be harmful.

Where is the study run from?
The trial will take place at approximately 35 centres in the UK. The main centre is Guy's & St Thomas' NHS Foundation Trust in London and will be coordinated from the London School of Hygiene and Tropical Medicine Clinical Trials Unit (LSHTM CTU).

When is the study starting and how long is it expected to run for?
Recruitment began in August 2013 and will continue until the end of April 2020. Follow-up will be for a minimum of two years, and the study is expected to finish in March 2022

Who is the main contact?
Ruth Canter
ruth.canter@lshtm.ac.uk

Previous plain English summary:
Background and study aims
In 2002, it was estimated that approximately 900,000 individuals in the United Kingdom had a diagnosis of heart failure and at least 1 in 20 of all deaths here were related to this condition. There is evidence of an increase in heart failure in the population, with the number of associated hospital admissions expected to increase by around 50% in the next 25 years. This is the likely consequence of a progressively aging population and improved survival from acute coronary syndromes, partly due to more efficient and timely revascularisation techniques. Patients with heart failure are traditionally treated with a combination of tablets and (in some cases) by insertion of a special pacemaker. Together these treatments are called Optimal Medical Therapy (OMT). In patients who have heart failure as well as narrowed heart arteries, several recent studies have suggested that treatment of the narrowed arteries by Percutaneous Coronary Intervention (PCI) or Coronary Artery Bypass Grafting (CABG) may improve heart muscle pumping strength and heart failure symptoms. However, most of these studies have been too small or have not been scientific enough to allow widespread use of PCI or CABG as a treatment for heart failure. The purpose of this study is to assess whether treatment of heart arteries by angioplasty and stenting (PCI) in combination with OMT can improve heart muscle function, quality of life and life expectancy of patients, compared to OMT alone.

Who can participate?
Patients at least 18 years of age with poor heart pumping function and diseased arteries of the heart.

What does the study involve?
Patients will be randomly allocated to two treatment groups - either Percutaneous Coronary Intervention (PCI) and Optimal Medical Therapy (OMT), or to OMT alone.

What are the possible benefits and risks of participating?
As the benefit of treating narrowed arteries has not been clearly established yet, patients should assume that there would be no direct benefit to them. There is a very small risk of major complications during or shortly after the PCI procedure (including damage to an artery, heart attack, stroke or death). PCI procedures involve exposure to radiation in the form of X-rays, which can potentially be harmful.

Where is the study run from?
The trial will take place at approximately 25 centres in the UK. The main centre is Guy's & St Thomas' NHS Foundation Trust in London and will be coordinated from the clinical trial unit at London School of Hygiene and Tropical Medicine (UK).

When is study starting and how long is it expected to run for?
Recruitment will begin in May 2013 and continue until May 2016. Follow-up will be for a minimum of two years, and the study is expected to finish in May 2018.

Who is funding the study?
NIHR Health Technology Assessment - HTA (UK).

Who is the main contact?
Richard Evans
richard.evans@lshtm.ac.uk

Contact information

Type(s)
Scientific

Contact name
Dr Divaka Perera

Contact details
King's College London
The Rayne Institute
Lambeth Wing
St Thomas' Hospital
London
United Kingdom
SE1 7EH
-
Divaka.Perera@kcl.ac.uk

Type(s)
Public

Contact name
Ms Ruth Canter

ORCID ID
https://orcid.org/0000-0003-0916-2551

Contact details
Clinical Trials Unit (CTU)
Department of Medical Statistics
London School of Hygiene & Tropical Medicine
Keppel Street
London
United Kingdom

WC1E 7HT
+44 20 7927 2071
Ruth.Canter@lshtm.ac.uk

Additional identifiers

ClinicalTrials.gov (NCT)
NCT01920048

Protocol serial number
HTA 10/57/67

Study information

Scientific Title
REVascularisation for Ischaemic VEntricular Dysfunction

Acronym
REVIVED

Study objectives
Current study hypothesis as of 21/04/2022:
Compared to medical therapy alone, PCI improves event-free survival in patients with ischaemic cardiomyopathy and viable myocardium.

More details and the latest version of the Protocol can be found at: https://www.journalslibrary.nihr.ac.uk/programmes/hta/105767#/

Previous study hypothesis:
Compared to medical therapy alone, PCI improves event-free survival in patients with ischaemic cardiomyopathy and viable myocardium.

More details can be found at: http://www.nets.nihr.ac.uk/projects/hta/105767
Protocol can be found at: http://www.nets.nihr.ac.uk/__data/assets/pdf_file/0016/81124/PRO-10-57-67.pdf

Ethics approval required
Old ethics approval format

Ethics approval(s)
Westminster Research Ethics Committee, 13/09/2010, bref: 10/H0802/46

Primary study design
Interventional

Study design
Multi-centre phase III randomised double-blind controlled trial

Study type(s)
Treatment

Health condition(s) or problem(s) studied
Cardiology, heart failure

Interventions
Patients are randomised to receive either Optimal Medical Therapy (OMT) alone or Percutaneous Coronary Intervention (PCI) and OMT.

Intervention Type
Procedure/Surgery

Primary outcome(s)
Current primary outcome measure as of 23/10/2018:
A composite of all-cause death and hospitalisation due to heart failure with a minimum follow-up of 2 years and maximum follow-up of approximately 8.5 years depending on time of randomisation.

Previous primary outcome measure as of 14/09/2018:
A composite of all-cause death and hospitalisation due to heart failure

Previous primary outcome measures:
1. All-cause death
2. Acute myocardial infarction or hospitalisation due to heart failure (hierarchy: death > MI > heart failure)

Key secondary outcome(s)
Current secondary outcome measures as of 21/04/2022:
1. Left Ventricular Ejection Fraction (LVEF) on echocardiography at 6 months and 1 year
2. Quality of life score:
2.1. Kansas City Cardiomyopathy questionnaire (KCCQ) up to 2 years
2.2. EuroQol EQ-5D-5L at 6 months and then yearly to the end of follow-up
3. New York Heart Association Functional (NYHA) Class up to 2 years
4. Cardiovascular death over the entire duration of follow-up
5. All-cause death over the entire duration of follow-up
6. Hospitalisation due to heart failure over the entire duration of follow-up
7. Acute myocardial infarction (MI) over the entire duration of follow-up
8. Appropriate implantable cardioverter defibrillator (ICD) therapy to 2 years
9. Unplanned further revascularisation over the entire duration of follow-up
10. Canadian Cardiovascular Society (CCS) up to 2 years
11. NHS resource use
12. Brain natriuretic peptide (BNP or NT-Pro BNP) up to 2 years
13. Major bleeding up to 2 years

Previous secondary outcome measures:
1. Cardiovascular death, MI, CVA or unplanned revascularisation at 30-days
2. Left ventricular ejection fraction at 6 months, 1 year
3. Cardiovascular death or myocardial infarction
4. Hospitalisation for heart failure
5. Appropriate ICD therapy
6. Unplanned further revascularisation
7. Acute coronary syndrome

Completion date
31/03/2022

Eligibility

Key inclusion criteria
Current inclusion criteria as of 13/08/2014:
1. LVEF ≤35%
2. Extensive coronary disease (BCIS-1 Jeopardy Score ≥6)
3. Viability in at least 4 dysfunctional segments, that can be revascularised by PCI

Previous inclusion criteria:
1. LVEF ≤30%
2. Extensive coronary disease (BCIS-1 Jeopardy Score ≥6)
3. Viable myocardium in ≥30% of dysfunctional segments

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group
Adult

Sex
All

Total final enrolment
700

Key exclusion criteria
Current exclusion criteria as of 14/09/2018:
1. Myocardial infarction <4 weeks previously
2. Decompensated heart failure requiring inotropic support, invasive or non-invasive ventilation or IABP/left ventricular assist device (LVAD) therapy <72 hours prior to randomisation
3. Sustained VT/VF or appropriate ICD discharges <72 hours prior to randomisation
4. Valve disease requiring intervention
5. Contraindications to PCI
6. Aged <18 years
7. eGFR < 25 ml/min, unless established on dialysis
8. Women who are pregnant
9. Previously enrolled in REVIVED-BCIS2 or current enrolment in other trial that may affect REVIVED-BCIS2 outcome data
10. Life expectancy <1 year due to non-cardiac pathology

Previous exclusion criteria as of 13/08/2014:
Specific exclusions:
1. Significant angina (≥CCS class 3)
2. Myocardial infarction < 4 weeks previously

General exclusions:
1. Decompensated heart failure requiring inotropic support or IABP/LVAD therapy <72 hours prior to randomisation
2. Sustained VT/VF or appropriate ICD discharges <72 hours prior to randomisation
3. More than mild aortic stenosis or more than mild aortic regurgitation on echocardiography
4. Contra-indications to PCI
5. Age <18 years
6. eGFR < 25 ml/min, unless established on dialysis
7. Women who are pregnant
8. Previously enrolled in REVIVED or current enrolment in other study
9. Life expectancy < 1 year due to non-cardiac pathology

Previous exclusion criteria:
Specific exclusions:
1. Significant angina (≥CCS class 3)
2. Myocardial infarction < 6 weeks previously

General exclusions:
1. Decompensated heart failure requiring inotropic support or IABP/LVAD therapy <72 hours prior to randomisation
2. Sustained VT/VF or appropriate ICD discharges <72 hours prior to randomisation
3. More than mild aortic stenosis or mild aortic regurgitation on echocardiography
4. Contra-indications to PCI, including contra-indications to Aspirin or Clopidogrel or Heparin
5. Age <18 years
6. Bleeding diathesis or Warfarin therapy with INR>3
7. Active internal bleeding (except menstruation)
8. Platelet count < 100,000 cells/mm3) at randomisation
9. Haemoglobin < 9 g/dl at randomisation
10. eGFR < 25 ml/min, unless established on dialysis
11. Women who are pregnant
12. Previously enrolled in REVIVED or current enrolment in other study
13. Life expectancy < 1 year due to non-cardiac pathology

Date of first enrolment
01/08/2013

Date of final enrolment
19/03/2020

Locations

Countries of recruitment
United Kingdom

England

Northern Ireland

Scotland

Wales

Study participating centre
King's College London
-
London
England
SE5 9RS

Study participating centre
Basingstoke and North Hampshire Hospital
Aldermaston Road
Basingstoke
England
RG24 9NA

Study participating centre
Blackpool Victoria Hospital
Whinney Heys Road
Blackpool
England
FY3 8NR

Study participating centre
Derriford Hospital
Derriford Road
Derriford
Plymouth
England
PL6 8DH

Study participating centre
Dorset County Hospital
Williams Avenue
Dorchester
England
DT1 2JY

Study participating centre
Freeman Hospital
Freeman Road

High Heaton
Newcastle upon Tyne
England
NE7 7DN

Study participating centre
Glan Clwyd Hospital
North Wales Cardiac Centre
Rhyl
Wales
LL30 1LB

Study participating centre
Glenfield Hospital
Groby Road
Leicester
England
LE3 9QP

Study participating centre
Golden Jubilee National Hospital
Agamemnon Street
Clydebank
Scotland
G81 4DY

Study participating centre
Great Western Hospital
-
Swindon
England
SN3 6BB

Study participating centre
Kettering General Hospital
Rothwell Road
Kettering
England
NN16 8UZ

Study participating centre
Leeds General Infirmary
Great George Street
Leeds
England
LS1 3EX

Study participating centre
Lister Hospital
Coreys Mill Lane
Stevenage
England
SG1 4AB

Study participating centre
Liverpool Heart and Chest Hospital
-
Liverpool
England
L14 3PE

Study participating centre
Manchester Royal Infirmary
Oxford Road
Manchester
England
M13 9WL

Study participating centre
New Cross Hospital, Heart and Lung Centre
Wolverhampton Road
Heath Town
Wolverhampton
England
WV10 0QP

Study participating centre
Ninewells Hospital
Ninewells Avenue
Dundee

Scotland
DD1 9SY

Study participating centre
Pinderfields Hospital
Aberford Road
Wakefield
England
WF1 4DG

Study participating centre
Queen Alexandra Hospital
Southwick Hill Road
Cosham
Portsmouth
England
PO6 3LY

Study participating centre
Royal Bournemouth Hospital
Castle Lane East
Bournemouth
England
BH7 7DW

Study participating centre
Royal Devon and Exeter Hospital
-
Exeter
England
EX2 3DW

Study participating centre
Royal Free Hospital
Rowland Hill Street
London
England
NW3 2PF

Study participating centre
Royal Infirmary of Edinburgh at Little France
51 Little France Crescent
Old Dalkeith Road
Edinburgh
Lothian
Scotland
EH16 4SA

Study participating centre
Royal Oldham Hospital
Rochdale Road
Oldham
England
OL1 2JH

Study participating centre
The Royal Victoria Hospital
274 Grosvenor Road
Belfast
Northern Ireland
BT12 6BA

Study participating centre
Salisbury District Hospital
Odstock Road
Salisbury
England
SP2 8BJ

Study participating centre
Southampton
Southampton General Hospital
Tremona Road
Southampton
England
SO16 6YD

Study participating centre
St Bartholomew's Hospital
-

London
England
EC1A 7BE

Study participating centre
St George’s Hospital
-
London
England
SW17 0QT

Study participating centre
St Thomas' Hospital
St. Thomas's Hospital
Westminster Bridge Road
London
England
SE1 7EH

Study participating centre
Sunderland Royal Hospital
-
Sunderland
England
SR4 7TP

Study participating centre
The James Cook University Hospital
Marton Road
Middlesbrough
England
TS4 3BW

Study participating centre
Coventry
University Hospital
Clifford Bridge Road
Coventry
England
CV2 2DX

Study participating centre
Worcestershire Royal Hospital
Charles Hastings Way
Worcester
England
WR5 1DD

Study participating centre
Worthing Hospital
Lyndhurst Road
Worthing
England
BN11 2DH

Study participating centre
Wythenshawe Hospital
Southmoor Road
Wythenshawe
Manchester
England
M23 9LT

Study participating centre
Bristol Royal Infirmary
Upper Maudlin Street
Bristol
England
BS2 8HW

Study participating centre
Birmingham Heartlands Hospital
Bordesley Green East
Bordesley Green
Birmingham
England
B9 5SS

Study participating centre

York Hospital
Wigginton Road
York
England
YO31 8HE

Study participating centre
Sheffield
Northern General Hospital
Herries Road
Sheffield
England
S5 7AU

Sponsor information

Organisation
King's College London

ROR
https://ror.org/0220mzb33

Organisation
Guy’s and St Thomas’ NHS Foundation Trust

Funder(s)

Funder type
Government

Funder Name
NIHR Health Technology Assessment - HTA (UK), ref: 10/57/67

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Primary composite outcome of death from any cause or hospitalization for heart failure	13/10/2022	22/12/2022	Yes	No
Protocol article		01/06/2018	19/11/2019	Yes	No
HRA research summary			28/06/2023	No	No
Protocol (other)	V8.3	03/08/2021	26/11/2026	No	No
Statistical Analysis Plan	V1.1	04/05/2022	26/11/2026	No	No
Study website		11/11/2025	11/11/2025	No	Yes