

A phase II/III, observer-blind, randomised, active controlled study to compare the safety and immunogenicity of a meningococcal A conjugate vaccine (PsA-TT) with meningococcal ACWY polysaccharide vaccine administered in healthy children 2 to 10 years of age

Submission date 14/08/2007	Recruitment status No longer recruiting	☒ Prospectively registered ☐ Protocol
Registration date 14/08/2007	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 01/03/2019	Condition category Infections and Infestations	☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Marie-Pierre Preziosi

Contact details

Initiative for Vaccine Research
World Health Organization (WHO)
Department of Immunisation, Vaccines and Biologicals (IVB)
20 Avenue Appia
Geneva
Switzerland
CH-1211
+41 (0)22 791 3744
preziosim@who.int

Type(s)

Scientific

Contact name

Dr Anand Pandit

Contact details

Department of Pediatrics
King Edward Memorial Hospital Research Centre
Sardar Moodliar Road
Rasta Peth
Pune
India
411011

Additional identifiers**Protocol serial number**

RPC219; PsA-TT-003a

Study information**Scientific Title**

A phase II/III, observer-blind, randomised, active controlled study to compare the safety and immunogenicity of a meningococcal A conjugate vaccine (PsA-TT) with meningococcal ACWY polysaccharide vaccine administered in healthy children 2 to 10 years of age

Study objectives

Bacterial meningitis is a life-threatening medical emergency. Morbidity includes hearing loss, chronic seizures, and learning disability. The principal pathogens of bacterial meningitis are *Neisseria meningitidis*, *Streptococcus pneumoniae* and *Haemophilus influenzae* type B. *Neisseria meningitidis* is particularly feared because it has the capability of causing large outbreaks of disease. Endemic meningococcal disease occurs worldwide and is mostly caused by meningococci of serogroups A, B, C, W135 and Y.

Hypothesis:

To compare the immunogenicity of a single dose of the PsA-TT vaccine with that of the Meningococcal A component of the PsACWY vaccine at 28 days after vaccination.

Ethics approval required

Old ethics approval format

Ethics approval(s)

King Edward Memorial Hospital Ethics Committee, Pune, India, 01/08/2007

Study design

Phase II/III observer-blind randomised active-controlled study

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Bacterial meningitis

Interventions

1. One 0.5 ml dose out of a decadose vial of PsA-TT vaccine will be injected intramuscularly (IM) in the right deltoid
2. One 0.5 ml dose of PsACWY vaccine will be injected IM in the right deltoid

Intervention Type

Biological/Vaccine

Phase

Phase II/III

Drug/device/biological/vaccine name(s)

PsA-TT, PsACWY

Primary outcome(s)

The percentage of subjects who show a seroconversion for anti-Meningococcal Polysaccharide A (MenPsA) antibodies, i.e., a four-fold increase in post-immunisation serum titre with respect to pre-immunisation serum titre, at 28 days after a single vaccine dose, as measured by a rabbit complement Serum Bactericidal Assay (rSBA).

Key secondary outcome(s)

Safety:

1. The percentage of subjects with local and systemic post-immunisation reactions during the first four days, adverse events and Serious Adverse Events (SAEs), as measured at 4 and 28 days after vaccination (reactogenicity and short-term safety)
2. The percentage of subjects with SAEs during the entire study duration, as measured at 182 days (6 months) and 364 days (1 year) (long-term safety)

Immunogenicity:

1. The percentage of subjects with anti-MenPsA titre greater than or equal to 1:8 (defined as seroprotection to MenA) at 28 days after a single vaccine dose, as measured by rSBA assay. The percentage of subjects with anti-MenPsA titer greater than or equal to 1:128 (defined as long-term seroprotection to MenA) will be also considered
2. Geometric Mean Titres (GMTs) for anti-MenPsA antibodies at 28 days after a single vaccine dose, as measured by rSBA assay
3. Evaluation of reverse cumulative distribution curves for MenPsA antibody titres at 28 days after a single vaccine dose, as measured by rSBA assay
4. The percentage of subjects who show a seroconversion for anti-MenPsA total Immunoglobulin G (IgG), i.e. a two-fold increase in post-immunisation serum concentration with respect to pre-immunisation serum concentration, at 28 days after a single vaccine dose, as measured by Enzyme-Linked Immunosorbent Assay (ELISA). The percentage of subjects with a four-fold increase in post-immunisation serum concentration with respect to pre-immunisation serum concentration will be also considered

Completion date

20/11/2007

Eligibility

Key inclusion criteria

1. Age 2 to 10 years of age (both included)
2. Written informed consent obtained from parents or legal guardian of the child
3. Free of obvious health problems as established by medical history including physical examination and clinical judgment of the investigator
4. Parents or legal guardian capable and willing to bring their child or to receive home visits (for their child) for all follow-up visits
5. Residence in the study area
6. Fully vaccinated according to the local Expanded Program on Immunisation (EPI) schedule

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

2 years

Upper age limit

10 years

Sex

All

Key exclusion criteria

1. Previous vaccination against *Neisseria meningitidis*
2. Known exposure to *Neisseria meningitidis* during the three previous months
3. History of allergic disease or known hypersensitivity to any component of the two study vaccines and/or following administration of vaccines included in the local program of immunisation
4. Administration of any other vaccine within 60 days prior to administration of study vaccines or planned vaccination during the first 28 days after the study vaccination
5. Use of any investigational or non-registered drug within 90 days prior to the administration of study vaccines
6. Administration of immunoglobulins and/or any blood products within 30 days prior to the administration of study vaccines or planned administration during the study period
7. Chronic administration (defined as more than 14 days) of immunosuppressants or other immune-modifying agents within 90 days prior to the administration of study vaccines (including systemic or inhaled corticosteroids, this means prednisone, or equivalent, greater than 0.5 mg/kg /day; topical steroids are allowed)
8. A family history of congenital or hereditary immunodeficiency
9. History of meningitis or seizures or any neurological disorder
10. Major congenital defects or serious chronic illness, including malnutrition (as per investigator's judgment)

11. Acute disease at the time of enrolment (acute disease is defined as the presence of a moderate or severe illness with or without fever) is a temporary exclusion
12. Acute or chronic, clinically significant pulmonary, cardiovascular, hepatic, or renal functional abnormality, as determined by medical history, physical examination or laboratory tests, which in the opinion of the investigator, might interfere with the study objectives
13. Any condition or criteria that in the opinion of the investigator might compromise the well being of the subject or the compliance with study procedures or interfere with the outcome of the study
14. Non residence in the study area or intent to move out within one year

Date of first enrolment

20/08/2007

Date of final enrolment

20/11/2007

Locations

Countries of recruitment

India

Switzerland

Study participating centre

Initiative for Vaccine Research

Geneva

Switzerland

CH-1211

Sponsor information

Organisation

Serum Institute of India Limited (SIIL)

Organisation

Program for Appropriate Technology in Health (PATH)

Organisation

Serum Institute of India (India)

ROR

Funder(s)

Funder type

Charity

Funder Name

Bill and Melinda Gates Foundation

Alternative Name(s)

Bill & Melinda Gates Foundation, Gates Foundation, Gates Learning Foundation, William H. Gates Foundation, BMGF, B&MGF, GF

Funding Body Type

Government organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United States of America

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	15/11/2015		Yes	No
Results article	results	15/11/2015		Yes	No
Results article	results	15/11/2015		Yes	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes