

N3 fatty acids as adjuvant therapy in rheumatoid arthritis (N3-fettsäuren als adjuvante therapie bei rheumatoider arthritis)

Submission date 01/12/2007	Recruitment status No longer recruiting	☐ Prospectively registered
Registration date 03/01/2008	Overall study status Completed	☐ Protocol
Last Edited 03/01/2008	Condition category Musculoskeletal Diseases	☐ Statistical analysis plan
		☐ Results
		☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Babak Bahadori

Contact details

State Hospital Muerzzuschlag
Department of Internal Medicine
Grazerstr. 63-65
Muerzzuschlag
Austria
8680
b.bahadori@gmx.at

Additional identifiers

Protocol serial number

10-184 ex 99/00

Study information

Scientific Title

Study objectives

Omega-3 fatty acids infusions may show earlier and superior antiinflammatory effects in patients with rheumatoid arthritis than oral administered ones (as shown in earlier trials), and we presumed the beneficial effects may be maintained by following oral omega-3 fatty acid administration.

Ethics approval required

Old ethics approval format

Ethics approval(s)

This trial was approved by the ethics committee of the Medical University of Graz on 10th July 2000.

Study design

Randomized, double-blinded, placebo-controlled trial.

Primary study design

Interventional

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Rheumatoid arthritis

Interventions

Intervention group: Infusions of omega-3 fatty acids once daily for 14 days (Infusion speed: 0.5 ml/kg body weight and hour), followed by oral administration of omega-3 fatty acids up to 6 months.

Control group: Placebo infusions once daily for 14 days (Infusion speed: 0.5 ml/kg body weight and hour), followed by oral administration of placebo up to 6 months.

Verum:

Infusions: Fish oil emulsion (Omegaven® Fresenius). 100 ml Omegaven® contained 10 g fish oil with 2.03 g EicosaPentaenoic Acid (EPA) and 2.26 g DocosaHexaenoic Acid (DHA) (and 0.02 mg a-tocopherol as antioxidant)

Capsules: Fish oil. One capsule contained 1 g fish oil with 0.20 g EPA and 0.23 g DHA (and 0.02 mg a-tocopherol as antioxidant)

Placebo:

Infusions: NaCl 0.9% (physiological sodium chloride solution)

Capsules: Paraffin wax

Randomisation was done by a hospital pharmacist who prepared the bottles and other infusion material, so that neither the patients nor the staff could identify them as verum or placebo. Additionally, the study conducting staff was not allowed to enter the inpatient area while the patients received their infusions.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Omega-3 fatty acids

Primary outcome(s)

The following were assessed at baseline, during infusion phase (after week 1 and after week 2) and during capsule phase (after two weeks [end of month 1] and then monthly until the end of month 6):

1. Swollen joints
2. Tender joints
3. Erythrocyte Sedimentation Rate (ESR)
4. C-Reactive Protein (CRP)

Key secondary outcome(s)

The following were assessed at baseline, during infusion phase (after week 1 and after week 2) and during capsule phase (after two weeks [end of month 1] and then monthly until the end of month 6):

1. Disease Activity Score (DAS)
2. Rheumatoid Arthritis Disease Activity Index (RADAI)
3. Health Assessment Questionnaire (HAQ)
4. 36-item Short Form health survey (SF-36)

Completion date

30/04/2004

Eligibility**Key inclusion criteria**

1. Patients at least 19 years of age and should not be older than 85 years
2. Patients suffering from Rheumatoid Arthritis (RA), according to American College of Rheumatology (ACR) criteria, for at least ($\geq$) 6 months
3. Patients with active RA who meet following criteria:
 - 3.1. $\geq$ 6 swollen joints
 - 3.2. $\geq$ 6 tender joints
 - 3.3. $\geq$ 1 of following criteria:
 - a. Joint morning stiffness $\geq$ 45 minutes
 - b. Erythrocyte Sedimentation Rate (ESR) $\geq$ 28 mm
 - c. C-Reactive Protein (CRP) $\geq$ 20 mg/l (normal range: 0 - 9)
4. No pregnant or breast feeding women allowed; monthly negative pregnancy test in women of reproductive age
5. Participant doesn't have a severe concomitant physical or mental illness
6. Participant has to be able to follow the study procedures
7. No severe malfunctioning of the liver, kidneys, bone marrow, blood clotting system
8. Stable dosages of RA medication for at least 1 month prior to baseline:
 - 8.1. DMARD therapy with methotrexate of at least 10 mg per week over 4 months
 - 8.2. Prednisone $\leq$ 10 mg daily
9. No injection of glucocorticoids into a joint within 3 month prior to baseline
10. Participant must not have legal conflict

11. Participant must not have alcohol or drug abuse problems
12. Written informed consent signed by the participant

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Known allergy against fish oil or any other substance of the study medication
2. Participant is younger than 19 years of age
3. Patient simultaneously participates in another trial
4. Participant doesn't sign the written informed consent.
5. No stable dosage of RA medication 1 month prior to baseline
6. Injection of glucocorticoids into a joint within 3 month prior to baseline
7. Pregnant or breast feeding women; positive pregnancy test in women of reproductive age
8. Patient has legal conflicts
9. Patient has alcohol or drug abuse problems
10. Severe concomitant physical or mental illness
11. Participant is physically or mentally unable to follow the study procedures
12. Hereditary disorder of lipid metabolism
13. Blood triglyceride levels >300 mg/dl
14. Patients with uncontrolled diabetes mellitus
15. Severe malfunctioning of the liver, kidneys, bone marrow, blood clotting system:
 - 15.1. Creatinin clearance ≤ 30 ml/min
 - 15.2. Elevated levels of ALanine aminoTransferase (ALT) 2 fold above the upper normal range
 - 15.3. Leucopenia $\leq 2,500/\text{ml}$, red blood cells ≤ 2.5 mill/ml, hemoglobin ≤ 9 mg/dl, platelets $\leq 70,000/\text{ml}$
16. Severe cardiovascular disease
17. Known neoplasms

Date of first enrolment

11/07/2000

Date of final enrolment

30/04/2004

Locations**Countries of recruitment**

Austria

Study participating centre
State Hospital Muerzzuschlag
Muerzzuschlag
Austria
8680

Sponsor information

Organisation
Medical University of Graz (Austria)

ROR
<https://ror.org/02n0bts35>

Funder(s)

Funder type
Other

Funder Name
Investigator-funded (This trial is a physician initiated study) (Austria)

Funder Name
Fresenius Kabi Austria (Provided Omega-3 infusions only)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration