

Remote ischaemic preconditioning to protect the myocardium during abdominal aortic aneurysm repair

Submission date 12/09/2003	Recruitment status No longer recruiting	☐ Prospectively registered
		☐ Protocol
Registration date 12/09/2003	Overall study status Completed	☐ Statistical analysis plan
		☒ Results
Last Edited 16/07/2009	Condition category Circulatory System	☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Ziad Ali

Contact details

Department of Vascular Surgery
Box 201
Addenbrooke's Hospital
Long Road
Cambridge
United Kingdom
CB2 2QQ

Additional identifiers

Protocol serial number

N0544122070

Study information

Scientific Title

Study objectives

Patients undergoing major vascular surgery are at significant risk of developing postoperative myocardial complications, particularly myocardial infarction. Protecting the heart during the perioperative period could provide a method to reduce morbidity and mortality after vascular surgery. Ischaemic preconditioning is a well recognised phenomenon, whereby a brief period of ischaemia followed by reperfusion prior to a prolonged ischaemic event can provide protection from cellular injury. Protection can be performed either by a stimulus to the myocardium itself or by ischaemia at a site distant to the heart. The aim of this study is to determine whether remote ischaemic preconditioning confers a reduction in myocardial damage in patients undergoing abdominal aortic aneurysm repair.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Aortic aneurysm repair

Interventions

Randomised controlled trial:

1. Abdominal aortic aneurysm repair alone
2. Abdominal aortic aneurysm repair with intermittent cross clamping/perfusion of common iliac artery

Postoperatively, patients will have cardiac troponin and creatine kinase MB (CKMB) recorded on days 1, 3 and 7. Holter monitoring is to be continued until 48 h post operation.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

A reduction in myocardial damage as assessed by serum measurement of cardiac troponin and Holter electrocardiogram.

Key secondary outcome(s)

Not provided at time of registration

Completion date

09/01/2006

Eligibility

Key inclusion criteria

72 Subjects aged 18-90.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Other

Lower age limit

18 Years

Upper age limit

90 Years

Sex

All

Key exclusion criteria

Does not meet inclusion criteria

Date of first enrolment

10/01/2003

Date of final enrolment

09/01/2006

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Department of Vascular Surgery

Cambridge

United Kingdom

CB2 2QQ

Sponsor information

Organisation

Department of Health (UK)

Funder(s)

Funder type

Research council

Funder Name

Cambridge Consortium - Addenbrookes (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	11/09/2007		Yes	No