

# Effectiveness of extracorporeal shock wave therapy in patients with proximal plantar fasciitis

**Submission date**  
02/07/2009

**Recruitment status**  
No longer recruiting

Prospectively registered

Protocol

**Registration date**  
18/09/2009

**Overall study status**  
Completed

Statistical analysis plan

Results

**Last Edited**  
27/08/2014

**Condition category**  
Musculoskeletal Diseases

Individual participant data

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Ms Hongying Chen

### Contact details

ST833  
The Hong Kong Polytechnic University  
Hong Kong  
China

## Additional identifiers

## Study information

### Scientific Title

Changes of proximal plantar fascia microcirculation after extracorporeal shock wave therapy in patients with proximal fasciitis: a double blinded randomised controlled trial

### Study objectives

1. There will be an increase in microcirculation at the proximal plantar fascia (PPF) in chronic plantar fasciitis patients

2. Short term and long term changes no microcirculation can be observed after application of extracorporeal shock wave therapy

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Ethical Committee of the Hong Kong Polytechnic University, 18/06/2009, ref: HSEARS20090618004

### **Study design**

Double-blinded randomised controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Chronic plantar fasciitis

### **Interventions**

Patients will be randomised by drawing cards to receive 3 or 6 sessions of radial extracorporeal shock wave therapy (ESWT) treatment (Storz Medical, Duolith SD, Switzerland), or no active treatment (control). The outcome measures will be taken before, immediately after, at 3, 6 and 12 months after intervention.

Contact details for patient information sheet:

Dr Fu Siu Ngor

The Hong Kong Polytechnic University

Rm ST533, Yuk Choi Road

Hung Hom, Kowloon

Hong Kong

Tel: +852 27666726

Email: rsamyfu@polyu.edu.hk

### **Intervention Type**

Other

### **Phase**

Not Applicable

### **Primary outcome(s)**

Microcirculation index, measured before, immediately after, at 3, 6 and 12 months after intervention

### **Key secondary outcome(s)**

1. Plantar fascia thickness, measured before and 6 and 12 months after intervention
2. Ankle range of motion, measured before and 6 and 12 months after intervention
3. Foot Function Index, measured before, immediately after, at 3, 6 and 12 months after

intervention

4. Visual Analogue Scale (VAS), measured before and 6 and 12 months after intervention

**Completion date**

01/09/2011

## Eligibility

**Key inclusion criteria**

1. Aged between 18 and 60 years (either sex)
2. Suffered from proximal heel pain for more than 3 months

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Upper age limit**

60 years

**Sex**

All

**Key exclusion criteria**

1. Surgery in the treatment area
2. Peripheral vessel diseases
3. Diabetes mellitus
4. Peripheral neuropathy
5. Foot fracture
6. Ankle sensation loss

**Date of first enrolment**

01/09/2009

**Date of final enrolment**

01/09/2011

## Locations

**Countries of recruitment**

China

## Study participating centre

ST833

Hong Kong

China

-

## Sponsor information

### Organisation

The Hong Kong Polytechnic University (China)

### ROR

<https://ror.org/0030zas98>

## Funder(s)

### Funder type

University/education

### Funder Name

The Hong Kong Polytechnic University (China) - Department of Rehabilitation Sciences

## Results and Publications

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/10/2013		Yes	No