

Different routes of progesterone administration for sustaining implantation in in vitro fertilisation patients: impact on pregnancy and live birth rates

Submission date 02/08/2023	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 15/08/2023	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 27/12/2024	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Progesterone administration is considered an essential therapy for luteal phase support in patients undergoing in-vitro fertilization (IVF) technologies. This study aims to compare the administration of a combined protocol of vaginal gel progesterone (90 mg/daily) and oral dydrogesterone (20 mg/daily), with the administration of micronized vaginal progesterone alone (800 mg daily) to support the luteal phase during embryo implantation in patients undergoing in vitro fertilization.

Who can participate

Infertile patients undergoing standard in-vitro fertilization (IVF) or intracytoplasmic sperm injection (ICSI) treatment at One Day Medical Center IVF Unit

What does the study involve

Ovarian stimulation:

All the patients will undergo a mild ovarian stimulation protocol:

Subcutaneous injections of recombinant follicle-stimulating hormone (rFSH) will be started at a dose of 75-150 IU (international unit). Growing ovarian follicles will be detected and measured using ultrasonography at two-day intervals. Estradiol blood level will be measured at two days intervals. Subcutaneous injections of Gonadotropin-releasing hormone antagonist (GnRH ant) (0.25 mg/day) will be administered when the leading follicle measures 14 mm. The rFSH dose will be adjusted when necessary according to the follicular size and estradiol level. Final oocyte maturation will be induced by subcutaneous injection of a dose of 10,000 IU of Human Chorionic Gonadotropin (hCG).

Oocyte recovery and embryo transfer

Oocyte retrieval will be carried out at 34-36 hours after hCG injection. Recovered oocytes will be inseminated by standard IVF or ICSI. Up to 3 embryos will be transferred into the patient's uterus on day three after the oocyte insemination.

What are the possible benefits and risks of participating?

Benefits: the applied supplementation therapy for luteal phase support after embryo replacement could increase the pregnancy and live birth rates.

Risks: the applied supplementation therapy for luteal phase support after embryo replacement could reduce the chance of pregnancy and live birth.

Where is the study run from?

One Day Medical Center s.r.l. (Italy)

When is the study starting and how long is it expected to run for?

October 2016 to December 2022

Who is funding the study?

One Day Medical Center s.r.l. (Italy)

Who is the main contact?

Leonardo Rinaldi, Leo.rinaldi@tiscali.it

Contact information

Type(s)

Principal investigator

Contact name

Dr Leonardo Rinaldi

Contact details

Via Attilio Ambrosini 114

Roma

Italy

00147

+39 (0)3356798557

leo.rinaldi@tiscali.it

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

oneday20161

Study information

Scientific Title

Oral dydrogesterone associated with vaginal micronized progesterone supplementation vs vaginal micronized progesterone alone supplementation for luteal phase support in in vitro fertilization patients: effects on pregnancy and live birth rates

Study objectives

To compare the administration of a combined protocol of progesterone vaginal gel and oral dydrogesterone, with the administration of micronized vaginal progesterone alone for luteal phase support in patients undergoing GnRH antagonist-mild stimulation regimen

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 10/12/2016, One Day Ethics Commission (Via Attilio Ambrosini 114, Roma, 00147, Italy; +39 (0)645212028; amministrazione@onedaymedicalcenter.it), ref: ethics12-2016

Study design

Prospective open randomized study

Primary study design

Interventional

Study type(s)

Treatment, Efficacy

Health condition(s) or problem(s) studied

Luteal phase supplementation in assisted reproduction

Interventions

All the patients are going to have a mild ovarian stimulation protocol: rFSH will be started at a dose of 75-150 IU on the base of AMH level. GnRH antagonist (0.25 mg /day) will be administered when the leading follicle measures 14 mm. The FSH dose will be adjusted when necessary according to the follicular size and estradiol level. Final oocyte maturation will be induced by a dose of 10,000 IU of hCG. Recovered oocytes will be inseminated by standard IVF or intracytoplasmic sperm injection (ICSI). Up to three embryos will be transferred on day 3 after oocyte insemination.

Randomization will be performed using a computer-generated random assignment schedule for each patient. Sealed and numbered envelopes will be used to conceal the treatment allocation until randomization. On the day of oocyte recovery, the patients will be randomized into two groups: Group A will receive vaginal administration of capsules of micronized progesterone (800 mg/day) for luteal phase support; Group B will receive a combination of vaginal administration of a gel of micronized progesterone (90 mg/day) and oral dydrogesterone 10 mg twice a day for luteal phase support. The duration of the treatment is 14 - 21 days; follow-up will be around 9 months for each patient, from the established pregnancy to the delivery.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Oral dydrogesterone, micronized vaginal progesterone

Primary outcome(s)

Clinical pregnancy measured using a blood test to evaluate the level of human chorionic gonadotropin (hCG), which indicates the eventual embryo implantation and pregnancy, 14 days after embryo replacement in the patient's uterus

Key secondary outcome(s)

Embryo implantation rate measured using ultrasonography at 4 weeks following a positive hCG test to confirm the clinical pregnancy

Completion date

23/12/2022

Eligibility

Key inclusion criteria

1. Infertility attributable to tubal factors, moderate endometriosis, male factor, or idiopathic infertility
2. Serum hormonal profile within the normal range (AMH >1 ng/ml)
3. Regular menstrual cycles
4. The presence of a normal uterine cavity
5. Body mass index (BMI) of 20–26 kg/m²

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

27 years

Upper age limit

41 years

Sex

Female

Total final enrolment

620

Key exclusion criteria

1. Infertility attributable to genetic factors or severely reduced ovarian reserve
2. Serum hormonal profile showing a reduced ovarian reserve (AMH <0.9 ng/ml)
3. Irregular menstrual cycles or amenorrhea
4. Congenital or acquired uterine malformation
5. A body mass index (BMI) <20 or >26 kg/m²

Date of first enrolment

01/01/2017

Date of final enrolment

01/11/2022

Locations

Countries of recruitment

Italy

Study participating centre

One Day Medical Center - PMA Unit

Via Attilio Ambrosini 114

Roma

Italy

00147

Sponsor information

Organisation

One Day Medical Center srl

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

One Day Medical Center s.r.l

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due to lack of consensus

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		21/12/2024	27/12/2024	Yes	No