

A Pilot Study of Continuous Negative Pressure (CNEP) In Bronchiolitis

Submission date 05/03/2004	Recruitment status Stopped	☐ Prospectively registered ☐ Protocol
Registration date 05/03/2004	Overall study status Stopped	☐ Statistical analysis plan ☐ Results
Last Edited 09/08/2021	Condition category Respiratory	☐ Individual participant data ☐ Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Martin Samuels

Contact details
Academic Department of Paediatrics, University Hospital of North Staffordshire
Stoke on Trent
United Kingdom
ST4 6QG
+44 (0)1782 552832
samuels@doctors.org.uk

Additional identifiers

Study information

Scientific Title
A Pilot Study of Continuous Negative Pressure (CNEP) In Bronchiolitis

Study objectives
Not provided at time of registration

Ethics approval required
Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Bronchiolitis

Interventions

Infants were randomised to either standard treatment, or standard with CNEP.

Inspired oxygen requirement was measured with a calibrated oxygen analyser placed close to the infant's nose in a headbox, in which the oxygen flow was adjusted to maintain SaO₂ 96-99% at rest, measured with a Nellcor N-200 pulse oximeter.

When an infant fulfilled the entry criteria, written informed consent was obtained from parents and baseline data was collected.

Patients were then randomised to either conventional therapy or conventional therapy plus CNEP. The randomisation was performed using a stratification scheme to achieve a measure of balance in the treatment groups.

Conventional therapy in these hospitals included the use of additional inspired oxygen and bronchodilators. Infants were intubated and positive pressure ventilation (PPV) was initiated in the presence of respiratory acidosis with a pH below 7.25, hypercapnia, hypoxaemia in spite of additional inspired oxygen, recurrent apnoea and respiratory fatigue.

CNEP was applied using purpose built systems (Horner and Wells Ltd, Chelmsford, UK, and DHB Tools Ltd, Leamington Spa, UK).

Treatment was begun with -4 cmH₂O of CNEP. If the FiO₂ required to achieve normal SaO₂ did not decrease within the next 30 min, CNEP was decreased to -6 cmH₂O. Weaning from CNEP was attempted after a treatment period of at least 24 hours and usually in the presence of an FiO₂<0.3.

Infants were fed by either nasogastric tube, or intravenously, according to the degree of respiratory distress.

Heart rate, respiratory rate and FiO₂ were recorded hourly if the infants were at rest when they were treated with additional inspired oxygen.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Not provided at time of registration

Key secondary outcome(s)

Not provided at time of registration

Completion date

31/12/2004

Reason abandoned (if study stopped)

Objectives no longer viable

Eligibility

Key inclusion criteria

Forty two infants admitted with bronchiolitis, who needed more than 40% oxygen therapy to maintain normal levels of arterial oxygen saturation. This was a pilot study. Forty two patients were enrolled between January 1991 and April 1992 in two paediatric hospitals in England (North Staffordshire Hospital and Royal Berkshire Hospital) if they fulfilled the following criteria:

1. Clinical diagnosis of bronchiolitis (irrespective of whether respiratory syncytial virus was isolated)
2. Age 1 year
3. Presence of respiratory failure with fractional inspired oxygen (FiO₂) 0.4 to achieve an arterial oxygen saturation (SaO₂) of 96-99%.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Neonate

Sex

Not Specified

Key exclusion criteria

Infants with bronchopulmonary dysplasia (BPD), congenital cardiac, pulmonary or neuromuscular diseases or signs of upper airway obstruction were excluded.

Date of first enrolment

01/01/2004

Date of final enrolment

31/12/2004

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Academic Department of Paediatrics, University Hospital of North Staffordshire
Stoke on Trent
United Kingdom
ST4 6QG

Sponsor information

Organisation

University Hospital of North Staffordshire (UK)

Funder(s)

Funder type

Government

Funder Name

North Staffordshire and Royal Berkshire Hospitals

Results and Publications

Individual participant data (IPD) sharing plan**IPD sharing plan summary**

Not provided at time of registration