

A randomised study of sulindac and epalrestat in diabetic retinopathy

Submission date 11/10/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 12/10/2006	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 16/10/2009	Condition category Eye Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mrs Yukiko Hattori

Contact details
Department of Molecular Oncology
Division of Molecular and Cellular Biology Institute on Aging and Adptation
Shinshu University School of Medicine
3-1-1 Asahi
Matsumoto
Japan
390-8621

Additional identifiers

Study information

Scientific Title

Acronym
RSSEDR

Study objectives

The primary purpose is to evaluate the effects of long term treatment with sulindac and epalrestat in diabetic retinopathy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Institutional Review Board of Asama General Hospital, date of Approval 01/08/1997 (reference number: AGH 97-01).

Study design

Interventional randomised single-blind uncontrolled parallel-assignment trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Diabetic Retinopathy

Interventions

1. The sulindac group: patients took 100 mg of sulindac twice a day
2. The epalrestat group: patients took 50 mg of epalrestat three times a day
3. The control group: patients took no additional medications

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Sulindac, epalrestat

Primary outcome(s)

1. Seven-field stereoscopic retinal photography at baseline, and yearly intervals
2. Endpoint if they were diagnosed with proliferative diabetic retinopathy

Key secondary outcome(s)

1. Glycosylated Hemoglobin (GHb) values determine at monthly intervals
2. Remain on their medications for three years
3. Endpoint if exhibited a dipstick proteinuria of more than or equal to 1+

Completion date

31/10/2000

Eligibility

Key inclusion criteria

1. Clinical diagnosis of type two diabetes
2. Age 20 years or older
3. Patient consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Having diabetes for less than five years
2. HbA1c less than 8.0%
3. Taking other medications except for oral hypoglycemic agents or insulin injections
4. Hypertension
5. Inability to understand the implications of the protocol

Date of first enrolment

01/10/1997

Date of final enrolment

31/10/2000

Locations**Countries of recruitment**

Japan

Study participating centre

Department of Molecular Oncology

Matsumoto

Japan

390-8621

Sponsor information**Organisation**

Asama General Hospital (Japan)

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Internally funded by participant centre, Asama General Hospital (Japan)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration