

Is a combination of individual consultations, text message reminders and interaction with a Facebook page more effective than educational sessions in encouraging university students to increase their physical activity levels?

Submission date 19/12/2019	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 05/03/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 17/07/2023	Condition category Other	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Despite the health benefits, physical activity levels among adults in Jordan are low. The aim of this study was to investigate whether a behavioural intervention could increase physical activity levels, reduce body mass index (BMI) and increase confidence in maintaining physical activity among students in universities in Jordan.

Who can participate?

University students in Jordan who are physically able to do moderate exercise

What does the study involve?

The participants will be allocated to one of two groups. One group will receive three educational lectures on the benefits of exercise, potential barriers to increasing physical activity and suggestions on how to change their behaviour to involve more physical activity. The other group will have a face-to-face consultation with individualised suggestions on how they could increase their physical activity. They will also receive 12 calls and regular text messages to encourage them to move more and have access to a Facebook page with tips and motivational messages. All participants will wear a pedometer for 6 months to count their daily steps. They will also have their blood pressure, height and weight measured before and after the 6-month intervention period, and will fill out questionnaires on their physical activity levels and confidence in exercising at these times.

What are the possible benefits and risks of participating?

Participants might benefit from the education and encouragement to increase their physical activity. There were no additional risks of participation.

Where is the study run from?
Philadelphia University (Jordan)

When is the study starting and how long is it expected to run for?
December 2015 to September 2016

Who is funding the study?
Philadelphia University (Jordan)

Who is the main contact?
Dr Eman Alsaleh, ealsaleh@philadelphia.edu.jo

Contact information

Type(s)
Scientific

Contact name
Dr Eman Alsaleh

Contact details
Philadelphia University
Jerash Road
Amman
Jordan
19392
+962 64799000
ealsaleh@philadelphia.edu.jo

Additional identifiers

Clinical Trials Information System (CTIS)
Nil known

Protocol serial number
Nil known

Study information

Scientific Title
A randomised controlled trial investigating the efficacy of a behavioural-based web and phone intervention to increase physical activity among university students in Jordan

Study objectives
A behavioural intervention that consists of multiple components such as goal-setting, self-monitoring and feedback in a combination of effective delivery systems based on mobile and Facebook motivates students to increase their physical activity levels.

Ethics approval required
Old ethics approval format

Ethics approval(s)

Approved November 2016, Ethical board at Philadelphia University (Deanship of Scientific Research and Graduate Studies, Jerash Rd, Amman 19392, Jordan; +962 6 4799 000 Ext. 2366 or 2376; kma@philadelphia.edu.jo), no reference number

Study design

Two-arm single-centre randomized controlled trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Physical activity in university students

Interventions

The study is two-group single-centre randomised controlled trial (n=150) in which a behavioural intervention was compared with usual care in increasing physical activity levels among healthy students in one Jordanian University. The participants were randomly allocated to the two groups (the control and intervention group) by opening opaque sealed sequence envelopes according to the participant's numbers, which contained the name of the group.

The intervention consisted of delivering behavioural strategies by a single face-to-face individualised consultation with the researcher and six telephone call consultations (one call each month) supplemented by motivational SMS text messages reminders (one SMS weekly). The tailored consultation aimed to help participants to integrate moderate physical activity into their daily routine by performing brisk walking of 30 min daily for a minimum of 5 days per week or at least 10,000 steps per day using a bracelet pedometer to count their steps. In addition, the participants were contacted by a Facebook page (Let's walk) that aimed to deliver a motivational tips and advice to increase their physical activity levels.

The educational sessions were provided as three scheduled educational lectures about the benefits of physical activity, potential barriers to physical activity and using behavioral change strategies to increase physical activity levels. The duration of each educational lecture was 1 h. The lectures were given at the first month, the third month and the fifth month.

Intervention Type

Behavioural

Primary outcome(s)

1. Steps per day measured using a bracelet pedometer worn by the participants around their wrists during the day. This bracelet pedometer measured individuals' steps per day. The international recommendation for steps per day is 10,000-step/ day to achieve the international recommendations of 30 minutes of brisk walking per day.
2. Frequency of walking assessed using the International Physical Activity Questionnaire (IPAQ) at baseline and 6 months
3. Duration of walking assessed using the International Physical Activity Questionnaire (IPAQ) at baseline and 6 months

4. Frequency of walking assessed using the International Physical Activity Questionnaire (IPAQ) at baseline and 6 months

Key secondary outcome(s)

1. Blood pressure was measured by the researchers using automated electronic blood pressure monitor. To ensure consistency with this device, blood pressures of the participants was measured by the same BP monitor at baseline and at 6 months. Blood pressure was measured based on the criteria of measuring blood pressure determined by the American Heart Association, the British Hypertension Society and the UK Medicines and Healthcare products Regulatory Agency (MHRA).
2. Body mass index calculated from height and weight. Body weight was measured with the same device at baseline and 6 months. Height was measured at baseline only.
3. Exercise self-efficacy assessed using the Exercise Self-Efficacy Scale (ESES) at baseline and 6 months. The scale was administered to identify the extent of the confidence of the subjects in performing the required levels of physical activity (regular moderate physical activity of 30 min 5 days a week) at a criterion level in specific situations by rating the level of their confidence.

Completion date

01/09/2016

Eligibility

Key inclusion criteria

1. Students of any subject at Philadelphia University
2. No medical condition that prevents them from doing moderate physical activity such as brisk walking

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

146

Key exclusion criteria

Illness or health conditions that prevent them from participating in physical activity

Date of first enrolment

01/02/2016

Date of final enrolment

01/03/2016

Locations

Countries of recruitment

Jordan

Study participating centre

Philadelphia University

Jerash Rd

Amman

Jordan

19392

Sponsor information

Organisation

Philadelphia University

Funder(s)

Funder type

University/education

Funder Name

Philadelphia University

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from the author (Dr Eman Alsaleh, email address: ealsaleh@philadelphia.edu.jo). The data includes the intervention components and materials such as the consultations contents and the motivational tips and messages. These data can be obtained by sending a request by email after the results have been published in a journal.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		28/06/2023	17/07/2023	Yes	No

