

# Efficacy and safety of praziquantel, tribendimidine and mebendazole in patients infected with *Clonorchis sinensis*

<b>Submission date</b> 03/05/2011	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
<b>Registration date</b> 19/05/2011	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 16/12/2015	<b>Condition category</b> Infections and Infestations	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Miss Lili Xu

**Contact details**  
207 Rui Jin Er Lu  
Shanghai  
China  
200025  
-  
luckylilyxu@yahoo.com.cn

## Additional identifiers

## Study information

**Scientific Title**  
Efficacy and safety of praziquantel, tribendimidine and mebendazole in patients infected with *Clonorchis sinensis*: a single-centre, open-label, randomised controlled study

**Study objectives**

Clonorchiasis is one of important food-borne trematodiasis which is highly prevalent in east and southeast Asia, especially in China, Korea, northern parts of Vietnam, and the far eastern part of Russia. Chemotherapy is the mainstay for the control of food-borne trematodiasis. Currently, praziquantel is the drug of choice for clonorchiasis. According to the recommendation from World Health Organisation (WHO), the appropriate treatment schedule is 25 mg/kg thrice daily for up to 2 days which results in cure rates of 94-100%. Promotion of this dose schedule of praziquantel in mass treatment of clonorchiasis may have certain difficulty, while administration of single dose or reduction of treatment course results in less or unstable efficacy. Therefore, it still needs to develop the new drugs against *C. sinensis*. In recent years, we found that oral administration of single-dose tribendimidine and mebendazole and multiple-dose albendazole are efficacious. Tribendimidine exhibits not only potential effect against *C. sinensis* but also potential effect against juvenile *C. sinensis*. Albendazole was reported to be effective against *C. sinensis* in rats, and similar results were also shown in experimentally infected dogs. Up to 2005, an experimental study indicated that albendazole and mebendazole showed a potential effect against adult *C. sinensis* in rats and their single complete curative dose was 150 mg/kg. We aim to assess the efficacy and safety of tribendimidine, mebendazole and albendazole compared with that of praziquantel in patients with parasitologically confirmed *Clonorchis sinensis*.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Ethical Review Committee of the National Institute of Parasitic Diseases, China CDC, 29/04/2011, no. 2011042901

### **Study design**

Single-centre open-label randomised controlled study

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Clonorchiasis

### **Interventions**

Current interventions as of 14/08/2012:

The intervention consists of four treatments, all the medicine is administered orally to each person

1. Multiple-dose praziquantel serves as control, and the total dose is 75 mg/kg (18.75mg/kg twice daily for two days consecutively).
2. Single-dose tribendimidine 400mg.
3. Two doses, on the same day, of Tribendimidine 200mg. The total dose is 400 mg.
4. Single-dose mebendazole 400mg.

All the participants are supervised after treatments, and are asked to report any potential drug related symptoms at 3h and 24h after each time of taking medicine.

Previous interventions until 14/08/2012:

The intervention consists of four treatments, all the medicine is administered orally to each person

1. Multiple-dose praziquantel serves as control, and the total dose is 75 mg/kg(18.75mg/kg twice daily for 2 days consecutively)
2. Single-dose tribendimidine(400mg)
3. Multiple-dose mebendazole, and the total dose is 1500mg (500mg once daily for 3 days consecutively )
4. Multiple-dose albendazole, the total dose is 3200mg( 400mg twice daily for 4 consecutive days).

All the participants are supervised after treatments, and are asked to report any potential drug related symptoms at 24h, 48h, 72h, 96h, 120h and 144h after the first dose

### **Intervention Type**

Drug

### **Phase**

Not Applicable

### **Drug/device/biological/vaccine name(s)**

Praziquantel, tribendimidine, mebendazole, albendazole

### **Primary outcome(s)**

1. Egg reduction rate (ERR) and cure rate (CR), measured using the Kato-Katz stool examination method
2. Adverse reactions, measured using a standardised questionnaire. The intensity of adverse events was graded as mild, moderate, severe and serious

Assessed at 4 weeks after treatment

### **Key secondary outcome(s)**

All the participants who are to receive the tribendimidine treatment will accept checks of blood, urine, electrocardiogram, liver function and kidney function before and after the treatment. The abnormal indexes will be supervised and recorded.

### **Completion date**

15/08/2012

## **Eligibility**

### **Key inclusion criteria**

Current inclusion criteria as of 14/08/2012:

Eligible patients are residents of Hunan province aged 15 to 65 years who have Clonorchis sinensis infections, both sexes.

Previous inclusion criteria until 14/08/2012:

Eligible patients are residents of Guangdong province aged 2 years and above who had Clonorchis sinensis infections, both sexes

### **Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Key exclusion criteria**

1. Pregnant or lactating women
2. Presence of any abnormal medical condition, such as heart disease, high blood pressure, severe malnutrition, severe liver and kidney disease, psychiatric and neurologic disorders
3. Use of praziquantel , tribendimidine, albendazole and mebendazole or any anthelmintic treatment within the past month
4. Enrolled in any other clinical investigation during the study

**Date of first enrolment**

25/06/2012

**Date of final enrolment**

15/08/2012

## **Locations**

**Countries of recruitment**

China

**Study participating centre**

207 Rui Jin Er Lu

Shanghai

China

200025

## **Sponsor information**

**Organisation**

The National Institute of Parasitic Diseases (China)

**ROR**

<https://ror.org/04wktzw65>

# Funder(s)

## Funder type

Government

## Funder Name

National Institute of Parasitic Diseases, China CDC (China)

# Results and Publications

## Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	14/08/2014		Yes	No