

# Early proprioceptive stimulations in mechanically ventilated critically ill patients

<b>Submission date</b> 05/03/2023	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 14/03/2023	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 29/03/2023	<b>Condition category</b> Other	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

After a stay in the intensive care unit (ICU), patients often have conditions unlinked to their initial reason for hospitalization. These conditions affect the physical, psychological, and cognitive aspects of patients' lives. They are grouped under the term post-intensive care syndrome (PICS). Recognition of this syndrome has triggered a paradigm shift in ICU practices in the past few years. One aspect that has been discussed a lot is prolonged bed rest. Helping patients be mobile or early mobilization in the ICU has then been studied, and its benefits have been proven. However, the timing, type, and intensity of the intervention still need to be defined. The present study focuses on a treatment called functional proprioceptive stimulations (FPS). This technique uses tendon vibrations to reproduce the signatures of the movements such as those used in walking (flexion/extension). It can be applied to unconscious or sedated patients, allowing it to be implemented early during the ICU stay. Some studies have already demonstrated the benefits of proprioceptive vibrations. However, these benefits have only been studied at the chronic stage of neurological diseases. The team hypothesize that early FPS might lessen the harmful effects of an ICU stay. In this study, better functional status and less muscle wasting are expected at discharge from the ICU. This study will help clarify the value of such an early and intensive intervention in mechanically ventilated critically ill patients.

### Who can participate?

Patients hospitalized in the ICU of a single center

### What does the study involve?

To test the hypothesis, a randomized controlled trial is being conducted on patients in ICU who have been mechanically ventilated for at least 48 hours. These patients are stratified into two groups according to the presence or absence of a central nervous system lesion. Within five days after admission, patients are included and either receive FPS added to standard mobilization or standard mobilization alone. FPS are delivered to the joints of the lower limbs four times a week during the ICU stay. The primary outcome measure assesses functional status at discharge from the ICU. Secondary outcomes include muscle strength, muscle wasting, cognitive impairment, and spasticity when appropriate. A smaller incidence of ICU-acquired weakness is expected in the experimental group than in the control group.

What are the possible benefits and risks of participating?

The expected benefits are a limitation of the harmful effects of prolonged bed rest and immobilization, especially on the functional status. The risks are minimal: apart from possible transient skin reddening, no side effect related to using the FPS is expected in this study.

Where is the study run from?

Hôpital Bicêtre (France)

When is the study starting and how long is it expected to run for?

May 2022 to May 2024

Who is funding the study?

1. The equipment used in this study belongs to the Bicêtre hospital. And one of the investigators is supported by a PhD grant (CIFRE fellowship N°2020/1605) co-funded by the National Association for Research in Technology (Association Nationale de la Recherche Technique; ANRT) and Techno Concept (Manosque, France).

2. Association MAPAR (Mise Au Point Anesthésie Réanimation) (France)

3. Assistance Publique - Hôpitaux de Paris/Public Assistance Hospital of Paris (France)

Who is the main contact?

Bernard Vigue, [bernard.vigue@aphp.fr](mailto:bernard.vigue@aphp.fr) (France)

## Contact information

### Type(s)

Principal investigator

### Contact name

Dr Bernard Vigué

### ORCID ID

<https://orcid.org/0000-0002-6244-1544>

### Contact details

Département d'Anesthésie réanimation - Hôpital Universitaire de Bicêtre

78, avenue du Général Leclerc

Le Kremlin Bicetre

France

94270

+33145213430

[bernard.vigue@aphp.fr](mailto:bernard.vigue@aphp.fr)

### Type(s)

Scientific

### Contact name

Mrs Florence Martinache

### ORCID ID

<https://orcid.org/0000-0003-1618-5532>

### Contact details

Département d'Anesthésie réanimation - Hôpital Universitaire de Bicêtre  
78, avenue du Général Leclerc  
Le Kremlin Bicetre  
France  
94270  
+33145213430  
florence.martinache@universite-paris-saclay.fr

### Type(s)

Public

### Contact name

Dr Bernard Vigué

### Contact details

Département d'Anesthésie réanimation - Hôpital Universitaire de Bicêtre  
78, avenue du Général Leclerc  
Le Kremlin Bicêtre  
France  
94270  
+33145213430  
bernard.vigue@aphp.fr

## Additional identifiers

### Protocol serial number

RCB 2022-A01277-36

## Study information

### Scientific Title

Assessment of the effects of early proprioceptive stimulations on the functional status of mechanically ventilated critically ill patients: a randomized controlled trial

### Acronym

REA-MOUV

### Study objectives

Early functional proprioceptive stimulations (FPS) mitigate the harmful effect of an ICU stay.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Approved 19/12/2022, Committee for the protection of individuals West 5 (Comité de protection des personnes Ouest V, CHU Pontchaillou - 9, Avenue de la Bataille Flandres Dunkerque Mai 1940, 35033 Rennes Cedex 09, France; +33 (0)2 99 28 25 56; cpp.ouestv@chu-rennes.fr), ref: 22.03429.000144

**Study design**

Single-center interventional randomized controlled study (examiner blinded)

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Early rehabilitation of critically ill patients

**Interventions**

The control group receives standard treatment usually provided in the ICU.

The intervention group receives standard treatment, and in addition, they receive "functional proprioceptive stimulations" (FPS) on the lower limbs (i.e., synchronized tendon vibration applied to the lower limbs). They receive 30-minute PFS sessions four times a week, from inclusion to discharge. FPS are delivered with the Vibramoov® system (Techno Concept, Manosque, France) by a trained physiotherapist. This system consists of twelve vibrators attached to the patients with the help of orthoses and controlled by a computer via Bluetooth. Treatment begins as soon as the physician in charge of the patient considers the patient clinically stable.

Patients are randomized according to a permuted block randomization scheme with a block size of 8. Moreover, patients are stratified into two groups according to the presence or absence of a central nervous system lesion.

**Intervention Type**

Device

**Phase**

Not Applicable

**Drug/device/biological/vaccine name(s)**

Vibramoov® system

**Primary outcome(s)**

Functional status measured using the Physical Function in ICU Test-scored (PFIT-s) at discharge from the ICU

**Key secondary outcome(s)**

1. Cognitive impairments measured using the Montreal cognitive assessment (MoCA) at discharge from the ICU and 3 months after discharge
2. Anxiety and depressive disorders measured using the Hospital anxiety and depression scale (HADS) at discharge from the ICU and 3 months after discharge
3. PTSD symptoms measured using the PTSD Checklist for DSM-5 (PCL-5) at discharge from the ICU and 3 months after discharge
4. Functional status measured using the Physical Function in ICU Test-scored (PFIT-s) 8 to 10 days from admission in the ICU and upon awakening
5. Functional status measured using the ICU Mobility Scale (IMS) upon awakening and on

discharge from the ICU

6. Functional independence measured using the Barthel Index at discharge from the ICU and 3 months after discharge
7. Correlations between functional tests (PFIT-s, IMS, Barthel Index) and cognitive tests (MoCA, HAD, PCL-5) measured using Pearson or Spearman coefficients using the timepoints defined
8. Muscular strength measured using the Medical Research Council-Sum Score (MRC-SS) and with a dynamometric measure of the quadriceps strength upon awakening and then once a week until discharge from the ICU
9. Changes in the quadriceps muscle thickness measured using ultrasound at inclusion, at 4 days post-inclusion, at 10 days post-inclusion, and then once a week until discharge from the ICU
10. Changes in the rectus femoris cross-section measured using ultrasound at inclusion, at 4 days post-inclusion, at 10 days post-inclusion, and then once a week until discharge from the ICU
11. Changes in the rectus femoris echogenicity measured using ultrasound at inclusion, at 4 days post-inclusion, at 10 days post-inclusion, and then once a week until discharge from the ICU
12. Changes in the spasticity measured using the modified Ashworth scale (MAS) and the modified Tardieu scale (MTS) on discharge from the ICU (only for patients with a central nervous system lesion)
13. Number of delirium days in ICU measured using the Confusion Assessment Method for ICU (CAM-ICU)
14. Number of days ventilation-free in the first 28 days in ICU measured using patient medical records at one timepoint at the end of the study
15. Incidence of ICU-Acquired Weakness (ICU-AW), diagnosed by an MRC-SS < 48/60, measured using patient medical records at one timepoint at the end of the study
16. Length of ICU stay measured in days using patient medical records at one timepoint at the end of the study
17. Length of hospital stay measured in days using patient medical records at one timepoint at the end of the study
18. Mortality rate in hospital at 3 months and 6 months measured using patient medical records at one timepoint at the end of the study

### **Completion date**

30/05/2024

## **Eligibility**

### **Key inclusion criteria**

1. Adult ( $\geq 18$  years old)
2. Hospitalised in the intensive care unit (ICU)
3. At least 48 hours of mechanical ventilation
4. Functionally independent two weeks before the admission to ICU (Barthel Index  $\geq 70$ , estimated by interviewing relatives)

### **Participant type(s)**

Patient

### **Healthy volunteers allowed**

No

### **Age group**

Adult

## **Lower age limit**

18 years

## **Sex**

All

## **Key exclusion criteria**

1. Hospitalised in ICU for more than 5 days
2. Hyperacute phase (absence of hemodynamic balance)
3. Orthopedic injury preventing the implementation of the protocol
4. Serious psychiatric or cognitive history
5. Irreversible disorder with a 6-month mortality greater than 50% or patient with a probable fatal outcome in the ICU
6. Rapidly progressive neuromuscular disease
7. Spinal cord injury
8. Severe traumatic brain injury (Glasgow Coma Scale score  $\leq 8$ )
10. Known pregnancy
11. Patients with phlebitis
12. Patients with active devices such as pacemakers, defibrillators, insulin pumps, neurostimulators, etc.
13. Patients with cardiac arrhythmia
14. Patients with epilepsy
15. Patients with fragile skin or open wounds
16. Not affiliated with the French social security system
17. Patients who speak neither French nor English
18. Persons deprived of liberty and under legal protection

## **Date of first enrolment**

27/02/2023

## **Date of final enrolment**

01/05/2024

## **Locations**

### **Countries of recruitment**

France

### **Study participating centre**

**Hôpital Bicêtre**

78, avenue du Général Leclerc

Le Kremlin Bicêtre

France

94270

## **Sponsor information**

**Organisation**

Association MAPAR (Mise Au Point Anesthésie Réanimation)

**Funder(s)****Funder type**

University/education

**Funder Name**

Association MAPAR (Mise Au Point Anesthésie Réanimation)

**Funder Name**

TechnoConcept

**Funder Name**

Assistance Publique - Hôpitaux de Paris/Public Assistance Hospital of Paris

**Alternative Name(s)**

Assistance Publique Hôpitaux de Paris, Assistance Publique–Hôpitaux de Paris, AP-HP

**Funding Body Type**

Private sector organisation

**Funding Body Subtype**

Trusts, charities, foundations (both public and private)

**Location**

France

**Funder Name**

Association Nationale Recherche Technologie/National Association for Research in Technology

**Results and Publications****Individual participant data (IPD) sharing plan**

The data-sharing plans for the current study are unknown and will be made available at a later date.

## IPD sharing plan summary

Data sharing statement to be made available at a later date

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol file</a>	version 5.0	14/12/2022	14/03/2023	No	No