

Comparison between intraarticular ozone and a placebo in the treatment of knee arthritis

Submission date 23/10/2010	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 08/11/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 12/01/2017	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Osteoarthritis (OA) is the most common type of arthritis, affecting millions of people worldwide. It occurs when the protective cartilage on the end of bones wears away. The bones then rub against one another, which can cause stiffness, pain and a reduction in a person's range of movement. One of the most common joints to be affected is the knee, and in many sufferers, the pain prevents people from moving around leading to muscle weakness and disability. There are a variety of treatments for knee OA, including exercise, medication and surgery (knee replacement). One promising new treatment is the injection of ozone into the knee joint to help alleviate pain. Ozone is a gas made up of three atoms of oxygen joined together, rather than the usual two atoms as found in the oxygen we breathe from the air. The aim of this study is to find out whether ozone injections into the affected knee joint are an effective treatment for knee OA.

Who can participate?

Patients aged between 60 and 85 who have knee OA.

What does the study involve?

Participants are randomly allocated to one of two groups. Those in the first group receive eight, weekly injections of ozone into the knee joint. Those in the second group have a placebo (dummy), consisting of normal air injected into the knee joint. At the start of the study and then again after four weeks, eight weeks and two months, patients in both groups have their pain levels assessed using a range of questionnaires.

What are the possible benefits and risks of participating?

Participants who receive the ozone injections may benefit from an improvement to their pain levels and OA symptoms. There is a small risk of pain in the knee during and after injections.

Where is the study run from?

1. Paulista School of Medicine – Federal University of São Paulo (Brazil)
2. Pró-Vida – Center for Total Health Assistance LLC (Brazil)
3. University of Santo Amaro – Medicine College (Brazil)

When is the study starting and how long is it expected to run for?
February 2010 to June 2015

Who is funding the study?
Sao Paulo Medical School (Brazil)

Who is the main contact?
Professor Carlos César Lopes de Jesus
caceloje@gmail.com

Contact information

Type(s)
Scientific

Contact name
Prof Carlos César Lopes de Jesus

Contact details
Rua Pedro de Toledo, 598
São Paulo
Brazil
04039-001
+55 (0)11 55752970 / 50850248
caceloje@gmail.com

Additional identifiers

ClinicalTrials.gov (NCT)
NCT29102010

Study information

Scientific Title
Comparison between intraarticular ozone and a placebo in the treatment of knee arthritis: A double blind, randomised placebo controlled trial

Acronym
INOPKA

Study objectives
Hypothesis as of 30/12/2016:
Ozone is more effective than a placebo in the treatment of knee osteoarthritis concerning pain relief, improvement of joint function and quality of life.

Original hypothesis:
Ozone is more affective than placebo in the treatment of knee arthritis.

Ethics approval required
Old ethics approval format

Ethics approval(s)

The Ethical Review Board of the Paulista School of Medicine – Federal University of Sao Paulo, 08/12/2010, ref: EPM-UNIFESP

Primary study design

Interventional

Study design

Randomised double-blinded placebo controlled clinical trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Knee osteoarthritis

Interventions

Interventions section as of 11/01/2017:

Study participants attend a baseline visit at which the following procedures are performed: medical history, physical examination, analysis of X-ray of the affected knee and application of the following questionnaires and tests: Visual Analogue Scale (VAS), Geriatric Pain Measure (GPM), Lequesne's Index, Timed Up and Go Test (TUG Test), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), and Short-Form Health Survey (SF-36). Eligible patients are fully informed of the purpose of the study. All patients that fulfill the inclusion criteria signed the informed consent prior to enrolment in the trial. They are instructed to continue their medical treatment according to their physicians' orientations.

With the objective of avoiding selection bias, all included participants are sequentially assigned by the researchers to receive OZ or PBO according to a pre-established computer-generated global randomization list. This is carried out in such a manner as to guarantee each patient an equal chance of receiving the intervention. The randomization list is generated using software ETCETERA, version 2.46, and constituted 98 numbers with the corresponding treatments. Prior to the beginning of the randomization it has been stipulated that group A would be the ozone group and B, the placebo group.

Ozone group (OZ): Participants receive one intra articular injection of ozone 20 µg/ml (10ml) per week for 8 consecutive weeks.

Placebo group (PBO): Participants receive one intra articular injection of 10 ml of air per week for 8 consecutive weeks.

Assessments are performed at baseline (visit 1), 4 weeks (visit 2), 8 weeks (visit 3), and 8 weeks after the end of the injections (visit 4). At the follow-up visits, the same procedures as those described for visit 1 are performed.

Original interventions section:

Patients will be randomized to receive either intra articular ozone, 20µg/ml once a week for 8 consecutive weeks, or intra articular air, once a week, for 8 consecutive weeks.

The follow-up will last for 2 months after the last session of treatment.

Intervention Type

Drug

Phase

Phase III

Drug/device/biological/vaccine name(s)

Ozone

Primary outcome(s)

Primary outcome measures as of 30/12/2016:

Reduction of pain measured according to

1. Visual Analogue Scale (VAS)
2. Lequesne Algofunctional Index
3. Geriatric Pain Measure (GPM)

All outcomes will be measured at baseline, 4 and 8 weeks and 2 months.

Original primary outcome measures:

Efficacy of treatment measured according to

1. Lequesne Algofunctional Index
2. Time Up and Go Test
3. Activities of Daily Living (ADLs)
4. Medical Outcomes Study 36-Item Short Form
5. Health Survey (MOS SF36 Health Survey)
6. Visual Analogue Scale (VAS)
7. Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC Index)

All outcomes will be measured at baseline, 4 and 8 weeks and 2 months.

Key secondary outcome(s)

Secondary outcome measures as of 30/12/2016:

Improvement of joint function and quality of life according to:

1. Time Up and Go Test
2. Medical Outcomes Study 36-Item Short Form
3. Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC Index)

All outcomes will be measured at baseline, 4 and 8 weeks and 2 months

Original secondary outcome measures:

Pain reduction according to

1. Lequesne Algofunctional Index
2. VAS

All outcomes will be measured at baseline, 4 and 8 weeks and 2 months.

Completion date

30/06/2015

Eligibility

Key inclusion criteria

Inclusion criteria as of 30/12/2016:

1. Patients from Outpatients Department of Geriatrics and Gerontology Discipline – Paulista School of Medicine – Federal University of Sao Paulo, Pro-Vida – Centro de Assistencia Integral a Saude, and Outpatients Department – Rheumatology Discipline – Medical College – Santo Amaro University
2. Aged between 60 years and 85 years
3. Clinically and radiologically confirmed knee osteoarthritis according to the American College of Rheumatology (ACR) criteria

Original inclusion criteria:

1. Patients from Outpatients Department of Geriatrics and Gerontology Discipline da Escola Paulista de Medicina - Universidade Federal de São Paulo
2. Aged more than 60 years or less than 85 years
3. Clinically and radiologically confirmed knee arthritis according to the American College of Rheumatology (ACR) criteria

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

All

Key exclusion criteria

Exclusion criteria as of 30/12/2016:

1. Patients from Outpatients Department of Geriatrics and Gerontology Discipline – Paulista School of Medicine – Federal University of Sao Paulo, Pro-Vida – Centro de Assistencia Integral a Saude, and Outpatients Department – Rheumatology Discipline – Medical College – Santo Amaro University,
2. Aged between 60 years and 85 years, and
3. Clinically and radiologically confirmed knee osteoarthritis according to the American College of Rheumatology (ACR) criteria.

Original exclusion criteria:

1. Patients aged less than 60 years or more than 75 years
2. Will not give their permission to be included in the study
3. Patients with mental and/or neurologic deficit
4. Diagnosis of secondary osteoarthrosis, patients with recent knee traumas or
5. Suspected associated knee lesion
6. Patients with coxofemoral articulation affections
7. In the acute phase of the disease

Date of first enrolment

19/11/2010

Date of final enrolment

23/03/2015

Locations

Countries of recruitment

Brazil

Study participating centre

Paulista School of Medicine – Federal University of São Paulo (EPM – UNIFESP)

Outpatients Department – Geriatrics and Gerontology Discipline

Rua Prof. Francisco de Castro, 105

São Paulo

Brazil

04039-001

Study participating centre

Pró-Vida – Center for Total Health Assistance LLC

Av. Paes de Barros, 411 – cj. 14 e 15

São Paulo

Brazil

03115-020

Study participating centre

University of Santo Amaro – Medicine College

Outpatients Department – Rheumatology Discipline

Rua Cássio de Campos Nogueira, 2031

São Paulo

Brazil

04829-310

Sponsor information

Organisation

Sao Paulo Medical School (Escola Paulista de Medicina) (Brazil)

ROR

<https://ror.org/036rp1748>

Funder(s)

Funder type

University/education

Funder Name

Sao Paulo Medical School (Escola Paulista de Medicina) (Brazil)

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Carlos César Lopes de Jesus (caceloje@gmail.com).

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results		11/01/2017	12/01/2017	No	No
Participant information sheet		31/12/2016	04/01/2017	No	Yes