ISRCTN56637846 https://doi.org/10.1186/ISRCTN56637846

STRategies in Early Arthritis Management

Submission date	Recruitment status	[ ] Prospectively registered
20/12/2005	No longer recruiting	[ ] Protocol
Registration date	**Overall study status**	[ ] Statistical analysis plan
20/12/2005	Completed	[ ] Results
Last Edited	**Condition category**	[ ] Individual participant data
17/12/2008	Musculoskeletal Diseases	[ ] Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr D. van Schaardenburg

Contact details
Jan van Breemen Instituut
Dr. Jan van Breemenstraat 2
Amsterdam
Netherlands
1056 AB
+31 (0)20 5896589
d.v.schaardenburg@janvanbreemen.nl

Additional identifiers

Study information

Scientific Title

Acronym
STREAM

Study objectives

After 2 years of treatment with a combination of anti-rheumatic drugs including adalimumab with the aim of achieving and maintaining remission in patients with mild arthritis, there is less radiographic progression than in patients treated with usual care.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Received from the local medical ethics committee

Primary study design
Interventional

Study design
Randomised single blind active controlled parallel group trial

Study type(s)
Treatment

Health condition(s) or problem(s) studied
Polyarthritis/rheumatoid arthritis

Interventions
Methotrexate, in case of insufficient response followed by adalimumab, and then by a combination of methotrexate, sulfasalazine, hydroxychloroquine and prednisone depending on achievement of remission versus usual care according to preference physician.

Intervention Type
Drug

Phase
Not Specified

Drug/device/biological/vaccine name(s)
Adalimumab, methotrexate, sulfasalazine, hydroxychloroquine, prednisone

Primary outcome(s)
Progression of radiographic damage score after 2 years.

Key secondary outcome(s)
1. Functional capacity
2. Quality of life
3. Disease activity

Completion date
01/07/2007

Eligibility

Key inclusion criteria
1. Aged 18+ years
2. Symptom duration less than 3 years
3. Swelling in two to five joints

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group
Adult

Lower age limit
18 Years

Sex
All

Key exclusion criteria
1. Earlier treatment with disease modifying antirheumatic drugs except hydroychloroquine
2. Prednisone use within 3 months
3. Bacterial arthritis, crystal induced arthritis, reactive arthritis, sarcoidosis, osteoarthritis or systemic autoimmune disease other than rheumatoid arthritis (RA)
4. Pregnancy
5. Erosive disease

Date of first enrolment
01/07/2004

Date of final enrolment
01/07/2007

Locations

Countries of recruitment
Netherlands

Study participating centre
Jan van Breemen Instituut
Amsterdam
Netherlands
1056 AB

Sponsor information

Organisation
Jan van Breemen Institute (Netherlands)

ROR
https://ror.org/00bp9f906

Funder(s)

Funder type
Industry

Funder Name
Abbott (Netherlands) - study is investigator driven

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration