

Tolerability of the intravenous immunoglobulin Octagam® 10%

Submission date 26/11/2008	Recruitment status No longer recruiting	☐ Prospectively registered
		☐ Protocol
Registration date 03/12/2008	Overall study status Completed	☐ Statistical analysis plan
		☒ Results
Last Edited 14/11/2022	Condition category Haematological Disorders	☐ Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Anette Debes

Contact details
Elisabeth-Selbert-Str. 11
Langenfeld
Germany
40764
-
anette.debes@octapharma.de

Additional identifiers

Protocol serial number
GAM10-05

Study information

Scientific Title
Tolerability of the intravenous immunoglobulin Octagam® 10%

Study objectives

Octagam® 10% is well tolerated in routine clinical use in the treatment of primary or secondary immunodeficiencies or in the immunomodulation of autoimmune diseases.

Ethics approval required

Old ethics approval format

Ethics approval(s)

As the procedures during this observational study do not interfere with the patient's usual treatment and monitoring of treatment, this study is not regarded as a clinical study as defined by EU Directive 2001/20/EC. Therefore, approval by an Independent Ethical Committee is not required.

Primary study design

Observational

Study design

Non-interventional prospective multi-centre observational cohort study

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Primary or secondary immunodeficiencies, immunomodulation in autoimmune diseases

Interventions

Treatment with Octagam® 10% will be documented. This includes data about the patient's disease, age, gender, weight, concomitant medication or illness. For each application, the date and duration of infusion, dose, batch number(s) and the absence or occurrence of an adverse drug reaction (ADR) will be recorded. In case of an ADR, additional detailed information about the reaction will be recorded. If available, laboratory data about the efficacy of treatment should also be documented. No investigations must be initiated for the purpose of this non-interventional trial.

The number of treatments with Octagam® 10% for each patient cannot be defined due to the different indications where it is used. Therefore some patients will be treated and observed for a few weeks and others for several months or years. For this reason, there are no specified timepoints for the outcomes of this trial.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Octagam® 10%

Primary outcome(s)

To observe the tolerability of Octagam® 10% in different indication groups in routine clinical practice

Key secondary outcome(s)

Data about the efficacy of Octagam® 10%

Completion date

14/09/2013

Eligibility**Key inclusion criteria**

Patients of any age and gender who receive treatment with Octagam® 10%

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

All

Sex

All

Key exclusion criteria

Patients with known contraindications to Octagam® 10%

Date of first enrolment

15/09/2008

Date of final enrolment

14/09/2013

Locations**Countries of recruitment**

Germany

Study participating centre

Octapharma GmbH

Langenfeld

Germany

40764

Sponsor information

Organisation

Octapharma GmbH (Germany)

ROR

<https://ror.org/002k5fe57>

Funder(s)**Funder type**

Industry

Funder Name

Octapharma GmbH (Germany)

Results and Publications**Individual participant data (IPD) sharing plan**

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	post-authorisation safety analysis	01/05/2018		Yes	No
Results article		08/03/2018	14/11/2022	Yes	No