

# Onset of analgesia with OxyNorm Instant in healthy volunteers

<b>Submission date</b> 19/01/2014	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 27/01/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 27/01/2014	<b>Condition category</b> Signs and Symptoms	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Acute and breakthrough pain (sudden additional acute pain on top of the long-term pain) is difficult to treat, especially since the pain requires immediate relief. Various drugs are available with different timings of pain relief. Here we compare the effect of a new painkiller drug in the form of a melt tablet, OxyNorm instant (oxycodone formulation), with an active placebo (dummy) tablet, a paracetamol melt tablet, to assess the speed of pain relief.

### Who can participate?

The study will be performed in twelve healthy female volunteers, aged 18 to 65.

### What does the study involve?

Pain will be induced experimentally through electrical and pressure pain stimulus. Participants will be randomly allocated to one of two groups. For pain relief, OxyNorm instant will be given to one group and paracetamol to the other. Pain will be recorded over 5 hours following drug intake. Participants are then crossed over (i.e., the treatment is swapped). A mathematical model will be used to analyse the data and get an indication of onset of pain relief.

### What are the possible benefits and risks of participating?

There will be no benefit for the participants. However, it will be useful for treating patients appropriately in future. As the drug doses are low, the risks of the study are minimal and include vomiting and itch.

### Where is the study run from?

The study is run from the Leiden University Medical Center, Netherlands.

### When is the study starting and how long is it expected to run for?

The study started in January 2013 and will run until February 2014.

### Who is funding the study?

Mundipharma Pharmaceuticals BV, Netherlands.

Who is the main contact?

Prof. Albert Dahan

a.dahan@lumc.nl

## Contact information

### Type(s)

Scientific

### Contact name

Prof Albert Dahan

### Contact details

Albinusdreef 2

Leiden

Netherlands

2333 ZA

## Additional identifiers

### Clinical Trials Information System (CTIS)

2012-002227-15

### Protocol serial number

POXY/12-124

## Study information

### Scientific Title

A randomized cross-over study on the onset of analgesia with OxyNorm Instant in healthy volunteers

### Acronym

Oxy Study

### Study objectives

Oxycodone produces greater analgesia than the active comparator and meaningful analgesia, as defined by a 15% increase in response thresholds, within 10 min.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Medical Ethics Committee (Commissie Medische Ethiek), Leiden University Medical Center, Leiden, The Netherlands, 01/08/2012, Ref: P12.124

### Study design

Randomized placebo-controlled double-blind crossover trial

**Primary study design**

Interventional

**Study type(s)**

Quality of life

**Health condition(s) or problem(s) studied**

Breakthrough pain

**Interventions**

Patients are randomised to two groups: administration of either oxycodone 20 mg or paracetamol 500 mg. Participants are then crossed over and the treatment is swapped. Antinociceptive responses to electrical and pressure pain are measured.

**Intervention Type**

Other

**Phase**

Not Applicable

**Primary outcome(s)**

1. Efficacy is measured as a difference in pain threshold and pain tolerance scores of the OxyNorm Instant treated subjects as compared to paracetamol treated subjects at different time points
2. To determine the onset of analgesia of OxyNorm Instant. Onset of analgesia is defined as the time point at which the OxyNorm Instant treated patients display significantly increased levels of pain threshold and/or pain tolerance

**Key secondary outcome(s)**

Not provided at time of registration

**Completion date**

01/02/2014

**Eligibility****Key inclusion criteria**

1. Females
2. Age of 18 to 65 years (inclusive)
3. Body Mass Index (BMI) between 18 and 35 kg/m<sup>2</sup> (inclusive) and body weight between 50 kg and 100 kg (inclusive)
4. Subject is able to read and understand the written consent form, complete study-related procedures, and communicate with the study staff
5. Subject is willing to comply with study restrictions

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

Female

**Key exclusion criteria**

1. Clinically relevant abnormal history of physical and mental health, as determined by medical history taking and physical examinations obtained during the screening visit and/or prior to the administration of the initial dose of the study drug (as judged by the investigator)
2. A semi-recumbent systolic blood pressure of >150 mmHg and/or diastolic blood pressure of >90 mmHg at screening
3. History of alcoholism or substance abuse within three years prior to screening
4. Positive pregnancy test
5. Subjects using more than 14 units of alcohol per week
6. Use of medication during the study period
8. Subject is not using oral contraceptives or is not post-menopausal (last menstrual period > 2 years ago and FSH > 25 IU/L) or surgically sterilized
9. Subject has a history of severe allergies, or has had an anaphylactic reaction or significant intolerance to prescription or non-prescription drugs or food
10. Participation in an investigational drug trial in the 3 months prior to administration of the initial dose of study drug or more than 4 times per year
11. Any other condition that in the opinion of the investigator would complicate or compromise the study, or the well being of the subject:  
OxyNorm is contra-indicated in case of hypersensitivity for oxycodone or one of its excipients or in any situation where opioids are contra-indicated. This can include the following situations:
  - 11.1. Respiratory depression
  - 11.2. Head injury
  - 11.3. Paralytic ileus
  - 11.4. Acute abdomen
  - 11.5. Chronic constipation
  - 11.6. Severe obstructive airways disease
  - 11.7. Severe bronchial asthma
  - 11.8. Cor pulmonale
  - 11.9. Hypercarbia
  - 11.10 Acute hepatic disease
  - 11.11. Severe hepatic impairment
  - 11.12. Severe renal impairment (creatinine clearance <10 ml/min)
  - 11.13. Cyanosis
  - 11.14. Concurrent administration of monoamine oxidase inhibitors or within 2 weeks of discontinuation of their use

**Date of first enrolment**

01/01/2013

**Date of final enrolment**

01/02/2014

# Locations

## Countries of recruitment

Netherlands

## Study participating centre

Albinusdreef 2

Leiden

Netherlands

2333 ZA

# Sponsor information

## Organisation

Leiden University Medical Center (Netherlands)

## ROR

<https://ror.org/05xvt9f17>

# Funder(s)

## Funder type

Other

## Funder Name

Investigator initiated and funded (Netherlands)

## Funder Name

Partly by Mundipharma Pharmaceuticals BV (Netherlands)

# Results and Publications

## Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration