

# Impact of treatment of chronic periodontitis on the serum levels of prohepcidin in patients with chronic kidney disease

<b>Submission date</b> 07/11/2010	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 18/11/2010	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 03/01/2012	<b>Condition category</b> Oral Health	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Eduardo Vilela

**Contact details**  
Rua Dr Geraldo Moutinho 55  
Jardins Imperiais  
Juiz de Fora  
Brazil  
36036348  
eduardo.vilela@ufjf.edu.br

## Additional identifiers

**Protocol serial number**  
# 942.248.2006/report # 327/2006

## Study information

**Scientific Title**  
Impact of treatment of chronic periodontitis on the serum levels of prohepcidin in patients with chronic kidney disease: Interventional controlled clinical assay

**Acronym**

UFJF

**Study objectives**

We hypothesized that part of the chronic inflammatory response seen in chronic kidney disease (CKD) patients stems from chronic periodontitis (CP), which, through the increase in the expression of inflammatory markers such as interleukin-6 (IL-6), stimulates hepcidin synthesis.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Ethics Committee in Research on Human Beings from UFJF approved on the 6th of December 2006 (ref: 942.248.2006 / report: 327/2006)

**Study design**

Interventional controlled clinical assay

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Chronic periodontitis (CP); chronic kidney disease (CKD); Prohepcidin

**Interventions**

Patients with CP were divided into two groups: CKD patients and controls  
Both groups received instructions on oral hygiene techniques, including the manual techniques of tooth and interproximal brushing, and on how to use dental floss and perform supragingival prophylaxis. The nonsurgical periodontal therapy (PT) consisted of radicular scraping and subgingival curettage, using standard instrumentation with Gracey curettes and ultrasound devices performed in 1 hour sessions over an average period of 4 weeks. Local anesthesia was used when necessary.

Upon concluding the periodontal treatment, participants were followed up after 15, 30, 60, and 90 days. At each return visit, instructions on oral hygiene and supragingival phrophylaxis were provided.

Blood samples were collected for biochemical analysis at baseline and 3 months after PT. Venous blood after 12 hours fasting was collected in vacuum tubes between 7:00am and 9:00am. Plasma samples with EDTA/heparin and serum samples were immediately placed in ice, aliquoted within 1 hour and stored at -80C until use.

**Results:**

The efficacy of PT was proven by the statistically significant decrease in the levels of inflammatory markers and the improvement of clinical parameters of CP observed 3 months after completion of the PT. Prohepcidin, IL-6 and us-PCR levels diminished significantly after PT

in both groups. In the control group, besides the decrease in the inflammatory markers, a significant increase could also be observed in the levels of haemoglobin and ferritin associated with PT.

### **Intervention Type**

Other

### **Phase**

Not Applicable

### **Primary outcome(s)**

Inflammatory markers, assessed at baseline and 3 months post-PT

1. C-reactive protein (us-CRP)
2. Interleukin-6 (IL-6)
3. Prohepcidin

### **Key secondary outcome(s)**

Biochemical analyses from blood sample in EDTA, at baseline and 3 months post-PT

1. Complete haemogram (automated Coulter STKS)
2. Serum iron (ferrozine)
3. Ferritin (electrochemiluminescence)
4. Transferrine saturation index (labtest ferrozine)

### **Completion date**

10/08/2010

## **Eligibility**

### **Key inclusion criteria**

1. Patients with CP, allocated to one of two groups depending on CKD status
  - 1.1. CKD group:
    - 1.1.1. Patients with CKD at stages 3 to 5 and undergoing conservative treatment
    - 1.1.2. Recruited from the PREVENRIM, a CKD prevention clinic at the Interdisciplinary Nucleus of Studies, Research and Treatment in Nephrology (NIEPEN) of the Universidade Federal de Juiz de Fora (UFJF)
  - 1.2. Control patients:
    - 1.2.1. Patients with no systemic disease from the Periodontology Clinic of the School of Dentistry at UFJF
2. Over 18 years of age, either sex
2. Minimum of 20 natural teeth and without periapical lesions
3. Have received no periodontal, antimicrobial, or anti-inflammatory treatment within the last 6 months
4. Have not used steroids or immunosuppressant drugs within the last 6 months

### **Participant type(s)**

Patient

### **Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Key exclusion criteria**

1. Pregnant or breast feeding women
2. Smokers or ex-smokers who had quit smoking within the last 10 years

**Date of first enrolment**

10/08/2008

**Date of final enrolment**

10/08/2010

**Locations****Countries of recruitment**

Brazil

**Study participating centre**

Rua Dr Geraldo Moutinho 55

Juiz de Fora

Brazil

36036348

**Sponsor information****Organisation**

Federal University of Juiz de Fora (Universidade Federal de Juiz de Fora [UFJF]) (Brazil)

**ROR**

<https://ror.org/04yqw9c44>

**Funder(s)****Funder type**

University/education

## Funder Name

Federal University of Juiz de Fora (Universidade Federal de Juiz de Fora [UFJF]) (Brazil)

# Results and Publications

## Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/07/2011		Yes	No